

DIPARTIMENTO DI MATEMATICA
“Francesco Brioschi”
POLITECNICO DI MILANO

**Quantitative Local Bounds for
Subcritical Semilinear Elliptic
Equations**

Bonforte, M.; Grillo, G.; Vazquez, J.L.

Collezione dei *Quaderni di Dipartimento*, numero **QDD 122**
Inserito negli *Archivi Digitali di Dipartimento* in data 10-5-2012



Piazza Leonardo da Vinci, 32 - 20133 Milano (Italy)

Quantitative Local Bounds for Subcritical Semilinear Elliptic Equations

Matteo Bonforte, Gabriele Grillo and Juan Luis Vazquez

Abstract. The purpose of this paper is to prove local upper and lower bounds for weak solutions of semilinear elliptic equations of the form $-\Delta u = cu^p$, with $0 < p < p_s = (d+2)/(d-2)$, defined on bounded domains of \mathbb{R}^d , $d \geq 3$, without reference to the boundary behaviour. We give an explicit expression for all the involved constants. As a consequence, we obtain local Harnack inequalities with explicit constants, as well as gradient bounds.

Mathematics Subject Classification (2010). 35B45, 35B65, 35K55, 35K65.

Keywords. Local bounds, semilinear elliptic equations, regularity, Harnack inequality.

Contents

| | |
|--|----|
| 1. Introduction | 2 |
| 2. Preliminaries. Local energy estimate | 3 |
| 2.1. More general nonlinearities | 6 |
| 3. Local Upper Bounds | 7 |
| 3.1. Local upper bounds I. The upper Moser iteration | 7 |
| 3.2. Local upper bounds II. Linear case with unbounded coefficients | 17 |
| 3.2.1. Energy Estimates and Reverse Poincaré inequalities | 17 |
| 3.2.2. Extending local upper bounds. A lemma by De Giorgi | 25 |
| 4. Lower bounds | 29 |
| 4.1. A short reminder about the spaces $M^p(\Omega)$. | 29 |
| 4.2. The John-Nirenberg Lemma and reverse Hölder inequalities. | 31 |
| 4.3. Lower Moser iteration | 34 |
| 4.4. Reverse Hölder inequalities and lower bounds when $1 < p < p_c$ | 37 |
| 5. Harnack inequalities | 41 |
| 6. Local Absolute bounds | 45 |
| 7. Regularity. Local bounds for the gradients | 46 |
| 8. Table of results | 52 |
| References | 53 |

1. Introduction

In this paper we obtain local upper and lower estimates for the weak solutions of semilinear elliptic equations of the form

$$-\Delta u = f(u) \tag{1.1}$$

posed in a bounded domain $\Omega \subset \mathbb{R}^d$. The choice of right-hand side we have in mind is $f(u) = \lambda u^p$ with $\lambda, p > 0$. The range of exponents of interest will be $1 < p < p_s := (d+2)/(d-2)$ if $d \geq 3$, or $p > 1$ if $d = 1, 2$. This problem is one of the most popular problems in nonlinear elliptic theory and enjoys a large bibliography [1, 7, 8, 12, 13, 14, 18, 19, 20, 21, 22, 23, 24, 26, 27, 32, 33, 34, 35, 36, 37] for different p , and [6, 10] for the limit case $p = p_s$.¹

We focus our attention on obtaining local estimates for solutions that are defined inside the domain without reference to their boundary behaviour. This is the notion of solution we use.

Definition 1.1. *A local weak solution to equation $-\Delta u = f(u)$ in Ω is defined as a function $u \in W_{loc}^{1,2}(\Omega)$ with $f(u) \in L_{loc}^1(\Omega)$ which satisfies*

$$\int_K [\nabla u \cdot \nabla \varphi - f(u)\varphi] \, dx = 0 \tag{1.2}$$

for any subdomain with compact closure $K \subset \Omega$ and all bounded $\varphi \in C_0^1(K)$.

Our aim is to contribute quantitative estimates in the form of upper bounds for solutions of any sign, lower bounds for positive solutions, and also local Harnack inequalities and gradient bounds. By quantitative estimates we mean keeping track of all the constants during the proofs. As far as we know, there does not exist in literature a systematic set of quantitative estimates of local upper and lower bounds, nor of the Harnack constant, in the explicit form we provide here. We recall that the quantitative control of the constants of such inequalities may have an important role in the applications; it is needed for instance in the results of [2] on the asymptotic properties of solutions of the fast diffusion equation in bounded domains.

Contents and main results. We start with a section devoted to basic energy estimates. We then consider in Section 3 the upper estimates for nonnegative solutions of the equation $-\Delta u = \lambda u^p$. The exponent range is $0 \leq p < p_s$; this is a main restriction of the theory, as it is already well known. See also [9] for L^∞ -bounds of different type for Equation (1.1) with more general nonlinearities.

Our first main result, Theorem 3.1, can be considered as a smoothing effect with very precise constants; it is much simpler for $p \leq 1$, but we also obtain the more complicated and novel estimates for $1 < p < p_s$. Next, we obtain local upper estimates for $-\Delta u = b(x)u$ with unbounded coefficient b in Theorem 3.8 and we apply them to the case $b(x) = u^{p-1}$ in Theorem 3.9.

¹We refrain from attempting to give a complete bibliography for this nowadays classical problem.

In Section 4 we prove quantitative lower estimates, Theorems 4.6, 4.8. We prove Harnack inequalities in Theorems 5.1, 5.2 and 5.3. All of these results appear to be well known from a qualitative point of view. Let us mention that, as far as we know, the Harnack inequality for solutions to (1.2) when $p > 1$ is not stated explicitly in the literature. The fact that the “constant” involved has to depend on u when $p_c \leq p < p_s$ is confirmed by the results of [4], [16] applied to separation of variable solutions of parabolic problems, see also the very recent monograph [17]. This is also related to the fact that, in the range $p_c \leq p < p_s$, there exist (very weak) singular solutions. Notice also that in such a range the notion of weak and very weak solution is really different, cf. [15, 25, 28, 29, 30, 31].

In Section 6 we derive quantitative absolute upper (for $1 < p < p_c$) and lower bounds (for $0 \leq p < 1$) which are new as far as we know, at least from a quantitative point of view, cf. Theorem 6.1. Universal (or absolute) upper bounds for weak solutions defined in an open subset of \mathbb{R}^d in the whole range $(1, p_s)$ follow as a consequence of the works of Dancer [11, 12] on classical solutions, and from the fact that weak solutions are indeed classical as can be proved by a standard bootstrap argument. Weak solutions are classical also when $p = p_s$, as proved by Brezis and Kato [5]. We do not obtain quantitative versions of these absolute upper bounds in the intermediate range $[p_c, p_s)$ with our methods.

The last section is devoted to quantitative gradient estimates, cf. Theorem 7.2, and absolute upper bounds for the gradient when $1 < p < p_c$, cf. Theorem 7.3.

Much of the known theory takes into account boundary conditions of different types: Dirichlet, Neumann, Robin, or other. Our results apply to all those cases. We will study the precise estimates for the Dirichlet problem in an upcoming paper [3].

Finally, the authors are grateful to A. Farina for relevant information on the topic of absolute bounds.

2. Preliminaries. Local energy estimate

We shall pursue in the sequel the well-known idea that local weak solutions satisfy reverse Sobolev or Poincaré inequalities. Such local reverse inequalities are the key to prove local upper and lower estimates in the next sections, and indeed they imply that such functions are Hölder continuous.

Lemma 2.1 (Energy Estimates). *Let $\Omega \subset \mathbb{R}^d$ be a bounded domain, and let $p \geq 0$ and $\lambda > 0$. Let u be a local nonnegative weak solution in Ω to $-\Delta u = \lambda u^p$. Then the following energy equality holds true for any $\delta > 0$, $\alpha \neq -1$ and any positive test function $\varphi \in C^2(\Omega)$ that is compactly supported in Ω :*

$$4\alpha \int_{\Omega} |\nabla \left((u + \delta)^{\frac{\alpha+1}{2}} \right)|^2 \varphi \, dx = \lambda(\alpha + 1)^2 \int_{\Omega} u^p (u + \delta)^\alpha \varphi \, dx + (\alpha + 1) \int_{\Omega} (u + \delta)^{\alpha+1} \Delta \varphi \, dx. \quad (2.1)$$

Moreover, for any $\delta \geq 0$ we have the Caccioppoli estimates

$$\lambda \int_{\Omega} \frac{u^p}{u + \delta} \varphi \, dx + \int_{\Omega} |\nabla \log(u + \delta)|^2 \varphi \, dx \leq \int_{\Omega} \frac{|\nabla \varphi|^2}{\varphi} \, dx. \quad (2.2)$$

Local subsolutions \underline{u} of $-\Delta \underline{u} \leq \lambda \underline{u}^p$ satisfy, for $\alpha \neq -1$ and $\delta > 0$:

$$\begin{aligned} 4\alpha \int_{\Omega} \left| \nabla \left((\underline{u} + \delta)^{\frac{\alpha+1}{2}} \right) \right|^2 \varphi \, dx &\leq \lambda(\alpha + 1)^2 \int_{\Omega} \underline{u}^p (\underline{u} + \delta)^{\alpha} \varphi \, dx \\ &+ (\alpha + 1) \int_{\Omega} (\underline{u} + \delta)^{\alpha+1} \Delta \varphi \, dx, \end{aligned} \quad (2.3)$$

while local supersolution $-\Delta \bar{u} \geq \lambda \bar{u}^p$ satisfy, for any $\alpha \neq -1$ and $\delta > 0$:

$$\begin{aligned} \frac{4\alpha}{(\alpha + 1)^2} \int_{\Omega} \left| \nabla \left((\bar{u} + \delta)^{\frac{\alpha+1}{2}} \right) \right|^2 \varphi \, dx &\geq \lambda \int_{\Omega} \bar{u}^p (\bar{u} + \delta)^{\alpha} \varphi \, dx \\ &+ \frac{1}{\alpha + 1} \int_{\Omega} (\bar{u} + \delta)^{\alpha+1} \Delta \varphi \, dx, \end{aligned} \quad (2.4)$$

and the Caccioppoli (2.2) estimates also work.

Remark. Notice that when $\alpha > -1$, we can let $\delta = 0$ in the energy identity (2.1) to get

$$4\alpha \int_{\Omega} \left| \nabla \left(u^{\frac{\alpha+1}{2}} \right) \right|^2 \varphi \, dx = \lambda(\alpha + 1)^2 \int_{\Omega} u^{p+\alpha} \varphi \, dx + (\alpha + 1) \int_{\Omega} u^{\alpha+1} \Delta \varphi \, dx. \quad (2.5)$$

The same remark applies to subsolutions:

$$4\alpha \int_{\Omega} \left| \nabla \left(\underline{u}^{\frac{\alpha+1}{2}} \right) \right|^2 \varphi \, dx \leq \lambda(\alpha + 1)^2 \int_{\Omega} \underline{u}^{p+\alpha} \varphi \, dx + (\alpha + 1) \int_{\Omega} \underline{u}^{\alpha+1} \Delta \varphi \, dx \quad (2.6)$$

Proof. Let $\varphi \in C^2(\Omega) \cap C_0^1(\bar{\Omega})$ and $\delta \geq 0$. Multiply $-\Delta u$ by $(u + \delta)^{\alpha} \varphi$, with $\alpha \neq -1$ and integrate by parts to get

$$\begin{aligned} - \int_{\Omega} \varphi (u + \delta)^{\alpha} \Delta u \, dx &= \int_{\Omega} \nabla \varphi \cdot (\nabla u) (u + \delta)^{\alpha} \, dx \\ &+ \alpha \int_{\Omega} \varphi (u + \delta)^{\alpha-1} |\nabla u|^2 \, dx \\ &= - \frac{1}{\alpha + 1} \int_{\Omega} (u + \delta)^{\alpha+1} \Delta \varphi \, dx \\ &+ \frac{4\alpha}{(\alpha + 1)^2} \int_{\Omega} \left| \nabla (u + \delta)^{\frac{\alpha+1}{2}} \right|^2 \varphi \, dx. \end{aligned} \quad (2.7)$$

For local weak solutions of $-\Delta u = \lambda u^p$, the above equality immediately gives the energy identity (2.1) for $\alpha \neq -1$. Similar considerations hold, in the stated range of α , for sub and supersolutions. To derive the Caccioppoli

estimate we use the test function $\varphi/(u + \delta)$ to get

$$\begin{aligned}
0 &\leq \lambda \int_{\Omega} \frac{u^p}{u + \delta} \varphi \, dx = - \int_{\Omega} \frac{\varphi}{u + \delta} \Delta u \, dx \\
&= - \int_{\Omega} \frac{\varphi}{(u + \delta)^2} |\nabla u|^2 \, dx + \int_{\Omega} \frac{\nabla \varphi \cdot \nabla u}{u + \delta} \frac{\sqrt{\varphi}}{\sqrt{\varphi}} \, dx \\
&\leq - \int_{\Omega} \varphi |\nabla \log(u + \delta)|^2 \, dx + \frac{1}{2} \int_{\Omega} \frac{|\nabla \varphi|^2}{\varphi} \, dx + \frac{1}{2} \int_{\Omega} |\nabla \log(u + \delta)|^2 \varphi \, dx \\
&\leq - \frac{1}{2} \int_{\Omega} \varphi |\nabla \log(u + \delta)|^2 \, dx + \frac{1}{2} \int_{\Omega} \frac{|\nabla \varphi|^2}{\varphi} \, dx,
\end{aligned}$$

where we have used the inequality $a \cdot b \leq (|a|^2 + |b|^2)/2$. \square

We shall also need the following particular computation.

Lemma 2.2. *Fix two balls $B_{R_1} \subset B_{R_0} \subset\subset \Omega$. Then there exists a test function $\varphi \in C_0^1(B_{R_0})$, with $\nabla \varphi \equiv 0$ on $\partial\Omega$, which is radially symmetric and piecewise C^2 as a function of r , satisfies $\text{supp}(\varphi) = B_{R_0}$ and $\varphi = 1$ on B_{R_1} , and moreover satisfies the bounds*

$$\|\nabla \varphi\|_{\infty} \leq \frac{4}{R_0 - R_1} \quad \text{and} \quad \|\Delta \varphi\|_{\infty} \leq \frac{4d}{(R_0 - R_1)^2}. \quad (2.8)$$

Proof. Consider the radial test function defined on B_{R_0}

$$\varphi(|x|) = \begin{cases} 1 & \text{if } 0 \leq |x| \leq R_1 \\ 1 - \frac{2(|x| - R_1)^2}{(R_0 - R_1)^2} & \text{if } R_1 < |x| \leq \frac{R_0 + R_1}{2} \\ \frac{2(R_0 - |x|)^2}{(R_0 - R_1)^2} & \text{if } \frac{R_0 + R_1}{2} < |x| \leq R_0 \\ 0 & \text{if } |x| > R_0 \end{cases} \quad (2.9)$$

for any $0 < R_1 < R_0$. We have

$$\nabla \varphi(|x|) = \begin{cases} 0 & \text{if } 0 \leq |x| \leq R_1 \text{ or if } |x| > R_0 \\ -\frac{4(|x| - R_1)}{(R_0 - R_1)^2} \frac{x}{|x|} & \text{if } R_1 < |x| \leq \frac{R_0 + R_1}{2} \\ -\frac{4(R_0 - |x|)}{(R_0 - R_1)^2} \frac{x}{|x|} & \text{if } \frac{R_0 + R_1}{2} < |x| \leq R_0 \end{cases}$$

and, recalling that $\Delta \varphi(|x|) = \varphi''(|x|) + (d-1)\varphi'(|x|)/|x|$,

$$\Delta \varphi(|x|) = \begin{cases} 0 & \text{if } 0 \leq |x| \leq R_1 \text{ or if } |x| > R_0 \\ -\frac{4}{(R_0 - R_1)^2} - \frac{d-1}{|x|} \frac{4(|x| - R_1)}{(R_0 - R_1)^2} & \text{if } R_1 < |x| \leq \frac{R_0 + R_1}{2} \\ -\frac{4}{(R_0 - R_1)^2} - \frac{d-1}{|x|} \frac{4(R_0 - |x|)}{(R_0 - R_1)^2} & \text{if } \frac{R_0 + R_1}{2} < |x| \leq R_0 \end{cases}$$

As a consequence we easily obtain the bounds (2.8). \square

Corollary 2.3 (Quantitative Caccioppoli Estimates). *Let $\delta \geq 0$. Let $\Omega \subset \mathbb{R}^d$ be a bounded domain, and let $p \geq 0$ and $\lambda > 0$. Let u be a local positive weak solution in Ω to $-\Delta u = \lambda u^p$. For any $B_R \subset B_{R_0} \subset \subset \Omega$ we have*

$$\lambda \int_{B_R} \frac{u^p}{u + \delta} dx + \int_{B_R} |\nabla \log(u + \delta)|^2 dx \leq \frac{8\omega_d R_0^d}{(R_0 - R)^2} \quad (2.10)$$

where ω_d denotes the volume of the unit ball in \mathbb{R}^d .

Proof. We use (2.2), using the test function φ of Lemma 2.2 with R replacing R_1 :

$$\begin{aligned} & \lambda \int_{B_R} \frac{u^p}{u + \delta} dx + \int_{B_R} |\nabla \log(u + \delta)|^2 dx \\ & \leq \lambda \int_{\Omega} \frac{u^p}{u + \delta} \varphi dx + \int_{\Omega} \varphi |\nabla \log(u + \delta)|^2 dx \\ & \leq \int_{\Omega} \frac{|\nabla \varphi|^2}{\varphi} dx \leq \frac{8|\text{supp}(\varphi)|}{(R_0 - R)^2} = \frac{8\omega_d R_0^d}{(R_0 - R)^2}. \quad \square \end{aligned}$$

Note that the case $\delta > 0$ follows immediately from the case $\delta = 0$ since $u \geq 0$.

Remark. Letting $\delta = 0$ in the Caccioppoli estimates (2.10) shows that

$$\lambda \int_{B_R} u^{p-1} dx \leq \frac{8\omega_d R_0^d}{(R_0 - R)^2} \quad (2.11)$$

When $p > 1$ this yields a *local absolute upper bound* for the local L^{p-1} -norm, a fact that will allow to conclude an absolute local L^∞ -bound in the range $1 < p < p_c := d/(d-2)$, as we shall see in Section 6. This absolute upper bound represents a novelty both because it is quantitative and because it is local: to our knowledge this is the first absolute local bound for elliptic equations. When $p = 1$ such absolute bound is easily seen to be impossible, while in the case $0 < p < 1$ we get an absolute lower bound for the local L^{p-1} -integral, which is new, at least as far as we know. It will be used below.

2.1. More general nonlinearities

As long as we deal with local estimates, we can apply the method to a larger class of operators and nonlinearities. (i) First of all, namely we can treat local solutions of:

$$-\nabla \cdot A(x, u, \nabla u) = \lambda u^p, \quad (2.12)$$

where A is a Carathéodory function such that

$$\nu_1 |\xi|^2 \leq A(x, u, \xi) \cdot \xi \leq \nu_2 |\xi|^2 \quad \text{and} \quad |A(x, u, \xi)|^2 \leq \nu_2 |\xi|^2$$

for suitable constants $0 < \nu_1 < \nu_2$. The proofs of the inequalities are the same, and the results contain ν_1 (resp. ν_2) depending on whether you consider subsolutions (resp. supersolutions).

(ii) Second we can consider supersolutions of the problem

$$-\nabla \cdot A(x, u, \nabla u) = f(x, u), \quad (2.13)$$

as long as $f(u) \geq a_0 u^p$ with $a_0 > 0$, since they are supersolutions of $-\nabla \cdot A(x, u, \nabla u) = a_0 u^p$.

(iii) We can consider subsolutions of (2.13) with $f(u) \leq a_1(u + b_1)^p$, and $a_1, b_1 \geq 0$. Then we can obtain an estimate for $v = u + b_1$.

The only thing that changes a bit are the energy estimates, and it is not so difficult to keep track of the new constants throughout the proof. We have decided here to consider the model case, to simplify the presentation and to focus on the main ideas.

(iv) Other semilinear problems of this type are treated in the literature. Thus, Ambrosetti and Prodi's book [1] discusses right-hand sides of the form $f(x, u) = \lambda u + c(u) + h(x)$, with $a \in \mathbb{R}$, $c(\cdot) \in C^0(\mathbb{R}) \cap L^\infty(\mathbb{R})$ and $h \in C^{0,\alpha}(\bar{\Omega})$, for some $\alpha \in (0, 1)$. Such nonlinearities can be treated with the methods presented here as well. We refrain from dealing with it in this work.

3. Local Upper Bounds

This section is devoted to the proof of the upper bounds and we will provide two kinds of estimates. We prove local upper bounds for nonnegative subsolutions, then by Kato's inequality it is easy to extend such results to solutions with any sign.

3.1. Local upper bounds I. The upper Moser iteration

The local upper bounds follow from the local Sobolev imbedding theorem on balls $B_R \subset \mathbb{R}^d$

$$\|f\|_{L^{2^*}(B_R)}^2 \leq \mathcal{S}_2^2 \left(\|\nabla f\|_{L^2(B_R)}^2 + \frac{1}{R^2} \|f\|_{L^2(B_R)}^2 \right) \quad (3.1)$$

where $\mathcal{S}_2 = \mathcal{S}_2(B_1)$ is the best constant and $2^* = 2d/(d-2)$. We are requiring hereafter without any further comment that $d \geq 3$. The Sobolev inequality combines with the energy inequalities of Lemma 2.1 which can be considered as local reverse Sobolev (or Poincaré) inequalities. The proof of the local upper bounds goes through the celebrated Moser iteration. We adopt the notation $\|f\|_{L^q(B_R)} = \|f\|_{q,R}$, we recall that $|B_R| = \omega_d R^d$ and that $\int_X f(x) dx = \int_X f(x) dx/|X|$. Throughout this section we are considering nonnegative subsolutions u to $-\Delta u = \lambda u^p$, unless otherwise explicitly stated.

Theorem 3.1 (Local Upper Estimates). *Let $\Omega \subseteq \mathbb{R}^d$ and let $\lambda > 0$. (i) Let $u \geq 0$ be a local weak subsolution to $-\Delta u = \lambda u^p$ in Ω , with $1 < p < p_s = 2^* - 1 = (d+2)/(d-2)$. Then, for any $q > \bar{q} := d(p-1)_+/2$ and for any $B_{R_\infty} \subset B_{R_0} \subseteq \Omega$, the following bound holds true*


$$\|u\|_{L^\infty(B_{R_\infty})} \leq I_{\infty,q} \left(\int_{B_{R_0}} u^q dx \right)^{\frac{1+(p-1)\mu}{q}} \left(\int_{B_{R_\infty}} u^{p-1} dx \right)^{-\mu} \quad (3.2)$$

where $\mu = d/(2q - d(p-1)) = d/2(q - \bar{q})$, and the constant $I_{\infty,q} > 0$ depends on d, p, q, R_0, R_∞ , but not on λ .

(ii) For $0 \leq p \leq 1$ the estimate simplifies into

$$\|u\|_{L^\infty(B_{R_\infty})} \leq I_{\infty,q} \left(\int_{B_{R_0}} u^q dx \right)^{1/q}. \quad (3.3)$$

valid for all $q > 0$. $I_{\infty,q} > 0$ has the same dependence as before, and it also depends on λ when $p = 1$, but not otherwise.



Exponents of the local upper estimates.

Remarks on the result. (i) Inequality (3.2) is a kind of reverse Hölder inequality, indeed we can rewrite it as:

$$\|u\|_{L^{\mu(p-1)}(B_{R_\infty})} \|u\|_{L^\infty(B_{R_\infty})} \leq C \|u\|_{L^q(B_{R_0})}^{1+\mu(p-1)}. \quad (3.4)$$

Written in this form, it is clear from Hölder's inequality that a constant which makes (3.4) true for a $q > \bar{q}$, make the same inequality true also for all $q' > q$. The same applies to (3.3).

(ii) The linear case $p = 1$ is well known, cf. [18, 22, 23].

Remarks on the constant. (i) The proof below allows to find the following expression for the constant:

$$I_{\infty,q} = \left[\frac{c_1 \mathcal{S}_2^2 \omega_d^{\frac{2(p-1)_+}{d(p-1)}}}{(1-\rho)^2} \right]^{\frac{d}{2q-d(p-1)_+}} \left\{ \left(\frac{d}{d-2} \right)^d \frac{2(d-2)}{(\sqrt{d}-\sqrt{d-2})^2} \right. \\ \left. \times \left[\Lambda_p + \frac{d-2}{q} + (1-\rho)^2 \max \left\{ \frac{d-2}{(dq)^2} |dq - (d-2)|, \frac{1}{4} \right\} \right] \right\}^{\frac{d}{2q-d(p-1)_+}} \quad (3.5)$$

where $\rho = R_\infty/R_0 < 1$ and we have used the convention $x_+/x = 0$ when $x = 0$ and, moreover, we have set $\Lambda_p = 2$ if $p \neq 1$, $\Lambda_p = \lambda/4$ if $p = 1$, with

$$c_1 := \begin{cases} \frac{(d-2)q}{(d-2)q-d} & \text{if } q > \frac{d}{d-2} \\ \max_{i=0,1} \frac{\left(\frac{d}{d-2}\right)^{k_0-1+i} \left[q - \frac{d(p-1)_+}{2} \right] + (p-1)_+ \frac{d-2}{2}}{\left| \left(\frac{d}{d-2}\right)^{k_0-1+i} \left[q - \frac{d(p-1)_+}{2} \right] + (p-1)_+ \frac{d-2}{2} - 1 \right|} & \text{if } 0 < q < \frac{d}{d-2}. \end{cases} \quad (3.6)$$

(iii) When q also satisfies $0 < q < d/(d-2)$, we will require in the proof the additional condition

$$\frac{\log \frac{2^* - d(p-1)_+}{2q - d(p-1)_+}}{\log \frac{d}{d-2}} \text{ is not an integer, and } k_0 := i.p. \left[\frac{\log \frac{2^* - d(p-1)_+}{2q - d(p-1)_+}}{\log \frac{d}{d-2}} \right], \quad (3.7)$$

(*i.p.* is the integer part of a real number). Notice that taking $q = p + 1 > d(p-1)/2$ is possible if and only if $p < p_s = (d+2)/(d-2)$.

(iv) Of course, condition (3.7) is not essential, in view of the remark after formula (3.4). In fact, let $q > \frac{d(p-1)_+}{2}$ be such that that $A(q) := \frac{\log \frac{2^* - d(p-1)_+}{2q - d(p-1)_+}}{\log \frac{d}{d-2}}$ is an integer. Take $\hat{q} \in (d(p-1)_+/2, q)$ such that $A(\hat{q})$ is not an integer. Then (3.2) is valid with \hat{q} instead of q .

Proof. We are going to use the energy identity (2.1) for any $\alpha > -1$, $\alpha \neq 0$, in the form (2.3) valid for subsolution, to prove $L^q - L^\infty$ local estimates via Moser iteration, keeping track of all the constants. We divide the proof in several steps.

• **STEP 1.** Let u as in Lemma 2.1 and φ the test function of Lemma 2.2, which is supported in B_{R_0} and such that $\varphi \equiv 1$ on B_{R_1} . The local Sobolev inequality (3.1) on the ball B_{R_1} applied to $f = u^{(\alpha+1)/2}$, together with the energy inequality (2.3) (we can take $\delta = 0$ as in (2.6)), gives

$$\begin{aligned} & \left[\int_{B_{R_1}} u^{\frac{2^*}{2}(\alpha+1)} dx \right]^{\frac{2}{2^*}} \leq S_2^2 \left(\int_{B_{R_1}} |\nabla u^{\frac{\alpha+1}{2}}|^2 dx + \frac{1}{R_1^2} \int_{B_{R_1}} u^{\alpha+1} dx \right) \\ & \leq S_2^2 \left(\int_{B_{R_0}} |\nabla u^{\frac{\alpha+1}{2}}|^2 \varphi dx + \frac{1}{R_1^2} \int_{B_{R_1}} u^{\alpha+1} dx \right) \\ & = S_2^2 \left(\frac{\lambda(\alpha+1)^2}{4|\alpha|} \int_{B_{R_0}} u^{p+\alpha} \varphi dx + \frac{\alpha+1}{4|\alpha|} \int_{B_{R_0}} u^{\alpha+1} \Delta \varphi dx \right. \\ & \quad \left. + \frac{1}{R_1^2} \int_{B_{R_1}} u^{\alpha+1} dx \right) \\ & \leq S_2^2 \left(\frac{\lambda(\alpha+1)^2}{4|\alpha|} \int_{B_{R_0}} u^{p+\alpha} dx + \left[\frac{(\alpha+1)\|\Delta \varphi\|_\infty}{4|\alpha|} + \frac{1}{R_1^2} \right] \int_{B_{R_0}} u^{\alpha+1} dx \right) \end{aligned}$$

$$\leq \mathcal{S}_2^2 \left(\frac{\lambda(\alpha+1)^2}{4|\alpha|} \int_{B_{R_0}} u^{p+\alpha} dx + \left[\frac{d(\alpha+1)}{|\alpha|(R_0-R_1)^2} + \frac{1}{R_1^2} \right] \int_{B_{R_0}} u^{\alpha+1} dx \right) \quad (3.8)$$

in the last step we have used the inequality $\|\Delta\varphi\|_\infty \leq 4d/(R_0-R_1)^2$ of Lemma 2.2.

• **STEP 2. Caccioppoli estimates and the first iteration step.** Now we need to split two cases, namely $0 \leq p \leq 1$ and $1 < p < p_s$, and in both cases we will use the Caccioppoli estimate (2.10) with $\delta = 0$ which holds for any $p > 0$ and reads

$$\lambda \frac{\|u\|_{p-1, R_\infty}^{p-1}}{|B_{R_0}|} \leq \frac{8}{(R_0-R_\infty)^2}. \quad (3.9)$$

Superlinear case: $1 < p < p_s$. We continue estimate (3.8) as follows:

$$\begin{aligned} & \left[\int_{B_{R_1}} u^{\frac{2^*}{2}(\alpha+1)} dx \right]^{\frac{2}{2^*}} \\ & \leq \mathcal{S}_2^2 \left(\frac{\lambda(\alpha+1)^2}{4|\alpha|} + \left[\frac{d(\alpha+1)}{|\alpha|(R_0-R_1)^2} + \frac{1}{R_1^2} \right] \frac{\int_{B_{R_0}} u^{\alpha+1} dx}{\int_{B_{R_0}} u^{p+\alpha} dx} \right) \int_{B_{R_0}} u^{p+\alpha} dx \\ & \leq_{(a)} \mathcal{S}_2^2 \left(\frac{\lambda(\alpha+1)^2}{4|\alpha|} + \left[\frac{d(\alpha+1)}{|\alpha|(R_0-R_1)^2} + \frac{1}{R_1^2} \right] \frac{|B_{R_0}|}{\int_{B_{R_0}} u^{p-1} dx} \right) \int_{B_{R_0}} u^{p+\alpha} dx \\ & = \frac{\mathcal{S}_2^2 |B_{R_0}|}{\|u\|_{p-1, R_0}^{p-1}} \left(\frac{\lambda(\alpha+1)^2}{4|\alpha|} \frac{\|u\|_{p-1, R_0}^{p-1}}{|B_{R_0}|} + \left[\frac{d(\alpha+1)}{|\alpha|(R_0-R_1)^2} + \frac{1}{R_1^2} \right] \right) \\ & \quad \times \int_{B_{R_0}} u^{p+\alpha} dx \\ & \leq_{(b)} \frac{\mathcal{S}_2^2 |B_{R_0}|}{\|u\|_{p-1, R_0}^{p-1}} \left(\frac{2(\alpha+1)^2}{|\alpha|(R_0-R_1)^2} + \left[\frac{d(\alpha+1)}{|\alpha|(R_0-R_1)^2} + \frac{1}{R_1^2} \right] \right) \\ & \quad \times \int_{B_{R_0}} u^{p+\alpha} dx \\ & = \frac{\mathcal{S}_2^2 |B_{R_0}|}{(R_0-R_1)^2 \|u\|_{p-1, R_0}^{p-1}} \left[\frac{1}{|\alpha|} (2(\alpha+1)^2 + d(\alpha+1)) + \frac{(R_0-R_1)^2}{R_1^2} \right] \\ & \quad \times \int_{B_{R_0}} u^{p+\alpha} dx \end{aligned} \quad (3.10)$$

where in (a) we have used the convexity in the variable $r > 0$ of the function $N(r) = \log \|u\|_r^r$, the incremental quotient is increasing, hence choosing $\alpha + 1 \geq \bar{\alpha} > 0$, we obtain

$$\frac{N(p-1+\bar{\alpha}) - N(\bar{\alpha})}{p-1} \leq \frac{N(\alpha+p) - N(\alpha+1)}{p-1}, \text{ namely } \frac{\|u\|_{p-1+\bar{\alpha}}^{p-1+\bar{\alpha}}}{\|u\|_{\bar{\alpha}}^{\bar{\alpha}}} \leq \frac{\|u\|_{\alpha+p}^{\alpha+p}}{\|u\|_{\alpha+1}^{\alpha+1}}$$

Then we have

$$\begin{aligned} \frac{\|u\|_{\alpha+p}^{\alpha+p}}{\|u\|_{\alpha+1}^{\alpha+1}} &\geq \frac{\|u\|_{p-1+\bar{\alpha}}^{p-1+\bar{\alpha}}}{\|u\|_{\bar{\alpha}}^{\bar{\alpha}}} = \frac{\|u\|_{p-1+\bar{\alpha}}^{\bar{\alpha}}}{\|u\|_{\bar{\alpha}}^{\bar{\alpha}}} \|u\|_{p-1+\bar{\alpha}}^{p-1} \\ &\geq |B_{R_0}|^{\frac{-(p-1)}{\bar{\alpha}+p-1}} |B_{R_0}|^{\frac{p-1}{\bar{\alpha}+p-1}-1} \|u\|_{p-1}^{p-1} = \frac{\|u\|_{p-1}^{p-1}}{|B_{R_0}|} \end{aligned}$$

since by Hölder inequality:

$$\frac{\|u\|_{p-1+\bar{\alpha}}}{\|u\|_{\bar{\alpha}}} \geq |B_R|^{\frac{-(p-1)}{\bar{\alpha}+p-1}} \quad \text{and} \quad \|u\|_{p-1+\bar{\alpha}} \geq |B_R|^{\frac{1}{\bar{\alpha}+p-1}-\frac{1}{p-1}} \|u\|_{p-1}.$$

In (b) we have used the Caccioppoli estimate (3.9).

Sublinear case: $0 \leq p \leq 1$. We first assume $0 \leq p < 1$, we discuss the case $p = 1$ separately. We continue estimate (3.8) as follows:

$$\begin{aligned} &\left[\int_{B_{R_1}} u^{\frac{2^*}{2}(\alpha+1)} dx \right]^{\frac{2}{2^*}} \\ &\leq S_2^2 \left(\frac{\lambda(\alpha+1)^2}{4|\alpha|} \frac{\int_{B_{R_0}} u^{p+\alpha} dx}{\int_{B_{R_0}} u^{\alpha+1} dx} + \frac{d(\alpha+1)}{|\alpha|(R_0-R_1)^2} + \frac{1}{R_1^2} \right) \int_{B_{R_0}} u^{\alpha+1} dx \\ &\leq S_2^2 \left(\frac{2(\alpha+1)^2}{|\alpha|(R_0-R_\infty)^2} + \frac{d(\alpha+1)}{|\alpha|(R_0-R_1)^2} + \frac{1}{R_1^2} \right) \int_{B_{R_0}} u^{\alpha+1} dx \\ &= \frac{S_2^2}{(R_0-R_\infty)^2} \left[\frac{1}{|\alpha|} (2(\alpha+1)^2 + d(\alpha+1)) + \frac{(R_0-R_1)^2}{R_1^2} \right] \int_{B_{R_0}} u^{\alpha+1} dx \end{aligned} \tag{3.11}$$

which follows by the convexity in the variable $r > 0$ of the function $N(r) = \log \|u\|_r^r$, which implies that the incremental quotient is increasing, hence choosing $\alpha+1 \geq \bar{\alpha} := \beta_0 > 0$, we obtain

$$\frac{N(p-1+\bar{\alpha}) - N(\bar{\alpha})}{p-1} \leq \frac{N(\alpha+p) - N(\alpha+1)}{p-1} \quad \text{namely} \quad \frac{\|u\|_{p-1+\bar{\alpha}}^{p-1+\bar{\alpha}}}{\|u\|_{\bar{\alpha}}^{\bar{\alpha}}} \leq \frac{\|u\|_{\alpha+p}^{\alpha+p}}{\|u\|_{\alpha+1}^{\alpha+1}}$$

hence

$$\begin{aligned} \frac{\int_{B_{R_0}} u^{p+\alpha} dx}{\int_{B_{R_0}} u^{\alpha+1} dx} &= \frac{\|u\|_{\alpha+p}^{\alpha+p}}{\|u\|_{\alpha+1}^{\alpha+1}} \leq \frac{\|u\|_{\bar{\alpha}-(1-p)}^{\bar{\alpha}-(1-p)}}{\|u\|_{\bar{\alpha}}^{\bar{\alpha}}} \\ &\leq \frac{|B_{R_0}|^{\frac{1-p}{\bar{\alpha}}}}{\|u\|_{\bar{\alpha}}^{1-p}} \leq \frac{\|u\|_{p-1,R_0}^{p-1}}{|B_{R_0}|} \leq \frac{8}{\lambda(R_0-R_\infty)^2} \end{aligned}$$

again by Hölder inequalities, we just stress on the last step in which we have used that

$$\frac{\|u\|_{p-1,R_0}}{|B_{R_0}|^{\frac{1}{p-1}}} \leq \frac{\|u\|_{\bar{\alpha}}}{|B_{R_0}|^{\frac{1}{\bar{\alpha}}}}, \quad \text{hence} \quad \frac{|B_{R_0}|^{\frac{1-p}{\bar{\alpha}}}}{\|u\|_{\bar{\alpha}}^{1-p}} \leq \frac{\|u\|_{p-1,R_0}^{p-1}}{|B_{R_0}|} \leq \frac{8}{\lambda(R_0-R_\infty)^2}$$

which is true since $p-1 < 0 < \bar{\alpha}$, and in the last step we have used the Caccioppoli estimate (3.9).

Notice that when $p = 1$, we obtain directly that

$$\begin{aligned}
& \left[\int_{B_{R_1}} u^{\frac{2^*}{2}(\alpha+1)} dx \right]^{\frac{2}{2^*}} \\
& \leq \mathcal{S}_2^2 \left(\frac{\lambda(\alpha+1)^2}{4|\alpha|} + \frac{d(\alpha+1)}{|\alpha|(R_0-R_1)^2} + \frac{1}{R_1^2} \right) \int_{B_{R_0}} u^{\alpha+1} dx \\
& = \frac{\mathcal{S}_2^2}{(R_0-R_\infty)^2} \left[\frac{1}{|\alpha|} \left(\frac{\lambda}{4}(\alpha+1)^2 + d(\alpha+1) \right) + \frac{(R_0-R_1)^2}{R_1^2} \right] \int_{B_{R_0}} u^{\alpha+1} dx
\end{aligned} \tag{3.12}$$

The first iteration step. We can write the first iteration step for all $p \geq 0$ in the following way: let $\beta = \alpha + 1 \geq \beta_0 > 0$ and recall that we are requiring $\beta \neq 1$ as well, then inequalities (3.10) and (3.11) can be written as

$$\left[\int_{B_{R_1}} u^{\frac{2^*}{2}\beta} dx \right]^{\frac{2}{2^*}} \leq I(p, \beta, R_1, R_0) \int_{B_{R_0}} u^{\beta+(p-1)_+} dx \tag{3.13}$$

where

$$I(p, \beta, R_1, R_0) = \frac{\mathcal{S}_2^2}{(R_0-R_1)^2} \frac{|B_{R_0}|}{\int_{B_{R_0}} u^{(p-1)_+} dx} \left[\frac{\Lambda_p \beta^2 + d\beta}{|\beta-1|} + \frac{(R_0-R_1)^2}{R_1^2} \right] \tag{3.14}$$

where $\Lambda_p = 2$ if $p \neq 1$ and $\Lambda_p = \lambda/4$ if $p = 1$.

• **STEP 3.** *The Moser iteration.* Let us define the sequence of exponents $\beta_n > 0$ so that

$$\beta_n + (p-1)_+ = \frac{2^*}{2} \beta_{n-1} \quad \text{that is} \quad \beta_n = \frac{2^*}{2} \beta_{n-1} - (p-1)_+$$

it turns out that, for any given β_0 and all $n \geq 1$:

$$\begin{aligned}
\beta_n &= \left[\frac{2^*}{2} \right]^n \left[\beta_0 - (p-1)_+ \sum_{k=0}^{n-1} \left(\frac{2^*}{2} \right)^{k-n} \right] \\
&= \left[\frac{2^*}{2} \right]^n \left[\beta_0 - (p-1)_+ \sum_{j=1}^n \left(\frac{2}{2^*} \right)^j \right] \\
&= \left[\frac{2^*}{2} \right]^n \left[\beta_0 - (p-1)_+ \frac{d-2}{2} \left(1 - \left(\frac{2}{2^*} \right)^n \right) \right] \\
&= \left[\frac{2^*}{2} \right]^n \left[\beta_0 - (p-1)_+ \frac{d-2}{2} \right] + (p-1)_+ \frac{d-2}{2}
\end{aligned} \tag{3.15}$$

since $\sum_{j=1}^k s^j = (1-s^k)s/(1-s)$. Moreover we have that for all $p \geq 1$,

$$\left(\frac{2^*}{2} \right)^{-n} \beta_n \xrightarrow{n \rightarrow \infty} \beta_0 - \frac{d-2}{2} (p-1)_+.$$

Requiring that $\beta_0 > (p-1)_+(d-2)/2$, which will be assumed from now on, then implies that $\beta_n \rightarrow +\infty$ as $n \rightarrow +\infty$. We shall also require that $\beta_n \neq 1$ for all n .

We will explicitly choose a decreasing sequence of radii $0 < R_\infty < \dots < R_n < R_{n-1} < \dots < R_0$ in the next step, in order to estimate explicitly the constants. The first iteration step then reads:

$$\begin{aligned} \|u\|_{\frac{2^*}{2}\beta_n, R_n} &= \left[\int_{B_{R_n}} u^{\frac{2^*}{2}\beta_n} dx \right]^{\frac{2}{2^*\beta_n}} \\ &\leq I(p, \beta_n, R_n, R_{n-1})^{\frac{1}{\beta_n}} \left[\int_{B_{R_{n-1}}} u^{(p-1)_+ + \beta_n} dx \right]^{\frac{1}{\beta_n}} \quad (3.16) \\ &:= I_n^{\frac{1}{\beta_n}} \|u\|_{\frac{\beta_n + (p-1)_+}{\beta_n}, R_{n-1}} = I_n^{\frac{1}{\beta_n}} \|u\|_{\frac{2^*}{2}\beta_{n-1}, R_{n-1}} \end{aligned}$$

where the constants $I(p, \beta, R_1, R_0)$ are defined in (3.14). Hence

$$\begin{aligned} I_n &= I(p, \beta_n, R_n, R_{n-1}) \\ &= \frac{S_2^2}{(R_{n-1} - R_n)^2} \frac{|B_{R_{n-1}}|}{\int_{B_{R_{n-1}}} u^{(p-1)_+} dx} \left[\frac{2\beta_n^2 + d\beta_n}{|\beta_n - 1|} + \frac{(R_{n-1} - R_n)^2}{R_n^2} \right] \quad (3.17) \end{aligned}$$

Iterating the above inequality yields

$$\begin{aligned} \|u_n\|_{\frac{2^*}{2}\beta_n, R_n} &\leq I_n^{\frac{1}{\beta_n}} \|u_n\|_{\frac{2^*}{2}\frac{\beta_{n-1}}{\beta_n}, R_{n-1}} \leq I_n^{\frac{1}{\beta_n}} I_{n-1}^{\frac{2^*}{2}\frac{1}{\beta_n}} \|u_n\|_{\frac{(\frac{2^*}{2})^2\frac{\beta_{n-2}}{\beta_n}}{2^*\beta_{n-2}}, R_{n-2}} \\ &\leq I_n^{\frac{1}{\beta_n}} I_{n-1}^{\frac{2^*}{2}\frac{1}{\beta_n}} \dots I_1^{\left(\frac{2^*}{2}\right)^{n-1}\frac{1}{\beta_n}} \|u_n\|_{\frac{(\frac{2^*}{2})^n\frac{\beta_0}{\beta_n}}{2^*\beta_0, R_0}} \quad (3.18) \\ &\leq \prod_{j=1}^n I_j^{\left(\frac{2^*}{2}\right)^{n-j}\frac{1}{\beta_n}} \|u_n\|_{\frac{(\frac{2^*}{2})^n\frac{\beta_0}{\beta_n}}{2^*\beta_0, R_0}} \end{aligned}$$

with

$$\beta_0 > \frac{d-2}{2}(p-1)_+ \quad \text{or} \quad \bar{q} := \frac{2^*}{2}\beta_0 > \frac{d(p-1)_+}{2}.$$

Taking the limit as $n \rightarrow \infty$ we obtain

$$\begin{aligned} \|u\|_{\infty, R_\infty} &= \lim_{n \rightarrow \infty} \|u\|_{\frac{2^*}{2}\beta_n, R_n} \leq \lim_{n \rightarrow \infty} \prod_{k=1}^n I_k^{\left(\frac{2^*}{2}\right)^{n-k}\frac{1}{\beta_n}} \|u\|_{\frac{\frac{\beta_0}{\beta_0 - \frac{d-2}{2}(p-1)_+}}{2^*\beta_0, R_0}} \\ &\leq \lim_{n \rightarrow \infty} \prod_{k=1}^n I_k^{\left(\frac{2^*}{2}\right)^{n-k}\frac{1}{\beta_n}} \|u\|_{\frac{\frac{\beta_0}{\beta_0 - \frac{d-2}{2}(p-1)_+}}{2^*\beta_0, R_0}} = I_\infty \|u\|_{\frac{2\bar{q}}{\bar{q}-d(p-1)_+}, R_0} \quad (3.19) \end{aligned}$$

notice that the penultimate passage follows because we shall see below that

$\prod_{k=1}^n I_k^{\left(\frac{2^*}{2}\right)^{n-k}\frac{1}{\beta_n}}$ has a limit I_∞ as $n \rightarrow +\infty$.

As a consequence of the above estimates $u \in L^\infty$, so that the above bounds holds for any $\bar{q} > d(p-1)_+/2$ as stated, provided we show that the constant I_∞ is finite and can be estimated as in (3.5).

• **STEP 4. Estimating all the constants.** Now it remains to estimate I_∞ . We will prove later that

$$I_k \leq I_0(p) \left[\frac{2^*}{2} \right]^{2k} \quad (3.20)$$

where $I_0(p)$ will have the explicit form given in formula (3.25). Using such bound we show that

$$\begin{aligned} I_\infty &= \lim_{n \rightarrow \infty} \prod_{k=1}^n I_k^{\left(\frac{2^*}{2}\right)^{n-k} \frac{1}{\beta_n}} = \lim_{n \rightarrow \infty} \exp \left[\sum_{k=1}^n \log \left(I_k^{\left(\frac{2^*}{2}\right)^{-k} \left(\frac{2^*}{2}\right)^n \frac{1}{\beta_n}} \right) \right] \\ &= \lim_{n \rightarrow \infty} \exp \left[\left(\frac{2^*}{2}\right)^n \frac{1}{\beta_n} \sum_{k=1}^n \left(\frac{2}{2^*}\right)^k \log(I_k) \right] \\ &\leq \lim_{n \rightarrow \infty} \exp \left[\left(\frac{2^*}{2}\right)^n \frac{1}{\beta_n} \sum_{k=1}^n \left(\frac{2}{2^*}\right)^k \log \left(I_0 \left[\frac{2^*}{2} \right]^{2k} \right) \right] \\ &= \lim_{n \rightarrow \infty} \exp \left[\left(\frac{2^*}{2}\right)^n \frac{1}{\beta_n} \left(\log(I_0) \sum_{k=1}^n \left(\frac{2}{2^*}\right)^k \right. \right. \\ &\quad \left. \left. + 2 \log \left(\frac{2^*}{2} \right) \sum_{k=1}^n \binom{k}{2} \right) \right] \\ &= \exp \left[\frac{1}{\beta_0 - \frac{d-2}{2}(p-1)_+} \left(\log(I_0) \sum_{k=1}^{+\infty} \left(\frac{2}{2^*}\right)^k + 2 \log \left(\frac{2^*}{2} \right) \sum_{k=1}^{+\infty} \binom{k}{2} \right) \right] \\ &= \exp \left[\frac{2}{2\beta_0 - (d-2)(p-1)_+} \left(\log(I_0) \frac{d-2}{2} + 2 \log \left(\frac{2^*}{2} \right) \frac{d(d-2)}{4} \right) \right] \\ &= \exp \left[\frac{d-2}{2\beta_0 - (d-2)(p-1)_+} \log(I_0) + \frac{d(d-2)}{2\beta_0 - (d-2)(p-1)_+} \log \left(\frac{2^*}{2} \right) \right] \\ &= I_0^{\frac{d-2}{2\beta_0 - (d-2)(p-1)_+}} \left(\frac{2^*}{2} \right)^{\frac{d(d-2)}{2\beta_0 - (d-2)(p-1)_+}} = \left[I_0 \left(\frac{2^*}{2} \right)^d \right]^{\frac{d-2}{2\beta_0 - (d-2)(p-1)_+}} \end{aligned}$$

We shall now obtain an explicit estimate for I_0 in order to finally obtain (3.5). *Estimating I_k .* We want to obtain estimates (3.20), and to this end we choose a decreasing sequence of radii $0 < R_\infty < \dots < R_k < R_{k-1} < \dots < R_0$ such that

$$(R_{k-1} - R_k)^2 = (R_0 - R_\infty)^2 \frac{c_0^2}{\beta_k} \quad \text{with} \quad c_0 = \left(\sum_{k=1}^{\infty} \sqrt{\frac{1}{\beta_k}} \right)^{-1} < +\infty$$

so that

$$\sum_{k=1}^{\infty} (R_{k-1} - R_k) = R_0 - R_\infty.$$

We now estimate I_k :

$$\begin{aligned}
I_k &= \frac{\mathcal{S}_2^2}{(R_{k-1} - R_k)^2} \frac{|B_{R_{k-1}}|}{\int_{B_{R_{k-1}}} u^{(p-1)+} dx} \left[\frac{\Lambda_p \beta_k^2 + d \beta_k}{|\beta_k - 1|} + \frac{(R_{k-1} - R_k)^2}{R_k^2} \right] \\
&= \frac{\mathcal{S}_2^2 \beta_k^2}{|\beta_k - 1| (R_{k-1} - R_k)^2} \frac{|B_{R_{k-1}}|}{\int_{B_{R_{k-1}}} u^{(p-1)+} dx} \\
&\quad \times \left[\Lambda_p + \frac{d}{\beta_k} + \frac{(R_{k-1} - R_k)^2}{R_k^2} \frac{|\beta_k - 1|}{\beta_k^2} \right] \\
&\leq^{(a)} \frac{\mathcal{S}_2^2 \beta_k^3}{c_0^2 |\beta_k - 1| (R_0 - R_\infty)^2} \frac{|B_{R_0}|}{\int_{B_{R_\infty}} u^{(p-1)+} dx} \\
&\quad \times \left[\Lambda_p + \frac{d}{\beta_0} + \frac{(R_0 - R_\infty)^2}{R_\infty^2} \max \left\{ \frac{|\beta_0 - 1|}{\beta_0^2}, \frac{1}{4} \right\} \right] \\
&\leq^{(b)} \frac{c_1 \mathcal{S}_2^2 \beta_k^2}{c_0^2 (R_0 - R_\infty)^2} \frac{|B_{R_0}|}{\int_{B_{R_\infty}} u^{(p-1)+} dx} \\
&\quad \times \left[\Lambda_p + \frac{d}{\beta_0} + \frac{(R_0 - R_\infty)^2}{R_\infty^2} \max \left\{ \frac{|\beta_0 - 1|}{\beta_0^2}, \frac{1}{4} \right\} \right] \\
&\leq^{(c)} \frac{2c_1 \mathcal{S}_2^2 [\beta_0 - (p-1) + \frac{d-2}{2}]}{c_0^2 (R_0 - R_\infty)^2} \frac{|B_{R_0}|}{\int_{B_{R_\infty}} u^{(p-1)+} dx} \\
&\quad \times \left[\Lambda_p + \frac{d}{\beta_0} + \frac{(R_0 - R_\infty)^2}{R_\infty^2} \max \left\{ \frac{|\beta_0 - 1|}{\beta_0^2}, \frac{1}{4} \right\} \right] \left[\frac{2^*}{2} \right]^{2n} \\
&\leq^{(d)} \frac{2(d-2)c_1 \mathcal{S}_2^2 |B_{R_0}|}{(\sqrt{d} - \sqrt{d-2})^2 (R_0 - R_\infty)^2} \frac{|B_{R_0}|}{\int_{B_{R_\infty}} u^{(p-1)+} dx} \\
&\quad \times \left[\Lambda_p + \frac{d}{\beta_0} + \frac{(R_0 - R_\infty)^2}{R_\infty^2} \max \left\{ \frac{|\beta_0 - 1|}{\beta_0^2}, \frac{1}{4} \right\} \right] \left[\frac{2^*}{2} \right]^{2n}
\end{aligned}$$

in (a) we have used that

$$\frac{|\beta_k - 1|}{\beta_k^2} \leq \max \left\{ \frac{|\beta_0 - 1|}{\beta_0^2}, \frac{1}{4} \right\}. \quad (3.21)$$

In (b) we have also used the inequality

$$\frac{\beta_k}{|\beta_k - 1|} \leq c_1 := \begin{cases} \frac{\beta_0}{\beta_0 - 1} & \text{if } \beta_0 > 1 \\ \max_{i=0,1} \frac{\beta_{k_0+i}}{|\beta_{k_0+i} - 1|} & \text{if } 0 < \beta_0 < 1, \end{cases} \quad (3.22)$$

with $k_0 = k_0 = i.p.$ $\left[\frac{\log \frac{1-(p-1) + \frac{d-2}{2}}{\beta_0 - (p-1) + \frac{d-2}{2}}}{\log \frac{d}{d-2}} \right]$. The inequality is stated in the general case $p \neq 1$ for later use and we shall now prove it. First notice that the numerical inequality

$$\frac{s}{|s-1|} \leq \max \left\{ \frac{a}{1-a}, \frac{b}{b-1} \right\}$$

holds true for all $0 < a < 1 < b < +\infty$ and all $s \in [0, a] \cup [b, \infty)$. When $\beta_0 > 1$ (3.22) follows applying such numerical inequality to $s = \beta_k$ and noticing that $\beta_k > \beta_0 = b > 1$ and that the function $x/|x-1|$ is decreasing when $x > 1$. Suppose instead that $0 < \beta_0 < 1$. Notice that, since we are also requiring that $\beta_0 > (p-1)_+(d-2)/2$, this is possible only when $0 < p < p_c = d/(d-2) < p_s$. We define k_0 to be the greatest integer for which $\beta_k < 1$, so that $\beta_{k_0+1} > 1$, so that

$$\beta_{k_0} < 1 < \beta_{k_0+1} \quad \text{with} \quad k_0 = i.p. \left[\frac{\log \frac{1-(p-1)_+\frac{d-2}{2}}{\beta_0-(p-1)_+\frac{d-2}{2}}}{\log \frac{d}{d-2}} \right]$$

and we shall take $\beta_0 \in (0, 1)$ such that

$$\frac{\log \frac{1-(p-1)_+\frac{d-2}{2}}{\beta_0-(p-1)_+\frac{d-2}{2}}}{\log \frac{d}{d-2}} \text{ is not an integer.} \quad (3.23)$$

The elementary properties of the function $x/|x-1|$ then show that, for all k :

$$\begin{aligned} \frac{\beta_k}{|\beta_k - 1|} &\leq \max_{i=0,1} \frac{\beta_{k_0+i}}{|\beta_{k_0+i} - 1|} \\ &= \max_{i=0,1} \frac{\left(\frac{d}{d-2}\right)^{k_0+i} \left[\beta_0 - (p-1)_+\frac{d-2}{2}\right] + (p-1)_+\frac{d-2}{2}}{\left|\left(\frac{d}{d-2}\right)^{k_0+i} \left[\beta_0 - (p-1)_+\frac{d-2}{2}\right] + (p-1)_+\frac{d-2}{2} - 1\right|} \\ &= \max_{i=0,1} \frac{\left(\frac{d}{d-2}\right)^{k_0-1+i} \left[\bar{q} - \frac{d(p-1)_+}{2}\right] + (p-1)_+\frac{d-2}{2}}{\left|\left(\frac{d}{d-2}\right)^{k_0-1+i} \left[\bar{q} - \frac{d(p-1)_+}{2}\right] + (p-1)_+\frac{d-2}{2} - 1\right|} \end{aligned}$$

as claimed, where we have put $\beta_0 = \frac{2}{2^*} \bar{q} = \frac{d-2}{d} \bar{q}$ and \bar{q} has to be chosen such that (3.23) holds.

In (c) we have used that $\beta_k = \beta_0(2^*/2)^k > \beta_0$

$$\begin{aligned} \beta_n &= \left[\frac{2^*}{2}\right]^n \left[\beta_0 - (p-1)_+\frac{d-2}{2}\right] + (p-1)_+\frac{d-2}{2} \\ &\leq 2 \left[\frac{2^*}{2}\right]^n \left[\beta_0 - (p-1)_+\frac{d-2}{2}\right] \end{aligned} \quad (3.24)$$

Finally in (d) we estimate $1/c_0^2$ as follows:

$$\begin{aligned} \frac{1}{c_0^2} &= \left(\sum_{k=1}^{\infty} \sqrt{\frac{1}{\beta_k}}\right)^2 \leq \left(\sum_{k=1}^{\infty} \frac{1}{(\beta_0 - (p-1)_+\frac{d-2}{2})^{1/2}} \left(\frac{2}{2^*}\right)^{\frac{k}{2}}\right)^2 \\ &= \frac{1}{(\beta_0 - (p-1)_+\frac{d-2}{2}) (\sqrt{d} - \sqrt{d-2})^2} \end{aligned}$$

since the explicit expression of β_k shows that

$$\beta_k \geq \left(\beta_0 - (p-1)_+ \frac{d-2}{2} \right) \left(\frac{2^*}{2} \right)^k$$

and

$$\sum_{k=1}^{+\infty} \left(\frac{2}{2^*} \right)^{k/2} = \sum_{k=1}^{+\infty} \left(\frac{d-2}{d} \right)^{k/2} = \frac{\sqrt{d-2}}{\sqrt{d} - \sqrt{d-2}}.$$

We conclude that we can take $I_0(p)$ as follows for any $p > 0$:

$$\begin{aligned} I_0(p) &= \frac{2(d-2)}{(\sqrt{d} - \sqrt{d-2})^2} \frac{c_1 \mathcal{S}_2^2}{(R_0 - R_\infty)^2} \frac{|B_{R_0}|}{\int_{B_{R_\infty}} u^{(p-1)_+} dx} \\ &\times \left[\Lambda_p + \frac{d}{\beta_0} + \frac{(R_0 - R_\infty)^2}{R_\infty^2} \max \left\{ \frac{|\beta_0 - 1|}{\beta_0^2}, \frac{1}{4} \right\} \right] \end{aligned} \quad (3.25)$$

and c_1 given by (3.22) and we recall that $\Lambda_p = 2$ if $p \neq 1$ and $\Lambda_p = \lambda/4$ if $p = 1$. The proof is concluded once we let $\beta_0 = 2\bar{q}/2^*$ as in the previous step. \square

3.2. Local upper bounds II. Linear case with unbounded coefficients

The local upper bounds for nonnegative subsolutions to

$$-\Delta u = b(x)u$$

with $b \in L^r(B_R)$ eventually unbounded, follow from the local Sobolev imbedding theorem on balls $B_R \subset \mathbb{R}^d$

$$\|f\|_{L^{2^*}(B_R)}^2 \leq \mathcal{S}_2^2 \left(\|\nabla f\|_{L^2(B_R)}^2 + \frac{1}{R^2} \|f\|_{L^2(B_R)}^2 \right) \quad (3.26)$$

where $\mathcal{S}_2 = \mathcal{S}_2(B_1)$ is the best constant and $2^* = 2d/(d-2)$. In the case $f \in W_0^{1,2}(B_R)$, we have

$$\|f\|_{L^{2^*}(B_R)}^2 \leq \mathcal{S}_2^2 \|\nabla f\|_{L^2(B_R)}^2. \quad (3.27)$$

We are requiring hereafter without any further comment that $d \geq 3$. We adopt the notation $\|f\|_{L^q(B_R)} = \|f\|_{q,R}$ and $|B_R| = \omega_d R^d$.

3.2.1. Energy Estimates and Reverse Poincaré inequalities.

Lemma 3.2. *Let $v \in L^{2^*}(B_R)$ and $b \in L^r(B_R)$ for some $r > d/2$. Then for any $\delta > 0$ the following inequality holds*

$$\begin{aligned} \int_{B_R} b(x)v^2(x) dx &\leq \delta \left[\int_{B_R} v^{2^*} dx \right]^{\frac{2}{2^*}} \\ &+ \frac{K_{r,d}^{(1)}}{\delta^{\frac{d+r(d-2)}{2r-d}}} |B_R|^{\frac{2}{2^*}} \left[\int_{B_R} b^r(x) dx \right]^{\frac{d}{2r-d}} \int_{B_R} v^2(x) dx \end{aligned} \quad (3.28)$$

where

$$K_{r,d}^{(1)} := \frac{2r-d}{rd} \left[\frac{rd}{d+r(d-2)} \right]^{\frac{d+r(d-2)}{2r-d}} \quad (3.29)$$

Proof. Let us estimate for any $0 < \varepsilon < 2$:

$$\begin{aligned}
\int_{B_R} bv^{(2-\varepsilon)+\varepsilon} dx &\leq_{(a)} \left[\int_{B_R} v^{(2-\varepsilon)\frac{2^*}{2}} dx \right]^{\frac{2}{2^*}} \left[\int_{B_R} b^{\frac{d}{2}} v^{\varepsilon\frac{d}{2}} dx \right]^{\frac{2}{d}} \\
&\leq_{(b)} |B_R|^{\frac{\varepsilon}{2^*}} \left[\int_{B_R} v^{2^*} dx \right]^{\frac{2-\varepsilon}{2^*}} \left[\int_{B_R} b^{\frac{d}{2}} v^{\varepsilon\frac{d}{2}} dx \right]^{\frac{2}{d}} \\
&\leq_{(c)} \frac{\delta_0(2-\varepsilon)}{2} \left[\int_{B_R} v^{2^*} dx \right]^{\frac{2}{2^*}} + \frac{\varepsilon}{2\delta_0^{\frac{2-\varepsilon}{\varepsilon}}} |B_R|^{\frac{2}{2^*}} \left[\int_{B_R} b^{\frac{d}{2}} v^{\varepsilon\frac{d}{2}} dx \right]^{\frac{4}{d\varepsilon}} \\
&\leq_{(d)} \delta_0 \frac{d+r(d-2)}{rd} \left[\int_{B_R} v^{2^*} dx \right]^{\frac{2}{2^*}} \\
&\quad + \frac{2(2r-d)}{2rd\delta_0^{\frac{d+r(d-2)}{2r-d}}} |B_R|^{\frac{2}{2^*}} \left[\int_{B_R} b^r dx \right]^{\frac{d}{2r-d}} \int_{B_R} v^2 dx \\
&\leq_{(e)} \delta \left[\int_{B_R} v^{2^*} dx \right]^{\frac{2}{2^*}} \\
&\quad + \frac{1}{\delta^{\frac{d+r(d-2)}{2r-d}}} \frac{2r-d}{rd} \left[\frac{rd}{d+r(d-2)} \right]^{\frac{d+r(d-2)}{2r-d}} |B_R|^{\frac{2}{2^*}} \left[\int_{B_R} b^r dx \right]^{\frac{d}{2r-d}} \int_{B_R} v^2 dx
\end{aligned}$$

where in the step (a) we have used Hölder inequality with the conjugate exponents $s = 2^*/2 = d/(d-2)$ and $s' = s/(s-1) = d/2$. In (b) we have used the inequality

$$\left[\int_{B_R} v^{(2-\varepsilon)\frac{2^*}{2}} dx \right]^{\frac{2}{2^*}} \leq |B_R|^{\frac{\varepsilon}{2^*}} \left[\int_{B_R} v^{2^*} dx \right]^{\frac{2-\varepsilon}{2^*}}$$

In (c) we have applied the Young inequality, valid for every $\sigma > 1$, $\delta_0 > 0$, $a, b \geq 0$:

$$ab \leq \frac{\delta_0}{\sigma} a^\sigma + \frac{\sigma-1}{\sigma} \frac{b^{\frac{\sigma}{\sigma-1}}}{\delta_0^{\frac{1}{\sigma-1}}}$$

with $\sigma = 2/(2-\varepsilon)$, so that $\sigma/(\sigma-1) = 2/\varepsilon$. In (d) we have used the estimate

$$\begin{aligned}
\left[\int_{B_R} b^{\frac{d}{2}} v^{\varepsilon\frac{d}{2}} dx \right]^{\frac{4}{d\varepsilon}} &\leq \left[\int_{B_R} b^r dx \right]^{\frac{d}{2r} \frac{4}{d\varepsilon}} \left[\int_{B_R} v^{\varepsilon\frac{d}{2} \frac{2r}{2r-d}} dx \right]^{\frac{2r-d}{2r} \frac{4}{d\varepsilon}} \\
&= \left[\int_{B_R} b^r dx \right]^{\frac{d}{2r-d}} \int_{B_R} v^2 dx
\end{aligned}$$

where in the first step we have used Hölder inequality with the conjugate exponents $s = 2r/d$ and $s' = s/(s-1) = 2r/(2r-d)$ (notice that we are assuming $r > d/2$, hence $s > 1$), while in the second step we have chosen $0 < \varepsilon = 2(2r-d)/(rd) \leq 2$. In (e) we have put

$$\delta = \delta_0 \frac{d+r(d-2)}{rd}$$

notice that $\delta > 0$ is in fact arbitrary since for every fixed r we can choose appropriately δ_0 to get any given value of δ by the above definition of δ . \square

Theorem 3.3 (Reverse Poincaré inequality for subsolutions). *Consider a non-negative weak subsolution u to $-\Delta u = b(x)u$ on B_R with $b \in L^r(B_R)$ with $r > d/2$. Suppose that $u \in L^{\alpha+1}(B_R)$. Then for any positive test function $\varphi \in C_0^2(B_R)$ with $|\nabla\varphi| \equiv 0$ on ∂B_R we have that for any $R > 0$ and $\alpha > 0$:*

$$\int_{B_R} |\nabla u^{\frac{\alpha+1}{2}}|^2 \varphi^2 \, dx \leq K^{(2)}[b] \int_{B_R} u^{\alpha+1} \, dx \quad (3.30)$$

with

$$\begin{aligned} K^{(2)}[b] &= K^{(2)}(b, R, \alpha, \varphi, r, d) \\ &:= \frac{\alpha+1}{\alpha} [2\|\varphi\|_\infty \|\Delta\varphi\|_\infty + \|\nabla\varphi\|_\infty^2 \\ &\quad + \mathcal{S}_2^{\frac{2[d+r(d-2)]}{2r-d}} \left(\frac{(\alpha+1)^2}{2\alpha} \right)^{\frac{r-d}{2r-d}} K_{r,d} \|\varphi\|_\infty^2 |B_R|^{\frac{2}{2^*}} \|b\|_r^{\frac{dr}{2r-d}}]. \end{aligned} \quad (3.31)$$

Remark. The requirement $u \in L^{1+\alpha}(B_R)$ will be dispensed with later, without further comment by using a Moser iteration technique.

Proof. It will be divided into several steps.

• **STEP 1. Energy estimates.** Proceeding as in (2.3), one shows that subsolutions to $-\Delta u \leq b(x)u$, satisfy, even for any $\alpha \neq -1$:

$$\frac{4\alpha}{(\alpha+1)^2} \int_{B_R} |\nabla u^{\frac{\alpha+1}{2}}|^2 \varphi^2 \, dx \leq \frac{1}{\alpha+1} \int_{B_R} u^{\alpha+1} \Delta\varphi^2 \, dx + \int_{B_R} b u^{\alpha+1} \varphi^2 \, dx. \quad (3.32)$$

• **STEP 2. Sobolev inequality in $W_0^{1,2}(B_R)$.** We apply inequality (3.28) of Lemma 3.2 to $v = u^{(\alpha+1)/2} \varphi \in W_0^{1,2}(B_R)$ so that for any $\delta > 0$:

$$\begin{aligned} \int_{B_R} b u^{\alpha+1} \varphi^2 \, dx &\leq \delta \left[\int_{B_R} \left(u^{\frac{\alpha+1}{2}} \varphi \right)^{2^*} \, dx \right]^{\frac{2}{2^*}} \\ &\quad + \frac{K_{r,d}}{\delta^{\frac{d+r(d-2)}{2r-d}}} |B_R|^{\frac{2}{2^*}} \left[\int_{B_R} b^r \, dx \right]^{\frac{d}{2r-d}} \int_{B_R} u^{\alpha+1} \varphi^2 \, dx \end{aligned} \quad (3.33)$$

where $K_{r,d}$ is given in (3.29). We notice that $v = u^{(\alpha+1)/2} \varphi \in W_0^{1,2}(B_R)$, so that the Sobolev inequality (3.27) reads

$$\begin{aligned} &\left[\int_{B_R} \left(u^{\frac{\alpha+1}{2}} \varphi \right)^{2^*} \, dx \right]^{\frac{2}{2^*}} \leq \mathcal{S}_2^2 \int_{B_R} |\nabla u^{\frac{\alpha+1}{2}} \varphi|^2 \, dx \\ &= \mathcal{S}_2^2 \left[\int_{B_R} |\nabla u^{\frac{\alpha+1}{2}}|^2 \varphi^2 \, dx + \int_{B_R} |\nabla\varphi|^2 u^{\alpha+1} \, dx + \frac{1}{2} \int_{B_R} \nabla\varphi^2 \cdot \nabla u^{\alpha+1} \, dx \right] \\ &= \mathcal{S}_2^2 \left[\int_{B_R} |\nabla u^{\frac{\alpha+1}{2}}|^2 \varphi^2 \, dx - \int_{B_R} \varphi(\Delta\varphi) u^{\alpha+1} \, dx \right] \end{aligned}$$

since $\Delta\varphi^2 = 2\varphi\Delta\varphi + 2|\nabla\varphi|^2$. We combine the above Sobolev inequality with (3.33) to get

$$\begin{aligned} \int_{B_R} b u^{\alpha+1} \varphi^2 \, dx &\leq \delta \mathcal{S}_2^2 \left[\int_{B_R} |\nabla u^{\frac{\alpha+1}{2}}|^2 \varphi^2 \, dx - \int_{B_R} \varphi (\Delta\varphi) u^{\alpha+1} \, dx \right] \\ &\quad + \frac{K_{r,d}}{\delta^{\frac{d+r(d-2)}{2r-d}}} |B_R|^{\frac{2}{2^*}} \left[\int_{B_R} b^r \, dx \right]^{\frac{d}{2r-d}} \int_{B_R} u^{\alpha+1} \varphi^2 \, dx \end{aligned} \quad (3.34)$$

where $K_{r,d}$ is given in (3.29).

• **STEP 3.** Putting the pieces together, i.e. combining inequalities (3.34) and (3.32) we obtain

$$\begin{aligned} \frac{4\alpha}{(\alpha+1)^2} \int_{B_R} |\nabla u^{\frac{\alpha+1}{2}}|^2 \varphi^2 \, dx &\leq \frac{1}{\alpha+1} \int_{B_R} u^{\alpha+1} \Delta\varphi^2 \, dx + \int_{B_R} b u^{\alpha+1} \varphi^2 \, dx \\ &\leq \frac{1}{\alpha+1} \int_{B_R} u^{\alpha+1} \Delta\varphi^2 \, dx + \delta \mathcal{S}_2^2 \left[\int_{B_R} |\nabla u^{\frac{\alpha+1}{2}}|^2 \varphi^2 \, dx - \int_{B_R} \varphi (\Delta\varphi) u^{\alpha+1} \, dx \right] \\ &\quad + \frac{K_{r,d}}{\delta^{\frac{d+r(d-2)}{2r-d}}} |B_R|^{\frac{2}{2^*}} \left[\int_{B_R} b^r \, dx \right]^{\frac{d}{2r-d}} \int_{B_R} u^{\alpha+1} \varphi^2 \, dx \end{aligned}$$

which thus implies

$$\begin{aligned} &\left(\frac{4\alpha}{(\alpha+1)^2} - \delta \mathcal{S}_2^2 \right) \int_{B_R} |\nabla u^{\frac{\alpha+1}{2}}|^2 \varphi^2 \, dx \\ &\leq \frac{1}{\alpha+1} \int_{B_R} u^{\alpha+1} \Delta\varphi^2 \, dx - \delta \mathcal{S}_2^2 \int_{B_R} \varphi (\Delta\varphi) u^{\alpha+1} \, dx \\ &\quad + \frac{K_{r,d}}{\delta^{\frac{d+r(d-2)}{2r-d}}} |B_R|^{\frac{2}{2^*}} \left[\int_{B_R} b^r \, dx \right]^{\frac{d}{2r-d}} \int_{B_R} u^{\alpha+1} \varphi^2 \, dx \\ &\leq \left[\left(\frac{2}{\alpha+1} + \delta \mathcal{S}_2^2 \right) \|\varphi\|_\infty \|\Delta\varphi\|_\infty + \frac{2}{\alpha+1} \|\nabla\varphi\|_\infty^2 \right. \\ &\quad \left. + \frac{K_{r,d} \|\varphi\|_\infty^2}{\delta^{\frac{rd}{2r-d}-1}} |B_R|^{\frac{2}{2^*}} \left(\int_{B_R} b^r \, dx \right)^{\frac{d}{2r-d}} \right] \int_{B_R} u^{\alpha+1} \, dx \end{aligned}$$

Letting $\delta \mathcal{S}_2^2 = \frac{2\alpha}{(\alpha+1)^2}$ gives the following reverse Poincaré inequality:

$$\int_{B_R} |\nabla u^{\frac{\alpha+1}{2}}|^2 \varphi^2 \, dx \leq \Lambda_0 \int_{B_R} u^{\alpha+1} \, dx$$

with the constant that we can estimate as follows

$$\begin{aligned}
\Lambda_0 &= \frac{(\alpha+1)^2}{2\alpha} \left[\frac{2(2\alpha+1)}{(\alpha+1)^2} \|\varphi\|_\infty \|\Delta\varphi\|_\infty + \frac{2}{\alpha+1} \|\nabla\varphi\|_\infty^2 \right. \\
&\quad \left. + \mathcal{S}_2^{\frac{2[d+r(d-2)]}{2r-d}} \frac{2\alpha}{(\alpha+1)^2} \left(\frac{(\alpha+1)^2}{2\alpha} \right)^{\frac{rd}{2r-d}} K_{r,d} \|\varphi\|_\infty^2 |B_R|^{\frac{2}{2^*}} \|b\|_r^{\frac{dr}{2r-d}} \right] \\
&\leq \frac{\alpha+1}{\alpha} [2\|\varphi\|_\infty \|\Delta\varphi\|_\infty + \|\nabla\varphi\|_\infty^2 \\
&\quad + \mathcal{S}_2^{\frac{2[d+r(d-2)]}{2r-d}} \left(\frac{(\alpha+1)^2}{2\alpha} \right)^{\frac{rd}{2r-d}} K_{r,d} \|\varphi\|_\infty^2 |B_R|^{\frac{2}{2^*}} \|b\|_r^{\frac{dr}{2r-d}}] \\
&\leq \frac{(\alpha+1)^{1+\frac{rd}{2r-d}}}{\alpha} [2\|\varphi\|_\infty \|\Delta\varphi\|_\infty + \|\nabla\varphi\|_\infty^2 \\
&\quad + \mathcal{S}_2^{\frac{2[d+r(d-2)]}{2r-d}} \left(\frac{\alpha+1}{2\alpha} \right)^{\frac{rd}{2r-d}} K_{r,d} \|\varphi\|_\infty^2 |B_R|^{\frac{2}{2^*}} \|b\|_r^{\frac{dr}{2r-d}}] =: K^{(2)}[b]. \quad \square
\end{aligned}$$

In fact, the last bound in the above formula for $K^{(2)}[b]$ could be avoided, but will make the following calculations somewhat easier.

The next calculus lemma, whose proof is straightforward, will be of great help in performing explicitly the Moser iteration.

Lemma 3.4 (Numerical Iteration). *Let $Y_n \geq 0$ be a sequence of numbers such that*

$$Y_n \leq I_{n-1}^{\sigma\theta^{n-1}} Y_{n-1} \quad \text{with} \quad I_{n-1} \leq I_0 C^{n-1} \quad (3.35)$$

for some $\sigma, I_0, C > 0$, $\theta \in (0, 1)$. Then $\{Y_n\}$ is a bounded sequence and one has

$$Y_\infty := \limsup_{n \rightarrow +\infty} Y_n \leq I_0^{\frac{\sigma}{1-\theta}} C^{\frac{\sigma\theta}{(1-\theta)^2}} Y_0. \quad (3.36)$$

Now we are ready to perform the Moser iteration, by combining a local Sobolev inequality with the reverse Poincaré inequality of Theorem 3.3 and then using the above numerical Lemma.

Theorem 3.5 (Moser Iteration). *Let $u \geq 0$ be a weak subsolution to $-\Delta u = bu$ on B_R with $b \in L^r(B_R)$ with $r > d/2$, and let $q > 1$, $R_\infty < R_0 < R$.*

$$\|u\|_{\infty, R_\infty} \leq \frac{K_q^{(3)}[b]}{(R_0 - R_\infty)^{\frac{d}{q}}} \|u\|_{q, R_0} \quad (3.37)$$

with constant

$$\begin{aligned}
K_q^{(3)}[b] &= \left(\frac{qd^d}{2^d} \right)^{\frac{rd^2}{2(2r-d)q}} \left[8 \frac{q(d+2)}{q-1} + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^2 \right. \\
&\quad \left. + \left(\frac{\mathcal{S}_2^2}{2} \right)^{\frac{rd}{2r-d}} \frac{2r-d}{rd} \left(\frac{qrd}{(q-1)[d+r(d-2)]} \right)^{1+\frac{rd}{2r-d}} \right. \\
&\quad \left. \times (R_0 - R_\infty)^2 |B_{R_0}|^{\frac{2}{2^*}} \|b\|_{L^r(B_{R_0})}^{\frac{rd}{2r-d}} \right]^{\frac{d}{2q}}.
\end{aligned} \quad (3.38)$$

Notice that in the case of bounded coefficients $b(x) \in L^\infty(B_{R_0})$ we can pass to the limit as $r \rightarrow \infty$ in the above expression of $K_q^{(3)}[b]$ to get

$$K_q^{(3)}[b] = \left(\frac{qd^d}{2^d}\right)^{\frac{d^2}{4q}} \left[8 \frac{q(d+2)}{q-1} + \left(\frac{\mathcal{S}_2^2}{2}\right)^{\frac{d}{2}} \frac{2}{d} \left(\frac{qd}{(q-1)(d-2)}\right)^{1+\frac{d}{2}} \right. \\ \left. \times (R_0 - R_\infty)^2 |B_{R_0}|^{\frac{2}{2^*}} \|b\|_{L^\infty(B_{R_0})}^{\frac{d}{2}} + \left(\frac{R_0 - R_\infty}{R_\infty}\right)^2 \right]^{\frac{d}{2q}}. \quad (3.39)$$

Proof. The proof is divided in several steps.

• **STEP 1. Sobolev and Reverse Poincaré inequalities.** We start choosing radii r_1, r_0 with $R_\infty < r_1 < r_0 < R_0$ and use the test function of Lemma 2.2 on the balls B_{r_1}, B_{r_0} . We use the Reverse Poincaré inequality (3.30) on the ball B_{r_0} and the fact that $\varphi \equiv 1$ on B_{r_1} to get

$$\int_{B_{r_1}} |\nabla u^{\frac{\alpha+1}{2}}|^2 dx \leq \int_{B_{r_0}} |\nabla u^{\frac{\alpha+1}{2}}|^2 \varphi^2 dx \leq K^{(2)}[b] \int_{B_{r_0}} u^{\alpha+1} dx$$

so that the local Sobolev inequality in $W^{1,2}(B_{r_1})$ applied to $f = v^{\frac{\alpha+1}{2}}$ for any $\alpha > 0$ yields

$$\left(\int_{B_{r_1}} u^{\frac{2^*}{2}(\alpha+1)} dx \right)^{\frac{2}{2^*}} \leq \mathcal{S}_2^2 \left[\int_{B_{r_1}} |\nabla u^{\frac{\alpha+1}{2}}|^2 dx + \frac{1}{r_1^2} \int_{B_{r_1}} u^{\alpha+1} dx \right] \\ \leq \mathcal{S}_2^2 \left(K^{(2)}[b] + \frac{1}{r_1^2} \right) \int_{B_{r_0}} u^{\alpha+1} dx \quad (3.40)$$

where the constant $K^{(2)}[b]$ is given by (3.30), and we can estimate it as follows:

$$K^{(2)}[b] = \frac{(\alpha+1)^{1+\frac{rd}{2r-d}}}{\alpha} \left[2\|\varphi\|_{\infty, r_0} \|\Delta\varphi\|_{\infty, r_0} + \|\nabla\varphi\|_{\infty, r_0}^2 \right. \\ \left. + \mathcal{S}_2^{\frac{2[d+r(d-2)]}{2r-d}} \left(\frac{\alpha+1}{2\alpha}\right)^{\frac{rd}{2r-d}} \frac{2r-d}{rd} \left(\frac{rd}{d+r(d-2)}\right)^{1+\frac{rd}{2r-d}} \right. \\ \left. \times \|\varphi\|_{\infty, r_0}^2 |B_{r_0}|^{\frac{2}{2^*}} \|b\|_{L^r(B_{r_0})}^{\frac{rd}{2r-d}} \right] \\ \leq_{(a)} \frac{\alpha+1}{\alpha} (\alpha+1)^{\frac{rd}{2r-d}} \left[\frac{8(d+2)}{(r_0-r_1)^2} + \mathcal{S}_2^{\frac{2[d+r(d-2)]}{2r-d}} \left(\frac{\alpha+1}{2\alpha}\right)^{\frac{rd}{2r-d}} \frac{2r-d}{rd} \right. \\ \left. \times \left(\frac{rd}{d+r(d-2)}\right)^{1+\frac{rd}{2r-d}} |B_{r_0}|^{\frac{2}{2^*}} \|b\|_{L^r(B_{r_0})}^{\frac{rd}{2r-d}} \right]$$

$$\begin{aligned} &\leq_{(b)} \frac{(\alpha + 1)^{\frac{rd}{2r-d}}}{(r_0 - r_1)^2} \left[8(d+2) \frac{\alpha + 1}{\alpha} + \frac{\mathcal{S}_2^{\frac{2[d+r(d-2)]}{2r-d}}}{2^{\frac{rd}{2r-d}}} \frac{2r-d}{rd} \right. \\ &\quad \left. \times \left(\frac{(\alpha + 1)rd}{\alpha[d+r(d-2)]} \right)^{1+\frac{rd}{2r-d}} (R_0 - R_\infty)^2 |B_{R_0}|^{\frac{2}{2^*}} \|b\|_{L^r(B_{R_0})}^{\frac{rd}{2r-d}} \right] \end{aligned}$$

where in (a) we have used the fact that the test function of Lemma 2.2 satisfies $\|\varphi\|_{\infty, r_0} = 1$, $\|\nabla\varphi\|_{\infty, r_0} \leq 4/(r_0 - r_1)$ and $\|\Delta\varphi\|_{\infty, r_0} \leq 4d/(r_0 - r_1)^2$, and in (b) the fact that $0 < R_\infty < r_1 < r_0 < R_0$. Finally we get:

$$\begin{aligned} \mathcal{S}_2^2 \left(K^{(2)}[b] + \frac{1}{r_1^2} \right) &\leq \mathcal{S}_2^2 \frac{(\alpha + 1)^{\frac{rd}{2r-d}}}{(r_0 - r_1)^2} \left[8(d+2) \frac{\alpha + 1}{\alpha} + \frac{\mathcal{S}_2^{\frac{2[d+r(d-2)]}{2r-d}}}{2^{\frac{rd}{2r-d}}} \frac{2r-d}{rd} \right. \\ &\quad \left. \times \left(\frac{(\alpha + 1)rd}{\alpha[d+r(d-2)]} \right)^{1+\frac{rd}{2r-d}} (R_0 - R_\infty)^2 |B_{R_0}|^{\frac{2}{2^*}} \|b\|_{L^r(B_{R_0})}^{\frac{rd}{2r-d}} + \frac{1}{r_1^2} \frac{(r_0 - r_1)^2}{(\alpha + 1)^{\frac{rd}{2r-d}}} \right] \\ &\leq \mathcal{S}_2^2 \frac{(\alpha + 1)^{\frac{rd}{2r-d}}}{(r_0 - r_1)^2} \left[8(d+2) \frac{\alpha + 1}{\alpha} + \frac{\mathcal{S}_2^{\frac{2[d+r(d-2)]}{2r-d}}}{2^{\frac{rd}{2r-d}}} \frac{2r-d}{rd} \right. \\ &\quad \left. \times \left(\frac{(\alpha + 1)rd}{\alpha[d+r(d-2)]} \right)^{1+\frac{rd}{2r-d}} (R_0 - R_\infty)^2 |B_{R_0}|^{\frac{2}{2^*}} \|b\|_{L^r(B_{R_0})}^{\frac{rd}{2r-d}} + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^2 \right] \end{aligned} \quad (3.41)$$

we have also used the fact that $\alpha > 0$.

• **STEP 2. The Moser iteration.** We now fix $\beta_0 = \alpha + 1 > 1$, and we define the sequence

$$\beta_n = \frac{2^*}{2} \beta_{n-1} = \left(\frac{2^*}{2} \right)^n \beta_0$$

Next we pick a sequence of radii $R_\infty = r_\infty < \dots < r_n < r_{n-1} < \dots < r_0 = R_0$, such that

$$(r_{n-1} - r_n)^2 = c_0^2 (R_0 - R_\infty)^2 \left(\frac{2}{2^*} \right)^{\frac{rdn}{2r-d}}$$

with

$$\begin{aligned} c_0 &= \left(\sum_{k=1}^{\infty} \left(\frac{2}{2^*} \right)^{\frac{rd}{2(2r-d)}k} \right)^{-1} = \left(\frac{2^*}{2} \right)^{\frac{rd}{2(2r-d)}} - 1 \geq \left(\frac{2^*}{2} - 1 \right)^{\frac{rd}{2(2r-d)}} \\ &= \left(\frac{2}{d-2} \right)^{\frac{rd}{2(2r-d)}} \end{aligned} \quad (3.42)$$

where the inequality in the above formula is easily shown to hold when $d \geq 3$ and $r > d/2$ as assumed, so that

$$\sum_{k=1}^{\infty} (r_{k-1} - r_k) = R_0 - R_\infty,$$

the above series being convergent. With these choices, inequality (3.40) in which $\alpha + 1$ is replaced by β_{n-1} , this being allowable since $\beta_n > 1$ for all n , and r_1, r_0 replaced by r_n, r_{n-1} reads, noticing in addition that $\beta_n/(\beta_n - 1) \leq \beta_0/(\beta_0 - 1)$ for all n ,

$$\begin{aligned}
& \left(\int_{B_{r_n}} u^{\frac{2^*}{2} \beta_{n-1}} dx \right)^{\frac{2}{2^*}} \leq \mathcal{S}_2^2 \left(K^{(2)}[b] + \frac{1}{r_n^2} \right) \int_{B_{r_{n-1}}} u^{\beta_{n-1}} dx \\
& \leq \frac{\mathcal{S}_2^2 \beta_{n-1}^{\frac{rd}{2r-d}}}{(r_{n-1} - r_n)^2} \left[8(d+2) \frac{\beta_0}{\beta_0 - 1} \right. \\
& \quad \left. + \frac{\mathcal{S}_2^{\frac{2[d+r(d-2)]}{2r-d}}}{2^{\frac{rd}{2r-d}}} \frac{2r-d}{rd} \left(\frac{\beta_0 rd}{(\beta_0 - 1)[d+r(d-2)]} \right)^{1+\frac{rd}{2r-d}} \right. \\
& \quad \left. \times (R_0 - R_\infty)^2 |B_{R_0}|^{\frac{2}{2^*}} \|b\|_{L^r(B_{R_0})}^{\frac{rd}{2r-d}} + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^2 \right] \int_{B_{r_{n-1}}} u^{\beta_{n-1}} dx \\
& := I_{n-1} \int_{B_{r_{n-1}}} u^{\beta_{n-1}} dx
\end{aligned}$$

Letting $Y_n := \|u\|_{\beta_n, R_n}$, we have obtained

$$\begin{aligned}
Y_n &= \|u\|_{\beta_n, R_n} \leq I_{n-1}^{\frac{1}{\beta_{n-1}}} \|u\|_{\beta_{n-1}, R_{n-1}} = I_{n-1}^{\frac{1}{\beta_{n-1}}} Y_{n-1} = I_{n-1}^{\frac{1}{\beta_0} \left(\frac{2}{2^*}\right)^{n-1}} Y_{n-1} \\
&= I_{n-1}^{\theta^{n-1}} Y_{n-1}
\end{aligned}$$

where we have set $\sigma = 1/\beta_0$ and $\theta = 2/2^* \in (0, 1)$. We shall prove that $I_n \leq I_0 C^n$. Indeed:

$$\begin{aligned}
I_{n-1} &= \frac{\mathcal{S}_2^2 \beta_{n-1}^{\frac{rd}{2r-d}}}{(r_{n-1} - r_n)^2} \left[8(d+2) \frac{\beta_0}{\beta_0 - 1} \right. \\
& \quad \left. + \frac{\mathcal{S}_2^{\frac{2[d+r(d-2)]}{2r-d}}}{2^{\frac{rd}{2r-d}}} \frac{2r-d}{rd} \left(\frac{\beta_0 rd}{(\beta_0 - 1)[d+r(d-2)]} \right)^{1+\frac{rd}{2r-d}} \right. \\
& \quad \left. \times (R_0 - R_\infty)^2 |B_{R_0}|^{\frac{2}{2^*}} \|b\|_{L^r(B_{R_0})}^{\frac{rd}{2r-d}} + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^2 \right] \\
& \leq \frac{\beta_0^{\frac{rd}{2r-d}}}{c_0^2 (R_0 - R_\infty)^2} \left[8(d+2) \frac{\beta_0}{\beta_0 - 1} \right. \\
& \quad \left. + \frac{\mathcal{S}_2^{\frac{2[d+r(d-2)]}{2r-d}}}{2^{\frac{rd}{2r-d}}} \frac{2r-d}{rd} \left(\frac{\beta_0 rd}{(\beta_0 - 1)[d+r(d-2)]} \right)^{1+\frac{rd}{2r-d}} \right. \\
& \quad \left. \times (R_0 - R_\infty)^2 |B_{R_0}|^{\frac{2}{2^*}} \|b\|_{L^r(B_{R_0})}^{\frac{rd}{2r-d}} + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^2 \right] \left(\frac{2^*}{2} \right)^{\frac{2rdn}{2r-d}}
\end{aligned}$$

$$\begin{aligned}
&\leq \left(\frac{d-2}{2}\right)^{\frac{rd}{2r-d}} \frac{\beta_0^{\frac{rd}{2r-d}}}{(R_0 - R_\infty)^2} \left[8(d+2) \frac{\beta_0}{\beta_0 - 1} \right. \\
&+ \frac{\mathcal{S}_2^{\frac{2[d+r(d-2)]}{2r-d}}}{2^{\frac{rd}{2r-d}}} \left(\frac{\beta_0 rd}{(\beta_0 - 1)[d+r(d-2)]} \right)^{1+\frac{rd}{2r-d}} \\
&\times \frac{2r-d}{rd} (R_0 - R_\infty)^2 |B_{R_0}|^{\frac{2}{2^*}} \|b\|_{L^r(B_{R_0})}^{\frac{rd}{2r-d}} + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^2 \left. \right] \left(\frac{2^*}{2} \right)^{\frac{2rd}{2r-d}n} \\
&:= I_0 C^{n-1}
\end{aligned} \tag{3.43}$$

where in the last inequality we estimated c_0 as in (3.42). Finally we use Lemma 3.4 with the above choices of σ and θ , thus proving that

$$Y_\infty \leq I_0^{1-\theta} C^{\frac{\sigma\theta}{(1-\theta)^2}} Y_0$$

namely

$$\|u\|_{\infty, R_\infty} \leq I_0^{\frac{d}{2\beta_0}} C^{\frac{d(d-2)}{4\beta_0}} \|u\|_{\beta_0, R_0} = K_q^{(3)}[b] \|u\|_{\beta_0, R_0}$$

which is exactly (3.37) with

$$\begin{aligned}
K_q^{(3)}[b] &= \left(\frac{d-2}{2}\right)^{\frac{rd^2}{2(2r-d)\beta_0}} \frac{\beta_0^{\frac{rd^2}{2(2r-d)\beta_0}}}{(R_0 - R_\infty)^{\frac{d}{\beta_0}}} \\
&\times \left[8(d+2) \frac{\beta_0}{\beta_0 - 1} + \frac{\mathcal{S}_2^{\frac{2[d+r(d-2)]}{2r-d}}}{2^{\frac{rd}{2r-d}}} \left(\frac{\beta_0 rd}{(\beta_0 - 1)[d+r(d-2)]} \right)^{1+\frac{rd}{2r-d}} \right. \\
&\times \frac{2r-d}{rd} (R_0 - R_\infty)^2 |B_{R_0}|^{\frac{2}{2^*}} \|b\|_{L^r(B_{R_0})}^{\frac{rd}{2r-d}} + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^2 \left. \right]^{\frac{d}{2\beta_0}} \\
&\times \left(\frac{d}{d-2} \right)^{\frac{rd^2}{(2r-d)\beta_0}} \left(\frac{d}{d-2} \right)^{\frac{rd^2(d-2)}{2\beta_0(2r-d)}} \\
&\leq \left(\frac{d}{2}\right)^{\frac{rd^3}{2(2r-d)\beta_0}} \frac{\beta_0^{\frac{rd^2}{2(2r-d)\beta_0}}}{(R_0 - R_\infty)^{\frac{d}{\beta_0}}} \left[8(d+2) \frac{\beta_0}{\beta_0 - 1} \right. \\
&+ \frac{\mathcal{S}_2^{\frac{2[d+r(d-2)]}{2r-d}}}{2^{\frac{rd}{2r-d}}} \left(\frac{\beta_0 rd}{(\beta_0 - 1)[d+r(d-2)]} \right)^{1+\frac{rd}{2r-d}} \\
&\times \frac{2r-d}{rd} (R_0 - R_\infty)^2 |B_{R_0}|^{\frac{2}{2^*}} \|b\|_{L^r(B_{R_0})}^{\frac{rd}{2r-d}} + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^2 \left. \right]^{\frac{d}{2\beta_0}},
\end{aligned}$$

as in (3.38). The proof is concluded once we let $\beta_0 = q > 1$. \square

3.2.2. Extending local upper bounds. A lemma by De Giorgi. In this section extend the local upper bound of the previous section. More precisely we show

that a bound of the type

$$\|u\|_{\infty,r} \leq \frac{A}{(R-r)^{\frac{d}{q}}} \|u\|_{q,R}$$

which is valid for any $q > 1$ and any $a \leq r < R \leq b$ indeed implies that

$$\|u\|_{\infty,a} \leq \frac{A}{(b-a)^{\frac{d}{q_0}}} \|u\|_{q_0,b}$$

for all $q_0 > 0$ and any $a \leq r < R \leq b$ maybe with a different constant A . The proof relies on the following lemma, originally due to E. De Giorgi, whose proof is contained in several books and papers, see for example [23], Lemma 6.1.

Lemma 3.6 (De Giorgi). *Let $Z(t)$ be a bounded non-negative function in the interval $[t_0, t_1]$. Assume that for $t_0 \leq t < s \leq t_1$ we have*

$$Z(t) \leq \theta Z(s) + \frac{A}{(s-t)^\alpha} \quad (3.44)$$

with $A \geq 0$, $\alpha > 0$ and $0 \leq \theta < 1$. Then

$$Z(t_0) \leq \frac{Ac(\alpha, \lambda, \theta)}{(t_1 - t_0)^\alpha} \quad (3.45)$$

where

$$c(\alpha, \lambda, \theta) = \frac{1}{(1-\lambda)^\alpha \left(1 - \frac{\theta}{\lambda^\alpha}\right)} \quad \text{for any } \lambda \in \left(\theta^{\frac{1}{\alpha}}, 1\right).$$

The above Lemma has important consequences, indeed it allows to prove that if a reverse Hölder inequality holds for some $0 < q < \bar{q}$, then it holds for any $0 < q_0 < \bar{q}$.

Lemma 3.7 (Extending Local Upper Bounds). *Assume that the following bounds holds true:*

$$\|u\|_{\bar{q},r} \leq \frac{K}{(R-r)^\gamma} \|u\|_{\underline{q},R} \quad (3.46)$$

for some $0 < \underline{q} < \bar{q}$, $\gamma > 0$ and for any $R_\infty \leq r < R \leq R_0$. Then we have that for all $0 < q_0 \leq \underline{q} < \bar{q}$

$$\|u\|_{\bar{q},R_\infty} \leq 3 \cdot 2^{\frac{\bar{q}(q-q_0)}{q_0(\bar{q}-\underline{q})}} \left[\left(4\gamma \frac{q(\bar{q}-q_0)}{q_0(\bar{q}-\underline{q})} \right)^\gamma \frac{K}{(R_0 - R_\infty)^\gamma} \right]^{\frac{q(\bar{q}-q_0)}{q_0(\bar{q}-\underline{q})}} \|u\|_{q_0,R_0}. \quad (3.47)$$

Proof. Define, for $t < R_0$, the bounded nonnegative function

$$Z(t) = \|u\|_{L^{\bar{q}}(B_t)} = \|u\|_{\bar{q},t}$$

then (3.46) reads, for $s > t$,

$$Z(t) = \|u\|_{\bar{q},t} \leq \frac{K}{(s-t)^\gamma} \|u\|_{\underline{q},s} \leq \frac{K}{(s-t)^\gamma} \|u\|_{q_0,s}^{1-\sigma} \|u\|_{\bar{q},s}^\sigma, \quad (3.48)$$

where in the last step we have used that for all $0 < q_0 \leq \underline{q} < \bar{q} \leq +\infty$

$$\|u\|_{\underline{q},s} \leq \|u\|_{q_0,s}^{1-\sigma} \|u\|_{\bar{q},s}^{\sigma} = \|u\|_{\frac{q_0(\bar{q}-\underline{q})}{\underline{q}(\bar{q}-q_0)}}^{\frac{q_0(\bar{q}-\underline{q})}{\underline{q}(\bar{q}-q_0)}} \|u\|_{\frac{\bar{q}(q-q_0)}{\underline{q}(\bar{q}-q_0)}}^{\frac{\bar{q}(q-q_0)}{\underline{q}(\bar{q}-q_0)}}, \quad \sigma = \frac{\bar{q}(q-q_0)}{\underline{q}(\bar{q}-q_0)} \in [0,1)$$

Inequality (3.48) gives then

$$\begin{aligned} Z(t) &= \|u\|_{\bar{q},t} \leq \frac{K}{(s-t)^{\gamma}} \|u\|_{q_0,s}^{1-\sigma} Z(s)^{\sigma} \leq \frac{1}{2} Z(s) + \frac{(2^{\sigma} K)^{\frac{1}{1-\sigma}}}{(s-t)^{\frac{\gamma}{1-\sigma}}} \|u\|_{q_0,s} \\ &\leq \frac{1}{2} Z(s) + \frac{(2^{\sigma} K)^{\frac{1}{1-\sigma}}}{(s-t)^{\frac{\gamma}{1-\sigma}}} \|u\|_{q_0,R_0} \end{aligned} \tag{3.49}$$

where we have used Young's inequality valid for any $\nu > 1$, $a, b \geq 0$, $\varepsilon > 0$:

$$ab \leq \frac{\varepsilon}{\nu} a^{\nu} + \frac{\nu-1}{\nu} \frac{b^{\frac{\nu}{\nu-1}}}{\varepsilon^{\frac{1}{\nu-1}}} \leq \varepsilon a^{\nu} + \frac{b^{\frac{\nu}{\nu-1}}}{\varepsilon^{\frac{1}{\nu-1}}}$$

with the choices

$$\varepsilon = 1/2 \quad a = Z(s)^{\sigma}, \quad \nu = \frac{1}{\sigma} > 1 \quad \text{and} \quad b = \frac{K}{(s-t)^{\gamma}} \|u\|_{q_0,R_0}^{1-\sigma}.$$

Inequality (3.49) is of the form appearing in Lemma 3.6 with $\alpha = \gamma/(1-\sigma) > 0$, $\theta = 1/2$ and $A = (2^{\sigma} K)^{\frac{1}{1-\sigma}} \|u\|_{q_0,R_0}$. Thus we get

$$\begin{aligned} \|u\|_{\bar{q},R_{\infty}} &= Z(R_{\infty}) \leq \frac{c(\alpha, \lambda, \theta) (2^{\sigma} K)^{\frac{1}{1-\sigma}}}{(R_0 - R_{\infty})^{\frac{\gamma}{1-\sigma}}} \|u\|_{q_0,R_0} \\ &\leq 3 \left(\frac{4\gamma}{1-\sigma} \right)^{\frac{1}{1-\sigma}} \frac{(2^{\sigma} K)^{\frac{1}{1-\sigma}}}{(R_0 - R_{\infty})^{\frac{\gamma}{1-\sigma}}} \|u\|_{q_0,R_0} \\ &= 3 \cdot 2^{\frac{\bar{q}(q-q_0)}{q_0(\bar{q}-\underline{q})}} \left[\left(4\gamma \frac{q(\bar{q}-q_0)}{q_0(\bar{q}-\underline{q})} \right)^{\gamma} \frac{K}{(R_0 - R_{\infty})^{\gamma}} \right]^{\frac{q_0(\bar{q}-\underline{q})}{\bar{q}(q-q_0)}} \|u\|_{q_0,R_0} \end{aligned}$$

noticing that

$$\frac{\sigma}{1-\sigma} = \frac{\bar{q}(q-q_0)}{q_0(\bar{q}-\underline{q})}, \quad \text{and} \quad \alpha = \frac{\gamma}{1-\sigma} = \gamma \frac{q(\bar{q}-q_0)}{q_0(\bar{q}-\underline{q})}$$

which is the desired bound, once we notice that whenever $\theta < \lambda^{\alpha} < 1$,

$$\begin{aligned} c(\alpha, \lambda, \theta) &= \frac{1}{(1-\lambda)^{\alpha} \left(1 - \frac{\theta}{\lambda^{\alpha}}\right)} = \frac{2(1+\theta)}{\left[2^{\frac{1}{\alpha}} - (1+\theta)^{\frac{1}{\alpha}}\right]^{\alpha} (1-\theta)} \\ &= \frac{12}{\left[4^{\frac{1}{\alpha}} - 3^{\frac{1}{\alpha}}\right]^{\alpha}} \leq 12 \frac{4^{\alpha} \alpha^{\alpha}}{4} = 3(4\alpha)^{\alpha} \end{aligned}$$

since we can choose $1/2 = \theta < \lambda^{\alpha} = (1+\theta)/2 < 1$, and since $\alpha = \gamma/(1-\sigma) > 1$, $(4^{1/\alpha} - 3^{1/\alpha})^{\alpha} \geq \frac{4}{4^{\alpha} \alpha^{\alpha}}$, since we know that $a^{1/\alpha} - b^{1/\alpha} \geq a^{1/\alpha} (a-b)/(\alpha a)$, for all $a \geq b \geq 0$ and $\alpha \geq 1$. \square

The above lemma can be used to extend the local upper bounds (3.50) of Theorem 3.5.

Theorem 3.8 (Local Upper bounds, unbounded coefficient). *Consider a non-negative weak subsolution u to $-\Delta u = bu$ in B_R with $b \in L^r(B_R)$ and $r > d/2$. Let $0 < R_\infty < R_0 < R$. Then, for any $q_0 > 0$ the following bound holds true*

$$\|u\|_{\infty, R_\infty} \leq \frac{A_{q_0}^{(1)}}{(R - R_\infty)^{\frac{d}{q_0}}} \left[A_{q_0}^{(2)} + A_{q_0}^{(3)} \|b\|_{L^r(B_{R_0})}^{\frac{rd}{2r-d}} \right]^{\frac{d}{2q_0}} \|u\|_{q_0, R} \quad (3.50)$$

with

$$A_{q_0}^{(1)} := \begin{cases} \left(\frac{q_0 d^d}{2^d} \right)^{\frac{rd^2}{2(2r-d)q_0}}, & \text{if } q_0 > 1, \\ 3 \cdot 2^{\frac{2d+1}{q_0}} \left(\frac{d}{q_0} \right)^{\frac{d}{q_0}} \left(\frac{(q_0+1)d^d}{2^d} \right)^{\frac{rd^2}{2(2r-d)q_0}}, & \text{if } 0 < q_0 \leq 1, \end{cases} \quad (3.51)$$

$$A_{q_0}^{(2)} := \begin{cases} 8 \frac{q_0(d+2)}{q_0-1} + \left(\frac{R-R_\infty}{R_\infty} \right)^2, & \text{if } q_0 > 1, \\ 8 \frac{(q_0+1)(d+2)}{q_0} + \left(\frac{R-R_\infty}{R_\infty} \right)^2, & \text{if } 0 < q_0 \leq 1, \end{cases} \quad (3.52)$$

$$A_{q_0}^{(3)} := \begin{cases} \left(\frac{S_2^2}{2} \right)^{\frac{rd}{2r-d}} \frac{2r-d}{rd} \left(\frac{q_0 rd}{(q_0-1)[d+r(d-2)]} \right)^{1+\frac{rd}{2r-d}} (R - R_\infty)^2 |B_R|^{\frac{2}{2^*}}, & q_0 > 1, \\ \left(\frac{S_2^2}{2} \right)^{\frac{rd}{2r-d}} \frac{2r-d}{rd} \left(\frac{(q_0+1)rd}{q_0[d+r(d-2)]} \right)^{1+\frac{rd}{2r-d}} (R - R_\infty)^2 |B_R|^{\frac{2}{2^*}}, & 0 < q_0 \leq 1. \end{cases} \quad (3.53)$$

Remark. In other words, we have the interior estimate $u(x) = O(d(x)^{-\frac{d}{q_0}})$, where $d(x)$ is distance to the boundary.

Proof. The upper bounds (3.50) of Theorem 3.5 can be rewritten as

$$\|u\|_{\infty, r} \leq \frac{K_q^{(3)}[b]}{(R-r)^{\frac{d}{q}}} \|u\|_{q, R} \quad (3.54)$$

for any $q > 1$ and $R_\infty \leq r < R \leq R_0$, where $K_q^{(3)}[b]$ is given by (3.38). It is clear that inequality (3.54) guarantees that we can use Lemma 3.7 with $0 < q = q < +\infty = \bar{q}$, $\gamma = d/q > 1$, $K = K_q^{(3)}[b]$ and for any $R_\infty \leq r < R \leq R_0$. Then we have that for all $0 < q_0 \leq \bar{q} = q$

$$\begin{aligned} \|u\|_{\infty, R_\infty} &\leq 3 \cdot 2^{\frac{q-q_0}{q_0}} \left[\left(4 \frac{d}{q} \frac{q}{q_0} \right)^{\frac{d}{q}} \frac{K_q^{(3)}[b]}{(R_0 - R_\infty)^{\frac{d}{q}}} \right]^{\frac{q}{q_0}} \|u\|_{q_0, R_0} \\ &= 3 \cdot 2^{\frac{2d+1}{q_0}} \left(\frac{d}{q_0} \right)^{\frac{d}{q_0}} \frac{K_{q_0+1}^{(3)}[b]^{\frac{q_0+1}{q_0}}}{(R_0 - R_\infty)^{\frac{d}{q_0}}} \|u\|_{q_0, R_0} \end{aligned}$$

since we can always choose $q = q_0 + 1 > 1$. Finally we notice that we can rewrite the upper bound for all $q_0 > 0$ in the following form:

$$\|u\|_{\infty, r} \leq \frac{A_{q_0}^{(1)}}{(R_0 - R_\infty)^{\frac{d}{q_0}}} \left[A_{q_0}^{(2)} + A_{q_0}^{(3)} \|b\|_{L^r(B_{R_0})}^{\frac{rd}{2r-d}} \right]^{\frac{d}{2q_0}} \|u\|_{q_0, R}$$

where $A_q^{(j)}$ are as in (3.51), (3.52) and (3.53) respectively. \square

The above Theorem has the following important consequence, when applied to the equation $-\Delta u = \lambda u^p$.

Theorem 3.9 (Local Upper bounds, second form). *Consider a nonnegative weak subsolution u to $-\Delta u = \lambda u^p$ on B_R , with $\lambda > 0$, $1 < p < p_s = 2^* - 1 = (d+2)/(d-2)$. Let $0 < R_\infty < R_0 < R$. If $u \in L^{\bar{r}}(B_{R_0})$ with $\bar{r} > d(p-1)/2 := \bar{q}$ then the following bound holds true for any $q_0 > 0$*

$$\|u\|_{\infty, r} \leq \frac{A_{q_0}^{(1)}}{(R_0 - R_\infty)^{\frac{d}{q_0}}} \left[A_{q_0}^{(2)} + A_{q_0}^{(3)} \lambda^{\frac{d(p-1)}{2\bar{r}-d(p-1)}} \|u\|_{L^{\bar{r}}(B_{R_0})}^{\frac{d(p-1)\bar{r}}{2\bar{r}-d(p-1)}} \right]^{\frac{d}{2q_0}} \|u\|_{q_0, R} \quad (3.55)$$

where $A_q^{(j)}$ are as in (3.51), (3.52) and (3.53) respectively.

Proof. Since u is a subsolution to $-\Delta u = \lambda u^p = bu$ with $b = \lambda u^{p-1}$, we need to assume that $u^{p-1} \in L^r$ with $r > d/2$, which amounts to require $u \in L^{\bar{r}}$ with $\bar{r} = r(p-1) > d(p-1)/2$, so that

$$\|b\|_{L^r(B_{R_0})}^{\frac{rd}{2r-d}} = \left(\lambda \int_{B_{R_0}} u^{r(p-1)} dx \right)^{\frac{d}{2r-d}} = \lambda^{\frac{d(p-1)}{2\bar{r}-d(p-1)}} \|u\|_{L^{\bar{r}}(B_{R_0})}^{\frac{d(p-1)\bar{r}}{2\bar{r}-d(p-1)}}$$

Finally, we can apply the bounds of Theorem 3.8 to get the bounds (3.55) with the constants written above. \square

4. Lower bounds

The lower bounds for nonnegative supersolutions can be obtained in two steps: first we perform a Moser iteration, then we need reverse Hölder inequalities, which are a consequence of the celebrated John-Nirenberg Lemma.

4.1. A short reminder about the spaces $M^p(\Omega)$.

We recall here some basic definitions and properties of suitable functional spaces, that will be used in the sequel. We omit the proofs, but we give appropriate references.

We say that a measurable function on $\Omega \subseteq \mathbb{R}^d$ belong to the space $M^p(\Omega)$ if and only if there exists a constant $K \geq 0$ such that

$$\int_{\Omega \cap B_R(x_0)} |f| dx \leq KR^{\frac{d(p-1)}{p}} \quad \text{for all } B_R(x_0),$$

and we define the norm on $M^p(\Omega)$ as follows

$$\|f\|_{M^p(\Omega)} = \inf \left\{ K > 0 : \int_{\Omega \cap B_R(x_0)} |f| \, dx \leq KR^{\frac{d(p-1)}{p}} \quad \text{for all } B_R(x_0) \right\}.$$

One can easily check the strict inclusion $L^p(\Omega) \subset M^p(\Omega)$ for all $1 < p < \infty$, and when Ω is bounded, the equalities $L^1(\Omega) = M^1(\Omega)$ and $L^\infty(\Omega) = M^\infty(\Omega)$. Moreover it is easy to check that when Ω is bounded one has:

$$\|f\|_{L^1(\Omega)} \leq \text{diam}(\Omega)^{\frac{d(p-1)}{p}} \|f\|_{M^p(\Omega)}. \quad (4.1)$$

We now proceed with a series of results that relate the M^p norm with the Riesz potential

$$\mathcal{V}_\mu[f](x) := \int_{\Omega} \frac{f(y)}{|x-y|^{d(1-\mu)}} \, dy \quad \text{with } \mu \in (0, 1]. \quad (4.2)$$

We collect hereafter some well known results, whose proof can be found for instance in [22].

Lemma 4.1. *Let \mathcal{V}_μ be defined as above. Then the following holds.*

(i) *The operator \mathcal{V}_μ maps continuously $L^s(\Omega)$ into $L^r(\Omega)$ for any $1 \leq r \leq \infty$ satisfying*

$$0 \leq \frac{1}{s} - \frac{1}{r} < \mu.$$

Moreover, for any $f \in L^p(\Omega)$,

$$\|\mathcal{V}_\mu f\|_r \leq \left(\frac{s(r+1)-r}{s(\mu r+1)-r} \right)^{\frac{s(r+1)-r}{sr}} \omega_d^{1-\mu} |\Omega|^{\frac{s(\mu r+1)-r}{sr}} \|f\|_s.$$

(ii) *Let $f \in M^p(\Omega)$, with $p > 1/\mu \geq 1$. Then*

$$|\mathcal{V}_\mu[f](x)| \leq \frac{p-1}{p\mu-1} \text{diam}(\Omega)^{\frac{d}{p}(p\mu-1)} \|f\|_{M^p(\Omega)}.$$

(iii) A ‘‘potential’’ version of the Morrey inequality. *Let Ω be a convex bounded subset of \mathbb{R}^d . Then for all $f \in W^{1,1}(\Omega)$ the following inequality holds*

$$|f(x) - f_{\Omega'}| \leq \frac{\text{diam}(\Omega)^d}{d|\Omega'|} \left| \mathcal{V}_{\frac{1}{d}}[|\nabla f|](x) \right| \quad (4.3)$$

for any measurable $\Omega' \subseteq \Omega$ with

$$f_{\Omega'} = \int_{\Omega'} f \frac{dx}{|\Omega'|}$$

Proof. Part (i) is exactly Lemma 7.12 of [22], part (ii) is exactly Lemma 7.18 of [22] and part (iii) is exactly Lemma 7.16 of [22]. \square

4.2. The John-Nirenberg Lemma and reverse Hölder inequalities.

The Caccioppoli estimates proved in Corollary 2.3 show that the gradient of the logarithm of the solution belongs to the space $M^d(\Omega)$, see Proposition 4.5 below. Such M^d -regularity then guarantees the validity of the celebrated John-Nirenberg lemma which as a consequence give a reverse Hölder inequality of the form

$$\frac{\|u\|_{q,R_0}}{\|u\|_{-q,R_0}} \leq \kappa_1^{2/q}$$

for some $0 < q < 1$ and some constant κ_1 . We recall that $\|f\|_{r,Q_0} := \left(\int_{Q_0} |f(y)|^r dy\right)^{1/r}$ also for negative values of r , provided the integral is finite.

We need a lemma concerning estimates on the Riesz potential \mathcal{V}_μ defined in (4.2). It is a quantified version of Lemma 7.20 of [22].

Lemma 4.2 (A “potential” version of the Moser-Trudinger imbedding.) *Let $f \in M^p(\Omega)$ with $p > 1$ and suppose $\|f\|_{M^p(\Omega)} \leq K$. Then there exist two constants κ_2 and κ_3 such that*

$$\int_{\Omega} \exp \left[\frac{|\mathcal{V}_{\frac{1}{p}}[f](x)|}{\kappa_2 K} \right] dx \leq \kappa_3. \quad (4.4)$$

One can take

$$\kappa_2 > (p-1)e \quad \text{and} \quad \kappa_3 = |\Omega| + \frac{\text{diam}(\Omega)^d}{\sqrt{2\pi}} \frac{pe\omega_d}{\kappa_2 - (p-1)e}.$$

Proof. Let $q \geq 1$, $\mu = 1/p$ and $g = \mathcal{V}_\mu[f]$. Then

$$|x - y|^{d(\mu-1)} = |x - y|^{\frac{d}{q}(\frac{\mu}{q}-1)} |x - y|^{d(1-\frac{1}{q})(\frac{\mu}{q}+\mu-1)}$$

and by Hölder inequality we obtain

$$|\mathcal{V}_\mu[f]| \leq \left| \mathcal{V}_{\frac{\mu}{q}}[f] \right|^{\frac{1}{q}} \left| \mathcal{V}_{\mu+\frac{\mu}{q}}[f] \right|^{1-\frac{1}{q}}. \quad (4.5)$$

Applying now estimates (i) of Lemma 4.1 with $s = r = 1$, to $\mathcal{V}_{\frac{\mu}{q}}[f]$, we obtain,

$$\begin{aligned} \|\mathcal{V}_{\frac{\mu}{q}}f\|_1 &\leq \frac{q\omega_d^{1-\frac{\mu}{q}}}{\mu} |\Omega|^{\frac{\mu}{q}} \|f\|_1 \leq pq\omega_d^{1-\frac{1}{pq}} |\Omega|^{\frac{1}{pq}} \text{diam}(\Omega)^{\frac{d(p-1)}{p}} \|f\|_{M^p(\Omega)} \\ &\leq pq\omega_d \text{diam}(\Omega)^{d(1-\frac{1}{p}+\frac{1}{pq})} \|f\|_{M^p(\Omega)} \leq pq\omega_d \text{diam}(\Omega)^{d(1-\frac{1}{p}+\frac{1}{pq})} K \end{aligned} \quad (4.6)$$

where we have used inequality (4.1) together with the fact that $|\Omega| \leq \omega_d \text{diam}(\Omega)^d$. Next we apply estimates (ii) of Lemma 4.1 to $\mathcal{V}_{\mu+\frac{\mu}{q}}[f]$ (the operator \mathcal{V}_ν is well-defined on L^1 , if Ω is bounded, for $\nu > 1$ as well) and we

obtain

$$\begin{aligned} \left| \mathcal{V}_{\mu+\frac{\mu}{q}}[f](x) \right| &\leq \frac{p-1}{p\left(\mu+\frac{\mu}{q}\right)-1} \text{diam}(\Omega)^{\frac{d}{p}\left(p\left(\mu+\frac{\mu}{q}\right)-1\right)} \|f\|_{M^p(\Omega)} \\ &\leq q(p-1) \text{diam}(\Omega)^{\frac{d}{pq}} K \end{aligned} \quad (4.7)$$

for all $x \in \Omega$, hence the same bound is valid for the $L^\infty(\Omega)$ -norm, provided $p(\mu+\mu/q) > 1$ which indeed holds true since $\mu = 1/p$. Joining now inequalities (4.5), (4.6) and (4.7), we obtain

$$\|\mathcal{V}_\mu[f]\|_q^q \leq \left\| \mathcal{V}_{\mu+\frac{\mu}{q}}[f] \right\|_{L^\infty}^{q-1} \left\| \mathcal{V}_{\frac{\mu}{q}}[f](x) \right\|_{L^1(\Omega)} \leq \frac{p\omega_d}{p-1} [(p-1)Kq]^q \text{diam}(\Omega)^d$$

Letting now $1 \leq q = k \in \mathbb{N}$ we get, for k_2 as in the statement,

$$\begin{aligned} \int_\Omega \sum_{k=1}^{\infty} \frac{|g|^k}{k!(\kappa_2 K)^k} dx &\leq \frac{p\omega_d}{p-1} \text{diam}(\Omega)^d \sum_{k=1}^{\infty} \frac{[(p-1)Kk]^k}{k!(\kappa_2 K)^k} \\ &\leq \frac{p\omega_d}{p-1} \text{diam}(\Omega)^d \sum_{k=1}^{\infty} \left[\frac{p-1}{\kappa_2} \right]^k \frac{k^k}{k!} \\ &\leq \frac{p\omega_d}{p-1} \text{diam}(\Omega)^d \sum_{k=1}^{\infty} \left[\frac{(p-1)e}{\kappa_2} \right]^k \frac{1}{\sqrt{2\pi k}} \\ &\leq \frac{p\omega_d}{p-1} \frac{\text{diam}(\Omega)^d}{\sqrt{2\pi}} \frac{(p-1)e}{\kappa_2 - (p-1)e} \\ &= \frac{\text{diam}(\Omega)^d}{\sqrt{2\pi}} \frac{pe\omega_d}{\kappa_2 - (p-1)e} \end{aligned}$$

we have used Stirling's formula:

$$n! = \sqrt{2\pi n} \left[\frac{n}{e} \right]^n e^{\alpha_n} \quad \text{with} \quad \frac{1}{12n+1} \leq \alpha_n \leq \frac{1}{12n}. \quad \square \quad (4.8)$$

We prove hereafter a simplified but quantitative version of the celebrated John-Nirenberg Lemma, which holds in convex domains. Indeed we will use it only on balls and in such case the constants simplify a bit.

Lemma 4.3 (John-Nirenberg). *Let $f \in W^{1,1}(\Omega)$ where Ω is convex, and suppose there exists a constant K such that*

$$\int_{B_R \cap \Omega} |\nabla f| dx \leq K R^{d-1} \quad \text{for all balls } B_R$$

Then the following inequality holds true

$$\int_\Omega \exp \left[\frac{|f - f_\Omega|}{\kappa_0 K} \right] dx \leq \kappa_1 \quad (4.9)$$

where for any $\kappa_2 > (d-1)e$

$$\kappa_0 = \frac{d|\Omega|}{\text{diam}(\Omega)^d \kappa_2} \quad \kappa_1 = \frac{\omega_d \text{diam}(\Omega)^d (\kappa_2 + e)}{\kappa_2 - (d-1)e} \quad \text{and} \quad f_\Omega = \int_\Omega f \frac{dx}{|\Omega|}.$$

Proof. The proof relies on the previous Lemma 4.2 in the special case $p = d$. Indeed inequality (4.4) in that case takes the form

$$\int_{\Omega} \exp \left[\frac{\left| \mathcal{V}_{\frac{1}{d}}[|\nabla f|](x) \right|}{\kappa_2 K} \right] dx \leq \frac{\text{diam}(\Omega)^d}{\sqrt{2\pi}} \frac{de\omega_d}{\kappa_2 - (d-1)e} + |\Omega| \leq \kappa_3 \quad (4.10)$$

where

$$\kappa_2 > (d-1)e,$$

$$\kappa_3 = \omega_d \text{diam}(\Omega)^d \left[\frac{de}{\kappa_2 - (d-1)e} + 1 \right] = \frac{\omega_d \text{diam}(\Omega)^d (\kappa_2 + e)}{\kappa_2 - (d-1)e}.$$

We combine this latter inequality with inequality (4.3) (which requires convexity of the domain) with $\Omega' = \Omega$ and $|\nabla f| \in M^d(\Omega)$. \square

The John-Nirenberg Lemma has an important consequence when applied to $f = \log(u + \delta)$:

Proposition 4.4 (Reverse Hölder inequalities). *Let $\delta \geq 0$ and u be a positive measurable function such that $\log(u + \delta) \in W^{1,1}(\Omega)$, where Ω is convex, and suppose there exists a constant K such that*

$$\int_{B_R \cap \Omega} |\nabla \log(u + \delta)| dx \leq K R^{d-1} \quad \text{for all balls } B_R. \quad (4.11)$$

Then the following inequality

$$\frac{\|u + \delta\|_{q,\Omega}}{\|u + \delta\|_{-q,\Omega}} \leq \kappa_1^{2/q} \quad \text{holds true for any} \quad 0 < q \leq \frac{1}{\kappa_0 K} \quad (4.12)$$

where the constants κ_i are given in Lemma 4.3.

Proof. Let $\delta > 0$. The validity of (4.11) for u entails the validity of the same inequality for $u + \delta$. Notice now that

$$\frac{\|u + \delta\|_{q,\Omega}}{\|u + \delta\|_{-q,\Omega}} \leq \kappa \quad \iff \quad \left(\int_{\Omega} (u + \delta)^q dx \right) \left(\int_{\Omega} (u + \delta)^{-q} dx \right) \leq \kappa^q$$

Then, letting $f = \log(u + \delta)$:

$$\begin{aligned} & \left(\int_{\Omega} (u + \delta)^q dx \right) \left(\int_{\Omega} (u + \delta)^{-q} dx \right) \\ &= \left(\int_{\Omega} e^{[q \log(u + \delta)]} dx \right) \left(\int_{\Omega} e^{[-q \log(u + \delta)]} dx \right) \\ &= \left(\int_{\Omega} e^{qf} dx \right) \left(\int_{\Omega} e^{-qf} dx \right) = \left(\int_{\Omega} e^{q(f - f_{\Omega})} dx \right) \left(\int_{\Omega} e^{-q(f - f_{\Omega})} dx \right) \\ &\leq \left(\int_{\Omega} e^{q|f - f_{\Omega}|} dx \right)^2 \leq \kappa_1^2 \end{aligned}$$

where we used (4.9) for $f = \log(u + \delta)$, and have assumed $q \leq 1/(\kappa_0 K)$ in order to ensure its validity. The case $\delta = 0$ is also true, just by taking the limit $\delta \rightarrow 0$. \square

We conclude this section by showing that reverse Hölder inequalities holds for local supersolutions to our problem, as a consequence of Caccioppoli estimates.

Proposition 4.5 (Reverse Hölder inequalities for supersolutions). *Let $\Omega \subset \mathbb{R}^d$ and let $\lambda > 0$. Let u be a local weak supersolution to $-\Delta u = \lambda u^p$, with $1 \leq p < p_s = 2^* - 1 = (d+2)/(d-2)$. Then for any $\varepsilon > 0$ the following inequality holds true for any $\delta \geq 0$*

$$\left[\frac{\varepsilon}{2^d (e d + \varepsilon)} \right]^{2/q} \frac{\|u + \delta\|_{q, R_0}}{|B_{R_0}|^{1/q}} \leq \frac{\|u + \delta\|_{-q, R_0}}{|B_{R_0}|^{-1/q}}, \quad 0 < q \leq \frac{2^{\frac{d-3}{2}}}{d\omega_d^2 [e(d-1) + \varepsilon]}.$$

Proof. The Caccioppoli estimates (2.2) with R_0 replaced by $2r$ and R replaced by r imply the hypothesis of the above Lemma, in fact:

$$\begin{aligned} \int_{B_r \cap B_{R_0}} |\nabla \log(u + \delta)| \, dx &\leq \int_{B_r} |\nabla \log(u + \delta)| \, dx \\ &\leq |B_r|^{\frac{1}{2}} \left[\int_{B_r} |\nabla \log(u + \delta)|^2 \, dx \right]^{\frac{1}{2}} \leq 2^{\frac{d+3}{2}} \omega_d r^{d-1} \\ &:= K r^{d-1}. \end{aligned} \tag{4.13}$$

Therefore putting $K = 2^{\frac{d+3}{2}} \omega_d$, taking an $\varepsilon > 0$ and choosing $\kappa_2 = e(d-1) + \varepsilon$, we obtain that

$$\frac{1}{\kappa_0 K} = \frac{2^{\frac{d-3}{2}}}{d\omega_d^2 [e(d-1) + \varepsilon]}, \quad \kappa_1 = 2^d \omega_d R_0^d \frac{\varepsilon + e d}{\varepsilon} = |B_{R_0}| 2^d \frac{\varepsilon + e d}{\varepsilon}. \quad \square$$

4.3. Lower Moser iteration

Now we are ready to run the Moser iteration to obtain quantitative local lower bounds in the form:

Theorem 4.6 (Local Lower Estimates). *Let $\Omega \subseteq \mathbb{R}^d$ and let $\lambda > 0$. Let u be a nonnegative local weak supersolution in $B_{R_0} \subseteq \Omega$ to $-\Delta u = \lambda u^p$, with $0 \leq p < p_s = 2^* - 1 = (d+2)/(d-2)$. Then for any $\varepsilon > 0$ and for any*


$$0 < \underline{q} \leq \frac{2^{\frac{d-3}{2}}}{d\omega_d^2 [e(d-1) + \varepsilon]} = q_0 \tag{4.14}$$

the following bound holds true

$$\inf_{x \in B_{R_\infty}} u(x) = \|u\|_{-\infty, R_\infty} \geq I_{-\infty, \underline{q}} \frac{\|u\|_{\underline{q}, R_0}}{|B_{R_0}|^{\frac{1}{\underline{q}}}}. \tag{4.15}$$

where

$$I_{-\infty, \underline{q}} = \left[2^d \mathcal{S}_2^2 \left(\frac{dR_0^2}{(R_0 - R_\infty)^2} + \frac{R_0^2}{R_\infty^2} \right) \right]^{-\frac{d}{2\underline{q}}} \left[\frac{\varepsilon}{2^d (e d + \varepsilon) \sqrt{\omega_d}} \right]^{\frac{2}{\underline{q}}}. \tag{4.16}$$



Upper figure: Plot of $q_0(d)$ defined in (4.14), for $1 \leq d \leq 16$, with $\varepsilon = 0.1$. Lower figure: Zoom for the plot of the same $q_0(x)$ near its minimum that lies in $(5, 6)$.

Remark. One can see that when the dimension d is sufficiently low one has $q_0 < 1$ whereas $q_0 > 1$ in higher dimensions. Notice also that the equality $\inf_{x \in B_{R_\infty}} u(x) = \|u\|_{-\infty, R_\infty}$ holds since u is nonnegative.

Proof. The proof is divided in two steps. We always consider a local supersolution u of $-\Delta u \geq \lambda u^p$.

• **STEP 1.** In this step we consider $\alpha < 0$, and we want to prove $L^{-\alpha} - L^{-\infty}$ local estimates via Moser iteration. The the energy inequality (2.4) for $\alpha < -1$ and $\delta > 0$ gives the estimate

$$\begin{aligned} \int_{\Omega} |\nabla \left((u + \delta)^{\frac{\alpha+1}{2}} \right)|^2 \varphi \, dx &\leq \frac{\lambda(\alpha + 1)^2}{4\alpha} \int_{\Omega} u^p (u + \delta)^\alpha \varphi \, dx \\ &+ \frac{\alpha + 1}{4\alpha} \int_{\Omega} (u + \delta)^{\alpha+1} \Delta \varphi \, dx \\ &\leq \frac{\alpha + 1}{4\alpha} \int_{\Omega} (u + \delta)^{\alpha+1} |\Delta \varphi| \, dx \end{aligned} \tag{4.17}$$

Applying now the Sobolev inequality (3.1) on the ball B_{R_1} and the properties of the test function φ defined in Lemma 2.2, one gets

$$\begin{aligned}
& \left[\int_{B_{R_1}} (u + \delta)^{\frac{2^*}{2}(\alpha+1)} dx \right]^{\frac{2}{2^*}} \\
& \leq \mathcal{S}_2^2 \left(\int_{B_{R_1}} |\nabla(u + \delta)^{\frac{\alpha+1}{2}}|^2 dx + \frac{1}{R_1^2} \int_{B_{R_1}} (u + \delta)^{\alpha+1} dx \right) \\
& \leq \mathcal{S}_2^2 \left(\int_{\Omega} |\nabla(u + \delta)^{\frac{\alpha+1}{2}} \varphi|^2 dx + \frac{1}{R_1^2} \int_{B_{R_1}} (u + \delta)^{\alpha+1} dx \right) \\
& \leq \mathcal{S}_2^2 \left(\frac{\alpha+1}{4\alpha} \int_{\Omega} (u + \delta)^{\alpha+1} |\Delta \varphi| dx + \frac{1}{R_1^2} \int_{B_{R_1}} (u + \delta)^{\alpha+1} dx \right) \\
& \leq \mathcal{S}_2^2 \left(\frac{\alpha+1}{4\alpha} \|\Delta \varphi\|_{\infty} + \frac{1}{R_1^2} \right) \int_{B_{R_0}} (u + \delta)^{\alpha+1} dx \\
& \leq \mathcal{S}_2^2 \left(\frac{d}{(R_0 - R_1)^2} + \frac{1}{R_1^2} \right) \int_{B_{R_0}} (u + \delta)^{\alpha+1} dx
\end{aligned} \tag{4.18}$$

Let, for a given $\gamma_0 < 0$, $\gamma_n := \left[\frac{2^*}{2} \right]^n \gamma_0$ so that $\gamma_n = \frac{2^*}{2} \gamma_{n-1}$. Notice that $\gamma_n \rightarrow -\infty$ monotonically. Consider the above inequality for $\alpha = \alpha_n$ and let $\alpha_n + 1 = \gamma_{n-1}$ so that

$$\begin{aligned}
\|u + \delta\|_{\gamma_n, R_n} &= \|u + \delta\|_{\frac{2^*}{2} \gamma_{n-1}, R_n} = \left[\int_{B_{R_n}} (u + \delta)^{\frac{2^*}{2} \gamma_{n-1}} dx \right]^{\frac{2}{2^* \gamma_{n-1}}} \\
&\geq \left[\mathcal{S}_2^2 \left(\frac{d}{(R_{n-1} - R_n)^2} + \frac{1}{R_n^2} \right) \right]^{\frac{1}{\gamma_{n-1}}} \left[\int_{B_{R_{n-1}}} (u + \delta)^{\gamma_{n-1}} dx \right]^{\frac{1}{\gamma_{n-1}}} \\
&\geq \left[\mathcal{S}_2^2 \left(\frac{d}{(R_{n-1} - R_n)^2} + \frac{1}{R_n^2} \right) \right]^{\frac{1}{\gamma_{n-1}}} \|u + \delta\|_{\gamma_{n-1}, R_{n-1}} \\
&:= I_n^{\frac{1}{\gamma_{n-1}}} \|u + \delta\|_{\gamma_{n-1}, R_{n-1}}
\end{aligned} \tag{4.19}$$

Hence, iterating the above inequality:

$$\|u + \delta\|_{\gamma_n, R_n} \geq I_n^{\frac{1}{\gamma_{n-1}}} I_{n-1}^{\frac{1}{\gamma_{n-2}}} \dots I_1^{\frac{1}{\gamma_0}} \|u + \delta\|_{\gamma_0, R_0} = \prod_{k=1}^n I_k^{\frac{1}{\gamma_{k-1}}} \|u + \delta\|_{\gamma_0, R_0} \tag{4.20}$$

where we have chosen $0 < R_{\infty} < \dots < R_{n+1} < R_n < \dots < R_0$ such that

$$\sum_{k=1}^{\infty} (R_{k-1} - R_k) = R_0 - R_{\infty} \quad \text{and} \quad R_{k-1} - R_k = \frac{R_0 - R_{\infty}}{2^k}$$

so that

$$I_k = \mathcal{S}_2^2 \left(\frac{d}{(R_{n-1} - R_n)^2} + \frac{1}{R_n^2} \right) \leq \mathcal{S}_2^2 \left(\frac{d}{(R_0 - R_{\infty})^2} + \frac{1}{R_{\infty}^2} \right) 4^k := I_0 4^k$$

and

$$\begin{aligned}
\prod_{k=1}^n I_k^{\frac{1}{\gamma_{k-1}}} &= \exp \left[\sum_{k=1}^n \frac{1}{\gamma_{k-1}} \log I_k \right] = \exp \left[\frac{2^*}{2\gamma_0} \sum_{k=1}^n \left[\frac{2}{2^*} \right]^k \log I_k \right] \\
&= \exp \left[\frac{2^*}{2\gamma_0} \sum_{k=1}^n \left[\frac{2}{2^*} \right]^k \log I_0 + \frac{2^* \log 4}{2\gamma_0} \sum_{k=1}^n k \left[\frac{2}{2^*} \right]^k \right] \\
&\geq I_0^{\frac{2^*}{2\gamma_0} \sum_{k=1}^n \left[\frac{2}{2^*} \right]^k} 4^{\frac{d^2}{4\gamma_0}}.
\end{aligned}$$

Taking limits we obtain

$$\prod_{k=1}^{\infty} I_k^{\frac{1}{\gamma_{k-1}}} \geq I_0^{\frac{2^*}{2\gamma_0} \frac{d-2}{2}} 4^{\frac{d^2}{4\gamma_0}} = (2^d I_0)^{\frac{d}{2\gamma_0}}.$$

We can now take the limit in (4.20) to get for any $\gamma_0 < 0$:

$$\begin{aligned}
\|u + \delta\|_{-\infty, R_\infty} &\geq \prod_{k=1}^{\infty} I_k^{\frac{1}{\gamma_{k-1}}} \|u + \delta\|_{\gamma_0, R_0} \geq (2^d I_0)^{\frac{d}{2\gamma_0}} \|u + \delta\|_{\gamma_0, R_0} \\
&= \left[2^d \mathcal{S}_2^2 \left(\frac{d}{(R_0 - R_\infty)^2} + 1 \right) \right]^{\frac{d}{2\gamma_0}} \|u + \delta\|_{\gamma_0, R_0}.
\end{aligned} \tag{4.21}$$

Now we need some Reverse Hölder inequalities, which is the subject of the next step.

• **STEP 2. Reverse Hölder inequalities.** The John-Nirenberg lemma implies reverse Hölder inequalities for super-solutions, in the form of Proposition 4.5: for any $\varepsilon > 0$ the following inequality holds true

$$\left[\frac{\varepsilon}{2^d (e d + \varepsilon)} \right]^{\frac{2}{\underline{q}}} \frac{\|u + \delta\|_{\underline{q}, R_0}}{|B_{R_0}|^{\frac{2}{\underline{q}}}} \leq \|u + \delta\|_{-\underline{q}, R_0}, \quad 0 < \underline{q} \leq \frac{2^{\frac{d-3}{2}}}{d\omega_d^2 [e(d-1) + \varepsilon]}. \tag{4.22}$$

Joining inequality (4.21) and (4.22) and letting $\gamma_0 = -\underline{q}$ with \underline{q} as in (4.22) we obtain

$$\begin{aligned}
\|u + \delta\|_{-\infty, R_\infty} &\geq \left[2^d \mathcal{S}_2^2 \left(\frac{d}{(R_0 - R_\infty)^2} + \frac{1}{R_\infty^2} \right) \right]^{-\frac{d}{2\underline{q}}} \|u + \delta\|_{-\underline{q}, R_0} \\
&\geq \left[2^d \mathcal{S}_2^2 \left(\frac{d}{(R_0 - R_\infty)^2} + \frac{1}{R_\infty^2} \right) \right]^{-\frac{d}{2\underline{q}}} \left[\frac{\varepsilon}{2^d (e d + \varepsilon)} \right]^{\frac{2}{\underline{q}}} \frac{\|u + \delta\|_{\underline{q}, R_0}}{|B_{R_0}|^{\frac{2}{\underline{q}}}} \\
&= \left[2^d \mathcal{S}_2^2 \left(\frac{dR_0^2}{(R_0 - R_\infty)^2} + \frac{R_0^2}{R_\infty^2} \right) \right]^{-\frac{d}{2\underline{q}}} \left[\frac{\varepsilon}{2^d (e d + \varepsilon) \sqrt{\omega_d}} \right]^{\frac{2}{\underline{q}}} \frac{\|u + \delta\|_{\underline{q}, R_0}}{|B_{R_0}|^{\frac{1}{\underline{q}}}} \\
&:= I_{-\infty, \underline{q}} \frac{\|u + \delta\|_{\underline{q}, R_0}}{|B_{R_0}|^{\frac{1}{\underline{q}}}}.
\end{aligned} \tag{4.23}$$

Finally we observe that we can let $\delta \rightarrow 0^+$, and obtain the desired result. \square

4.4. Reverse Hölder inequalities and lower bounds when $1 < p < p_c$

In this section we will first prove more quantitative *reverse Hölder inequalities*, when $p > 1$. We have obtained a reverse smoothing effect from L^q to $L^{-\infty}$, for a suitable explicit q which may be close to zero, if we seek for a bound valid for any dimension. In order to be able to join local upper and lower estimates to get a clean form of Harnack inequality, we need to reach those values of q which are above $d(p-1)/2$, and this is possible only when $1 < p < p_c = d/(d-2)$.

Proposition 4.7 (Reverse Hölder inequalities for $1 < p < p_c$). *Let $\Omega \subseteq \mathbb{R}^d$ and let $\lambda > 0$. Let u be a nonnegative local weak supersolution in Ω to $-\Delta u = \lambda u^p$, with $1 < p < p_c = d/(d-2)$. Let $B_{\bar{R}} \subset B_{R_0} \subset \Omega$. Then we have that*

$$\frac{\|u\|_{\bar{q}, \bar{R}}}{|B_{\bar{R}}|^{\frac{1}{\bar{q}}}} \leq I_{\bar{q}, q_0} \frac{\|u\|_{q_0, R_0}}{|B_{R_0}|^{\frac{1}{q_0}}} \quad \forall q_0 \in (0, \bar{q}], \quad d(p-1)/2 < \bar{q} < d/(d-2) \quad (4.24)$$

where, if $\frac{d-2}{d}\bar{q} \leq q_0 \leq \bar{q}$

$$I_{\bar{q}, q_0} := \left[\frac{2d\bar{q}\mathcal{S}_2^2}{(2^* - 2\bar{q})} + \mathcal{S}_2^2 \frac{(R_0 - \bar{R})^2}{\bar{R}^2} \right]^{\frac{2^*}{2\bar{q}}} \left[\frac{\omega_d^{1/d} R_0}{R_0 - \bar{R}} \right]^{\frac{2^*}{\bar{q}}} \left[\frac{R_0}{\bar{R}} \right]^{\frac{d}{\bar{q}}}$$

whereas if $0 < q_0 < \frac{d-2}{d}\bar{q}$

$$I_{\bar{q}, q_0} := 3 \cdot 2^{\frac{(d-2)\bar{q}}{2q_0} - \frac{d}{2}} \left[\frac{2d\bar{q}\mathcal{S}_2^2}{(2^* - 2\bar{q})} \frac{\bar{R}^2}{(R_0 - \bar{R})^2} + \mathcal{S}_2^2 \right]^{\frac{\bar{q}-q_0}{\bar{q}q_0} \frac{d}{2}} \\ \times \left(4\omega_d^{\frac{1}{d}} \frac{\bar{q} - q_0}{q_0\bar{q}} \right)^{\frac{d}{q_0} - \frac{d}{\bar{q}}} \left[\frac{\bar{R}}{R_0} \right]^{\frac{d}{q_0}}.$$

Proof. Consider the energy identity for supersolutions with $-1 < \alpha < 0$ (we can take $\delta = 0$ in such a range of α), which gives the following estimate for any positive test function $\varphi \in C_0^2(\Omega)$ with $\nabla\varphi \equiv 0$ on $\partial\Omega$:

$$\frac{4|\alpha|}{(\alpha+1)^2} \int_{\Omega} |\nabla u^{\frac{\alpha+1}{2}}|^2 \varphi \, dx + \lambda \int_{\Omega} u^{p+\alpha} \varphi \, dx \leq \frac{1}{|\alpha+1|} \int_{\Omega} u^{\alpha+1} |\Delta\varphi| \, dx \quad (4.25)$$

that implies, using the test function φ of Lemma 2.2 with $R_{\infty} < R_0$

$$\int_{B_{R_{\infty}}} |\nabla u^{\frac{\alpha+1}{2}}|^2 \, dx \leq \frac{d|\alpha+1|}{|\alpha|(R_0 - R_{\infty})^2} \int_{B_{R_0}} u^{\alpha+1} \, dx \quad (4.26)$$

Applying now the Sobolev inequality (3.26) on the ball $B_{R_{\infty}}$ we arrive at

$$\left[\int_{B_{R_{\infty}}} u^{\frac{2^*}{2}(\alpha+1)} \, dx \right]^{\frac{2}{2^*}} \leq \mathcal{S}_2^2 \left[\frac{d|\alpha+1|}{|\alpha|(R_0 - R_{\infty})^2} + \frac{1}{R_{\infty}^2} \right] \int_{B_{R_0}} u^{\alpha+1} \, dx$$

Letting now $0 < \alpha + 1 = \beta < 1$ we get

$$\begin{aligned} \left[\int_{B_{R_\infty}} u^{\frac{2^*}{2}\beta} dx \right]^{\frac{2}{2^*\beta}} &\leq \left[\frac{\mathcal{S}_2}{R_0 - R_\infty} \right]^{\frac{2}{\beta}} \left[\frac{d|\beta|}{(1-\beta)} + \frac{(R_0 - R_\infty)^2}{R_\infty^2} \right]^{\frac{1}{\beta}} \\ &\times \left[\int_{B_{R_0}} u^\beta dx \right]^{\frac{1}{\beta}}. \end{aligned} \quad (4.27)$$

Choosing $\beta > (d-2)(p-1)/2$ is compatible with $\beta < 1$, if and only if $p < d/(d-2) = p_c$ and this is the point where the well known Serrin's exponent p_c enters. We now let $d(p-1)/2 < \bar{q} = 2^*\beta/2 < 2^*/2$ and we see that (4.27) implies

$$\begin{aligned} \|u\|_{\bar{q},r} &\leq \left[\frac{2d\bar{q}\mathcal{S}_2^2}{(2^* - 2\bar{q})} + \mathcal{S}_2^2 \frac{(R-r)^2}{r^2} \right]^{\frac{2^*}{2\bar{q}}} \frac{\|u\|_{\frac{2^*}{2}\bar{q},R}}{(R-r)^{\frac{2^*}{\bar{q}}}} \\ &\leq \left[\frac{2d\bar{q}\mathcal{S}_2^2}{(2^* - 2\bar{q})} + \mathcal{S}_2^2 \frac{(R_0 - R_\infty)^2}{R_\infty^2} \right]^{\frac{2^*}{2\bar{q}}} \frac{\|u\|_{\frac{2^*}{2}\bar{q},R}}{(R-r)^{\frac{2^*}{\bar{q}}}} \end{aligned} \quad (4.28)$$

for any $R_\infty \leq r < R \leq R_0$. Let $\underline{q} = 2\bar{q}/2^* < \bar{q}$. We consider separately the case $\underline{q} \leq q_0 \leq \bar{q}$ and the case $0 < q_0 < \underline{q} < \bar{q}$. In the first case we can use Hölder inequality in (4.28):

$$\begin{aligned} \|u\|_{\bar{q},r} &\leq \left[\frac{2d\bar{q}\mathcal{S}_2^2}{(2^* - 2\bar{q})} + \mathcal{S}_2^2 \frac{(R_0 - R_\infty)^2}{R_\infty^2} \right]^{\frac{2^*}{2\bar{q}}} \frac{\|u\|_{\frac{2^*}{2}\bar{q},R}}{(R-r)^{\frac{2^*}{\bar{q}}}} \\ &\leq \left[\frac{2d\bar{q}\mathcal{S}_2^2}{(2^* - 2\bar{q})} + \mathcal{S}_2^2 \frac{(R_0 - R_\infty)^2}{R_\infty^2} \right]^{\frac{2^*}{2\bar{q}}} \left[\frac{\omega_d^{1/d}}{R-r} \right]^{\frac{2^*}{\bar{q}}} \frac{|B_R|^{\frac{1}{\bar{q}}}}{|B_R|^{\frac{1}{q_0}}} \|u\|_{q_0,R} \end{aligned}$$

which is (4.24) when $\underline{q} \leq q_0 \leq \bar{q}$, once we let $R = R_0$ and $r = \bar{R}$. On the other hand, when $0 < q_0 < \underline{q} < \bar{q}$, we can use inequality (4.28) rewritten as

$$\|u\|_{\bar{q},r} \leq \left[\frac{2d\bar{q}\mathcal{S}_2^2}{(2^* - 2\bar{q})} + \mathcal{S}_2^2 \frac{(R_0 - R_\infty)^2}{R_\infty^2} \right]^{\frac{2^*}{2\bar{q}}} \frac{\|u\|_{\frac{2^*}{2}\bar{q},R}}{(R-r)^{\frac{2^*}{\bar{q}}}} := \frac{K}{(R-r)^{\frac{2^*}{\bar{q}}}} \|u\|_{\frac{2^*}{2}\bar{q},R} \quad (4.29)$$

so that Lemma 3.7 with $\gamma = 2^*/\bar{q}$ gives that for all $0 < q_0 \leq \underline{q} < \bar{q}$ (recall that $\underline{q} = 2\bar{q}/2^*$)

$$\begin{aligned}
\|u\|_{\bar{q}, R_\infty} &\leq 3 \cdot 2^{\frac{\bar{q}(q_0 - \bar{q})}{q_0(\bar{q} - \underline{q})}} \left[\left(4\gamma \frac{q(\bar{q} - q_0)}{q_0(\bar{q} - \underline{q})} \right)^\gamma \frac{K}{(R_0 - R_\infty)^\gamma} \right]^{\frac{q(\bar{q} - q_0)}{q_0(\bar{q} - \underline{q})}} \|u\|_{q_0, R_0} \\
&= 3 \cdot 2^{\frac{(d-2)\bar{q}}{2q_0} - \frac{d}{2}} \left[\left(4 \frac{2d}{(d-2)\bar{q}} \frac{d-2}{d} \frac{(\bar{q} - q_0)}{\bar{q}} \frac{K}{q_0 \frac{2}{d} \bar{q}} \right)^\gamma \frac{K}{(R_0 - R_\infty)^\gamma} \right]^{\frac{q(\bar{q} - q_0)}{q_0(\bar{q} - \underline{q})}} \\
&\quad \times \|u\|_{q_0, R_0} \\
&= 3 \cdot 2^{\frac{(d-2)\bar{q}}{2q_0} - \frac{d}{2}} K^{\frac{q(\bar{q} - q_0)}{q_0(\bar{q} - \underline{q})}} \left(4d \frac{\bar{q} - q_0}{q_0 \bar{q}} \frac{1}{R_0 - R_\infty} \right)^\gamma \frac{q(\bar{q} - q_0)}{q_0(\bar{q} - \underline{q})} \|u\|_{q_0, R_0} \\
&= 3 \cdot 2^{\frac{(d-2)\bar{q}}{2q_0} - \frac{d}{2}} K^{\frac{\bar{q} - q_0}{q_0} \frac{d-2}{2}} \left(4d \frac{\bar{q} - q_0}{q_0 \bar{q}} \frac{\omega_d^{1/d} R_0}{R_0 - R_\infty} \right)^{\frac{d}{q_0} - \frac{d}{\bar{q}}} \\
&\quad \times \frac{|B_{R_0}|^{\frac{1}{\bar{q}}}}{|B_{R_0}|^{\frac{1}{q_0}}} \|u\|_{q_0, R_0} \\
&= 3 \cdot 2^{\frac{(d-2)\bar{q}}{2q_0} - \frac{d}{2}} \left[\frac{2d\bar{q}\mathcal{S}_2^2}{(2^* - 2\bar{q})} + \mathcal{S}_2^2 \frac{(R_0 - R_\infty)^2}{R_\infty^2} \right]^{\frac{\bar{q} - q_0}{\bar{q} q_0} \frac{d}{2}} \\
&\quad \times \left(4d \frac{\bar{q} - q_0}{q_0 \bar{q}} \frac{\omega_d^{1/d} R_0}{R_0 - R_\infty} \right)^{\frac{d}{q_0} - \frac{d}{\bar{q}}} \left[\frac{R_\infty}{R_0} \right]^{\frac{d}{q_0}} \frac{|B_{R_0}|^{\frac{1}{\bar{q}}}}{|B_{R_\infty}|^{\frac{1}{q_0}}} \|u\|_{q_0, R_0} \\
&= 3 \cdot 2^{\frac{(d-2)\bar{q}}{2q_0} - \frac{d}{2}} \left[\frac{2d\bar{q}\mathcal{S}_2^2}{(2^* - 2\bar{q})} \frac{R_\infty^2}{(R_0 - R_\infty)^2} + \mathcal{S}_2^2 \right]^{\frac{\bar{q} - q_0}{\bar{q} q_0} \frac{d}{2}} \\
&\quad \times \left(4d\omega_d^{\frac{1}{d}} \frac{\bar{q} - q_0}{q_0 \bar{q}} \right)^{\frac{d}{q_0} - \frac{d}{\bar{q}}} \left[\frac{R_\infty}{R_0} \right]^{\frac{d}{q_0}} \frac{|B_{R_0}|^{\frac{1}{\bar{q}}}}{|B_{R_\infty}|^{\frac{1}{q_0}}} \|u\|_{q_0, R_0}
\end{aligned} \tag{4.30}$$

whence the statement follows upon relabeling R_∞ as \bar{R} . \square

As a first consequence of the above inequalities, we can improve the local lower bounds of Theorem 4.6 in this good supercritical range.

Theorem 4.8 (Local Lower Estimates when $1 < p < p_c$). *Let $\Omega \subseteq \mathbb{R}^d$ and let $\lambda > 0$. Let u be a nonnegative local weak supersolution in $B_{R_0} \subseteq \Omega$ to $-\Delta u = \lambda u^p$, with $1 < p < p_c = d/(d-2)$.*

$$\inf_{x \in B_{R_\infty}} u(x) = \|u\|_{-\infty, R_\infty} \geq \frac{I_{-\infty, \underline{q}}}{I_{\bar{q}, \underline{q}}} \frac{\|u\|_{\bar{q}, \bar{R}}}{|B_{\bar{R}}|^{\frac{1}{\bar{q}}}} \text{ with } d(p-1)/2 < \bar{q} < d/(d-2)
\tag{4.31}$$

for any $0 < R_\infty < \bar{R} < R_0$, where $\underline{q} \in (0, q_0 \wedge \bar{q}]$, q_0 and $I_{-\infty, \underline{q}}$ are given in (4.33) and $I_{\bar{q}, \underline{q}}$ is given by (4.35), (4.36).

Proof. We use the local lower bounds of Theorem 4.6 for $\underline{q} \in (0, q_0]$, $\varepsilon = e$, with the definition of q_0 to be recalled below, so that

$$\inf_{x \in B_{R_\infty}} u(x) = \|u\|_{-\infty, R_\infty} \geq I_{-\infty, \underline{q}} \frac{\|u\|_{\underline{q}, R_0}}{|B_{R_0}|^{\frac{1}{\underline{q}}}}. \quad (4.32)$$

where

$$\underline{q} \leq q_0 := \frac{2^{\frac{d-3}{2}}}{d^2 \omega_d^2 e} \quad (4.33)$$

$$I_{-\infty, \underline{q}} = \left[2^d \mathcal{S}_2^2 \left(\frac{dR_0^2}{(R_0 - R_\infty)^2} + \frac{R_0^2}{R_\infty^2} \right) \right]^{-\frac{d}{2\underline{q}}} \left[\frac{e}{2^d e (d+1) \sqrt{\omega_d}} \right]^{\frac{2}{\underline{q}}}.$$

Recall the reverse Hölder inequalities of Proposition 4.7

$$\frac{\|u\|_{\underline{q}, R_0}}{|B_{R_0}|^{\frac{1}{\underline{q}}}} \geq \frac{\|u\|_{\bar{q}, \bar{R}}}{I_{\bar{q}, \underline{q}} |B_{\bar{R}}|^{\frac{1}{\bar{q}}}}, \quad (4.34)$$

valid whenever $0 < \bar{R} < R_0$, $\underline{q} \in (0, \bar{q}]$, $d(p-1)/2 < \bar{q} < d/(d-2)$, where

$$I_{\bar{q}, \underline{q}} := \left[\frac{2d\bar{q}\mathcal{S}_2^2}{(2^* - 2\bar{q})} + \mathcal{S}_2^2 \frac{(R_0 - \bar{R})^2}{\bar{R}^2} \right]^{\frac{2^*}{2\bar{q}}} \left[\frac{\omega_d^{1/d} R_0}{R_0 - \bar{R}} \right]^{\frac{2^*}{\bar{q}}} \left[\frac{R_0}{\bar{R}} \right]^{\frac{d}{\bar{q}}} \quad (4.35)$$

if $\frac{d-2}{d}\bar{q} \leq \underline{q} \leq \bar{q}$,

$$I_{\bar{q}, \underline{q}} = 3 \cdot 2^{\frac{(d-2)\bar{q}}{2\underline{q}} - \frac{d}{2}} \left[\frac{2d\bar{q}\mathcal{S}_2^2}{(2^* - 2\bar{q})} \frac{\bar{R}^2}{(R_0 - \bar{R})^2} + \mathcal{S}_2^2 \right]^{\frac{\bar{q}-\underline{q}}{\bar{q}\underline{q}} \frac{d}{2}} \left(4\omega_d^{\frac{1}{d}} \frac{\bar{q}-\underline{q}}{\underline{q}\bar{q}} \right)^{\frac{d}{\underline{q}} - \frac{d}{\bar{q}}} \left[\frac{R_0}{\bar{R}} \right]^{\frac{d}{\bar{q}}}, \quad (4.36)$$

if $0 < \underline{q} < \frac{d-2}{d}\bar{q}$, with q_0 as in (4.33). Combining inequalities (4.32) and (4.34) we obtain (4.31). \square

Remark. The above lower bounds turn out to be important when applied to solutions, since they will imply directly a clean form of Harnack inequality when $1 < p < p_c$ and then local absolute bounds, which is a novelty and a typical feature of the “good” superlinear case $1 < p < p_c$. We stress the fact that in the upper range $p_c \leq p < p_s$ such absolute bounds can not be true, as explicit counter-examples show. We will give more details on these counterexamples in the next section.

5. Harnack inequalities

In this section we will show in a quantitative way how upper and lower bounds can be joined to form Harnack inequalities for solutions, and to obtain as a consequence absolute local upper ($1 < p < p_c$) and absolute local lower bounds ($0 < p < 1$), which are new, as far as we know. We first join local bounds of Theorems 3.1, 3.9 (upper) and (4.6) (lower), to obtain a general form for Harnack inequalities, which at a first sight appear to be weaker than what expected, because its constant depends on local L^q -norms of the

solution itself. This is the only form of Harnack inequality that can hold for all $0 \leq p < p_s = (d+2)/(d-2)$. To eliminate this quotient and to obtain Harnack inequalities in a more classical form one has to assume that $0 < p < p_c = d/(d-2)$.

This fact might seem puzzling, but there are very weak (distributional) solutions in the range $p_c \leq p < p_s$ that are not bounded, cf. [25, 28, 29, 30, 31], even when one prescribes zero Dirichlet boundary conditions. According to Mazzeo and Pacard [25], in this range there are solutions with a singularity of the type $|x - x_0|^{-2/(p-1)}$ at a point $x_0 \in \Omega$. Such solutions are not locally in L^q with $q > d(p-1)_+/2$ if $p > p_c$, hence the local upper estimate fails for them when applied to a ball that contains the singularity. In this range there appears in a clear form the difference between weak and very weak solutions, which helps understanding these critical exponents. Regarding boundary behaviour, the range to consider is $p_1 \leq p < p_s$, where $p_1 = (d+1)/(d-1)$ is the exponent introduced by Brezis and Turner [7]. In this range there exist very weak solutions which are not weak (energy) solutions and can have a singularity at some points of the boundary and satisfy elsewhere on the boundary the prescribed condition in a suitable trace sense, not necessarily in a continuous fashion, cf. del Pino et al. [15].

Theorem 5.1 (Harnack inequality for $0 \leq p < p_s$). *Let $\Omega \subseteq \mathbb{R}^d$ and let $\lambda > 0$. Let u be a nonnegative local weak solution in $B_{R_0} \subseteq \Omega$ to $-\Delta u = \lambda u^p$, with $0 \leq p < p_s = (d+2)/(d-2)$. Given $R_\infty < R_0$ and $\varepsilon > 0$ we assume*

$$0 < \underline{q} \leq q_0 := \frac{2^{\frac{d-3}{2}}}{d\omega_d^2[e(d-1) + \varepsilon]}, \quad \bar{q} > \frac{d(p-1)_+}{2}. \quad (5.1)$$

If $0 < \bar{q} < d/(d-2)$ we also assume

$$\left[\frac{\log \frac{2^* - d(p-1)_+}{2\bar{q} - d(p-1)_+}}{\log \frac{d}{d-2}} \right] \text{ not integer.}$$

Then the following bound holds true

$$\sup_{x \in B_{R_\infty}} u(x) \leq \mathcal{H}_p[u] \inf_{x \in B_{R_\infty}} u(x) \quad (5.2)$$

where $\mathcal{H}_p[u]$ depends on u through some local norms as follows

$$\begin{aligned} \mathcal{H}_p[u] &= \mathcal{H}_p[u](d, \bar{q}, \underline{q}, \varepsilon, R_0, R_\infty) \\ &= \frac{I_{\infty, \bar{q}}}{I_{-\infty, \underline{q}}} \left(\frac{\left(\int_{B_{R_0}} u^q dx \right)^{\frac{(p-1)_+}{q}}}{\int_{B_{R_\infty}} u^{(p-1)_+} dx} \right)^{\frac{d}{2\bar{q} - d(p-1)_+}} \frac{\left(\int_{B_{R_0}} u^{\bar{q}} dx \right)^{\frac{1}{\bar{q}}}}{\left(\int_{B_{R_0}} u^{\underline{q}} dx \right)^{\frac{1}{\underline{q}}}}. \end{aligned} \quad (5.3)$$

with $I_{\infty, \bar{q}}$ given by (3.5), $I_{-\infty, \underline{q}}$ is given by (4.16).

Proof. We recall the local upper bounds of Theorem 3.1: for any $B_{R_\infty} \subset B_{R_0} \subseteq \Omega$

$$\|u\|_{\infty, R_\infty} \leq I_{\infty, \bar{q}} \left(\frac{\left(\int_{B_{R_0}} u^q dx \right)^{\frac{(p-1)_+}{q}}}{\int_{B_{R_\infty}} u^{(p-1)_+} dx} \right)^{\frac{d}{2\bar{q}-d(p-1)_+}} \frac{\|u\|_{\bar{q}, R_0}}{|B_{R_0}|^{\frac{1}{\bar{q}}}} \quad (5.4)$$

for any $\bar{q} > \frac{d(p-1)_+}{2}$, where $I_{\infty, \bar{q}}$ is given by (3.5) and when $0 < \bar{q} < d/(d-2)$ we require the additional condition (3.7) on \bar{q} . We also recall the lower bounds of Theorem 4.6: for any $\varepsilon > 0$ and for any \underline{q} as in (5.1), the following bound holds true

$$\frac{\inf_{x \in B_{R_\infty}} u(x) |B_{R_0}|^{\frac{1}{\underline{q}}}}{I_{-\infty, \underline{q}} \|u\|_{\underline{q}, R_0}} \geq 1. \quad (5.5)$$

where $I_{-\infty, \underline{q}}$ is given by (4.16). Joining (5.4) and (5.5) gives (5.2). \square

Theorem 5.2 (Harnack inequality, $0 \leq p \leq 1$). *Let $\Omega \subseteq \mathbb{R}^d$ and let $\lambda > 0$. Let u be a nonnegative local weak solution in $B_{R_0} \subseteq \Omega$ to $-\Delta u = \lambda u^p$, with $0 \leq p \leq 1$. For all $R_\infty < R_0$ the following bound holds true*

$$\sup_{x \in B_{R_\infty}} u(x) \leq \mathcal{H}_p \inf_{x \in B_{R_\infty}} u(x)$$

where \mathcal{H}_p does not depend on u , and is given by

$$\begin{aligned} \mathcal{H}_p &= \left[\frac{2^d S_2^4 R_0^2}{(R_0 - R_\infty)^2} \left(\frac{d R_0^2}{(R_0 - R_\infty)^2} + \frac{R_0^2}{R_\infty^2} \right) \right]^{\frac{d}{2q_0}} \\ &\times \left[\frac{2^d \left(\left(\frac{d}{d-2} \right)^{n_0 - \frac{1}{2}} \frac{2^{\frac{d-3}{2}}}{d\omega_d^2} + e \right) \sqrt{\omega_d}}{\left(\frac{d}{d-2} \right)^{n_0 - \frac{1}{2}} \frac{2^{\frac{d-3}{2}}}{d\omega_d^2} - e(d-1)} \right]^{\frac{2}{q_0}} \\ &\times \left\{ \left(\frac{d}{d-2} \right)^d \frac{2(d-2)\sqrt{d}}{(\sqrt{d} - \sqrt{d-2})^3} \right. \\ &\left. \left[\Lambda_p + \frac{d-2}{q_0} + \frac{(R_0 - R_\infty)^2}{R_\infty^2} \max \left\{ \frac{d-2}{(dq_0)^2} |dq_0 - (d-2)|, \frac{1}{4} \right\} \right] \right\}^{\frac{d}{2q_0}} \end{aligned} \quad (5.6)$$

with

$$q_0 = \left(\frac{d-2}{d} \right)^{n_0 - \frac{1}{2}} \quad \text{and} \quad n_0 = i.p. \left[\frac{\log \left(e(d-1) \frac{d\omega_d^2}{2^{\frac{d-3}{2}}} \right)}{\log \frac{d}{d-2}} + \frac{3}{2} \right] \quad (5.7)$$

Proof. The goal of the proof is to simplify the quotient of L^q -norms in the expression of the constant $\mathcal{H}_p[u]$ of the Harnack inequality (5.2). Since we

are dealing with the range $0 \leq p \leq 1$, we can choose

$$0 < \bar{q} = \underline{q} = q_0 = q_0(\varepsilon) := \frac{2^{\frac{d-3}{2}}}{d\omega_d^2[e(d-1) + \varepsilon]},$$

where $\bar{q} > 0$ with $\left\lceil \frac{\log \frac{2^*}{2\bar{q}}}{\log \frac{d}{d-2}} \right\rceil$ not an integer. In fact, we shall arrive, with a suitable choice of the parameter ε , to a value of q_0 smaller than $d/(d-2)$, so that the requirement $\left\lceil \log \frac{2^*}{2\bar{q}} / \log \frac{d}{d-2} \right\rceil$ not being integer is necessary. The latter condition means $q_0(\varepsilon) \neq [(d-2)/d]^n$ for all $n \in \mathbb{N}$, and this is possible since we can always choose ε

$$0 < \varepsilon = \left(\frac{d}{d-2} \right)^{n_0 - \frac{1}{2}} \frac{2^{\frac{d-3}{2}}}{d\omega_d^2} - e(d-1) \quad \text{so that} \quad q_0 = \left(\frac{d-2}{d} \right)^{n_0 - \frac{1}{2}}$$

where n_0 is the first integer n such that $\varepsilon(n) > 0$, which is

$$n_0 = i.p. \left[\frac{\log \left(e(d-1) \frac{d\omega_d^2}{2^{\frac{d-3}{2}}} \right)}{\log \frac{d}{d-2}} + \frac{1}{2} \right] + 1.$$

The constants become in this case

$$\begin{aligned} I_{\infty, \bar{q}} &= \left[\frac{c_1 \mathcal{S}_2^2 R_0^2}{(R_0 - R_\infty)^2} \right]^{\frac{d}{2q_0}} \left\{ \left(\frac{d}{d-2} \right)^d \frac{2(d-2)}{(\sqrt{d} - \sqrt{d-2})^2} \times \right. \\ &\quad \left. \times \left[\Lambda_p + \frac{d-2}{q_0} + \frac{(R_0 - R_\infty)^2}{R_\infty^2} \max \left\{ \frac{d-2}{(dq_0)^2} |dq_0 - (d-2)|, \frac{1}{4} \right\} \right] \right\}^{\frac{d}{2q_0}} \\ &= \left[\frac{\mathcal{S}_2^2 R_0^2}{(R_0 - R_\infty)^2} \right]^{\frac{d}{2q_0}} \left\{ \left(\frac{d}{d-2} \right)^d \frac{2(d-2)\sqrt{d}}{(\sqrt{d} - \sqrt{d-2})^3} \times \right. \\ &\quad \left. \times \left[\Lambda_p + \frac{d-2}{q_0} + \frac{(R_0 - R_\infty)^2}{R_\infty^2} \max \left\{ \frac{d-2}{(dq_0)^2} |dq_0 - (d-2)|, \frac{1}{4} \right\} \right] \right\}^{\frac{d}{2q_0}}, \end{aligned} \quad (5.8)$$

where $\Lambda_p = 2$ if $p \neq 1$, $\Lambda_p = \lambda/4$ if $p = 1$ and, since $q_0 < d/(d-2)$,

$$c_1 := \max_{i=0,1} \frac{q_0 \left(\frac{d}{d-2} \right)^{k_0 - 1 + i}}{\left| q_0 \left(\frac{d}{d-2} \right)^{k_0 - 1 + i} - 1 \right|} = \max_{i=0,1} \frac{\left(\frac{d}{d-2} \right)^{i + \frac{1}{2}}}{\left(\frac{d}{d-2} \right)^{i + \frac{1}{2}} - 1} = \frac{\sqrt{d}}{\sqrt{d} - \sqrt{d-2}} \quad (5.9)$$

since k_0 is given by:

$$k_0 = i.p. \left[\frac{\log \frac{2^*}{2q_0}}{\log \frac{d}{d-2}} \right] = i.p. \left[1 + \frac{\log \frac{1}{q_0}}{\log \frac{d}{d-2}} \right] = i.p. \left[1 + n_0 - \frac{1}{2} \right] = n_0 + 1$$

and the last step in (5.9) follows by an explicit calculation. Moreover $I_{-\infty, \underline{q}}$ given by formula (4.16) takes the form

$$\begin{aligned} I_{-\infty, q_0} &= \left[2^d \mathcal{S}_2^2 \left(\frac{dR_0^2}{(R_0 - R_\infty)^2} + \frac{R_0^2}{R_\infty^2} \right) \right]^{-\frac{d}{2q_0}} \left[\frac{\varepsilon}{2^d (e d + \varepsilon) \sqrt{\omega_d}} \right]^{\frac{2}{q_0}} \\ &= \left[2^d \mathcal{S}_2^2 \left(\frac{dR_0^2}{(R_0 - R_\infty)^2} + \frac{R_0^2}{R_\infty^2} \right) \right]^{-\frac{d}{2q_0}} \\ &\quad \times \left[\frac{\left(\frac{d}{d-2} \right)^{n_0 - \frac{1}{2}} 2^{\frac{d-3}{2}} \frac{2}{d\omega_d^2} - e(d-1)}{2^d \left(\left(\frac{d}{d-2} \right)^{n_0 - \frac{1}{2}} 2^{\frac{d-3}{2}} \frac{2}{d\omega_d^2} + e \right) \sqrt{\omega_d}} \right]^{\frac{2}{q_0}}. \end{aligned}$$

Hence we get the expression of $\mathcal{H}_p = I_{\infty, q_0} / I_{-\infty, q_0}$ given in (5.6). \square

When $p > 1$ we can not join the upper and the lower bound so easily, we need the improved lower bounds of Theorem 4.8, valid only when $p < p_c$.

Theorem 5.3 (Harnack Inequalities when $1 < p < p_c$). *Let $\Omega \subseteq \mathbb{R}^d$ and let $\lambda > 0$. Let u be a nonnegative local weak solution to $-\Delta u = \lambda u^p$ in $B_{R_0} \subseteq \Omega$, with $1 < p < p_c = d/(d-2)$. Then for any $0 < R_\infty < \bar{R} < R_0$ there exists an explicit constant $\mathcal{H}_p > 0$ such that*

$$\sup_{x \in B_{R_\infty}} u(x) \leq \mathcal{H}_p \inf_{x \in B_{R_\infty}} u(x) \quad (5.10)$$

where \mathcal{H}_p does not depend on u , and is given by

$$\mathcal{H}_p = I_{\infty, \bar{q}} \left(\frac{I_{\bar{q}, \underline{q}}}{I_{-\infty, \underline{q}}} \right)^{\frac{2\bar{q}}{2\bar{q} - d(p-1)}}, \quad \text{with} \quad \frac{d(p-1)}{2} < \bar{q} < \frac{d}{d-2} \quad (5.11)$$

where the constants $\underline{q} \in (0, q_0 \wedge \bar{q}]$, q_0 and $I_{-\infty, \underline{q}}$ are given in (4.33), $I_{\bar{q}, \underline{q}}$ is given by (4.35), (4.36), $I_{\infty, \bar{q}}$ is given by (3.5); moreover, since $\bar{q} < d/(d-2)$ we require the additional condition (3.7).

Proof. We first consider the lower bounds of Theorem 4.8. Let $\Omega \subseteq \mathbb{R}^d$ and let $\lambda > 0$. Let u be a nonnegative local weak supersolution in $B_{R_0} \subseteq \Omega$ to $-\Delta u = \lambda u^p$, with $1 < p < p_c = d/(d-2)$. Then

$$\frac{\|u\|_{\bar{q}, \bar{R}}}{|B_{\bar{R}}|^{\frac{1}{\bar{q}}}} \leq \frac{I_{\bar{q}, \underline{q}}}{I_{-\infty, \underline{q}}} \inf_{x \in B_{R_\infty}} u(x) \quad (5.12)$$

for any $0 < R_\infty < \bar{R} < R_0$, where $d(p-1)/2 < \bar{q} < d/(d-2)$, $\underline{q} \in (0, q_0 \wedge \bar{q}]$, q_0 and $I_{-\infty, \underline{q}}$ are given in (4.33) and $I_{\bar{q}, \underline{q}}$ is given by (4.35), (4.36). Then we

recall the upper bounds of Theorem 3.1 which we rewrite as

$$\begin{aligned}
\|u\|_{\infty, R_\infty} &\leq I_{\infty, \bar{q}} \left(\frac{\|u\|_{\bar{q}, \bar{R}}^{(p-1)_+}}{|B_{\bar{R}}|^{\frac{(p-1)_+}{\bar{q}}}} \frac{|B_{R_\infty}|}{\int_{B_{R_\infty}} u^{(p-1)_+} dx} \right)^{\frac{d}{2\bar{q}-d(p-1)_+}} \frac{\|u\|_{\bar{q}, \bar{R}}}{|B_{\bar{R}}|^{\frac{1}{\bar{q}}}} \\
&\leq I_{\infty, \bar{q}} \left(\frac{\|u\|_{\bar{q}, \bar{R}}}{|B_{\bar{R}}|^{\frac{1}{\bar{q}}}} \frac{1}{\inf_{x \in B_{R_\infty}} u(x)} \right)^{\frac{d(p-1)}{2\bar{q}-d(p-1)}} \frac{\|u\|_{\bar{q}, \bar{R}}}{|B_{\bar{R}}|^{\frac{1}{\bar{q}}}} \\
&\leq I_{\infty, \bar{q}} \left(\frac{I_{\bar{q}, q}}{I_{-\infty, q}} \right)^{\frac{d(p-1)}{2\bar{q}-d(p-1)}+1} \inf_{x \in \bar{B}_{R_\infty}} u(x)
\end{aligned} \tag{5.13}$$

for any $\bar{q} > \frac{d(p-1)_+}{2}$, where $I_{\infty, \bar{q}}$ is given by (3.5) and since $0 < \bar{q} < d/(d-2)$ we require the additional condition (3.7). In the third step we have used the lower bound (5.12). \square

Remark. Notice that the constant \mathcal{H}_p does not depend on u in the range $0 \leq p < p_c$, and it does not depend on $\lambda > 0$ when moreover $p \neq 1$.

6. Local Absolute bounds

In this section we will prove local absolute lower bounds when $0 < p < 1$ and local absolute upper bounds when $1 < p < p_c$ as a consequence of the Harnack inequalities of the previous section together with the Caccioppoli estimates (2.11).

Theorem 6.1 (Local Absolute bounds). *Let $\Omega \subseteq \mathbb{R}^d$ and let $\lambda > 0$. Let u be a local nonnegative weak solution to $-\Delta u = \lambda u^p$ in $B_{R_0} \subseteq \Omega$, with $0 < p < p_c = d/(d-2)$. Then for any $0 < R_\infty < \bar{R} < R_0$ there exists a constant $\mathcal{H}_p > 0$ that does not depend on u , such that*

$$\sup_{x \in B_{R_\infty}(x_0)} u(x) \leq \mathcal{H}_p \left(\frac{8R_0^d}{\lambda(R_0 - R)^2 R^d} \right)^{\frac{1}{p-1}} \quad \text{when } 1 < p < p_c = \frac{d}{d-2}, \tag{6.1}$$

and, if $u \not\equiv 0$ on B_{R_0}

$$\inf_{x \in \bar{B}_{R_\infty}(x_0)} u(x) \geq \mathcal{H}_p^{-1} \left(\frac{\lambda(R_0 - R)^2 R^d}{8R_0^d} \right)^{\frac{1}{1-p}} \quad \text{when } 0 < p < 1. \tag{6.2}$$

The constant \mathcal{H}_p is given by (5.6) when $0 < p < 1$ and by (5.11) when $1 < p < p_c$.

Remark. The way the estimate blows up as $R \rightarrow R_0$ is $(R_0 - R)^{-2/(p-1)}$ which is natural from scaling considerations and is predicted by Dancer in the papers [11, 12].

Proof. We combine the quantitative Harnack inequalities of Theorems 5.2 and 5.3 together with the quantitative Caccioppoli estimates (2.11)

$$\lambda \int_{B_R} u^{p-1} dx \leq \frac{8\omega_d R_0^d}{(R_0 - R)^2}$$

which implies, when $p > 1$,

$$\inf_{x \in \bar{B}_R} u(x) \leq \left(\frac{1}{|B_R|} \int_{B_R} u^{p-1} dx \right)^{\frac{1}{p-1}} \leq \left(\frac{8R_0^d}{\lambda(R_0 - R)^2 R^d} \right)^{\frac{1}{p-1}} \quad (6.3)$$

and when $0 < p < 1$ as

$$\begin{aligned} \left(\frac{\lambda(R_0 - R)^2 R^d}{8R_0^d} \right)^{\frac{1}{1-p}} &\leq \left(\frac{|B_R|}{\int_{B_R} u^{p-1} dx} \right)^{\frac{1}{1-p}} \leq \left(\frac{1}{\sup_{x \in \bar{B}_R} u(x)^{p-1}} \right)^{\frac{1}{1-p}} \\ &= \sup_{x \in \bar{B}_R} u(x) \end{aligned} \quad (6.4)$$

The above inequalities can be now combined with the corresponding Harnack inequalities of Theorems 5.2 and 5.3, which have the form

$$\sup_{x \in \bar{B}_R} u(x) \leq \mathcal{H}_p \inf_{x \in \bar{B}_R} u(x)$$

to obtain the desired bounds in both cases. The constant \mathcal{H}_p is given by (5.6) when $0 < p < 1$ and by (5.11) $1 < p < p_c$. \square

7. Regularity. Local bounds for the gradients

In this section we will prove L^∞ bounds for the gradients, to conclude that solutions to $-\Delta u = \lambda u^p$ are indeed local Lipschitz functions. The strategy to prove such results is to show that the incremental quotients $u_{h,i}$ satisfy the equation $-\Delta u_{h,i} \leq b(x)u_{h,i}$ for a suitable $b(x)$, so that we can apply the local L^∞ bounds of Theorem 3.8. We start with a numerical Lemma.

Lemma 7.1. *The following inequality holds for any $a, b \geq 0$*

$$(a - b)(a^p - b^p) \leq (p \vee 1) \max\{a^{p-1}, b^{p-1}\}(a - b)^2, \quad \text{and for any } p \geq 0. \quad (7.1)$$

Moreover the following inequality holds for any $a, b \geq 0$ and $p \geq 1$:

$$a^p - b^p \geq p b^{p-1}(a - b). \quad (7.2)$$

Proof. If $a \geq b$ the validity of (7.1) is equivalent, setting $x = \frac{b}{a}$, to the validity of $(1-x)(1-x^p) \leq p(1-x)^2$ for all $x \in [0, 1]$, that is to $1-x^p \leq p(1-x)$ for all $x \in [0, 1]$, which does in fact hold if $p \geq 1$ by the concavity of $g(x) := 1-x^p$, since the line $h(x) := p(1-x)$ is the tangent to g at $x = 1$. The case $a < b$ follows as well by interchanging the role of a and b . The case $0 < p < 1$ can be proven analogously: in fact the stated inequality is equivalent to $1-x^p \leq 1-x$ for any $x \in [0, 1]$, which holds true by the convexity of $h(x) = 1-x^p$ for any $p \in (0, 1)$.

The second inequality (7.2) follows by the inequality $x^p - 1 \geq p(x - 1)$ for all $x \geq 0$ which is valid since $x^p - 1$ is convex so that its graph lies above its tangent at $x = 1$. \square

Short reminder about incremental quotients in $W^{1,q}$. Here we follow Giusti [23]. It is well known that if $u \in W^{1,q}(\Omega)$ then its incremental quotients is defined as

$$u_{h,i} := \frac{u(x + he_i) - u(x)}{h}$$

where e_i denotes the unit vector in the direction x_i , cf. [18, 23]. Let us recall some properties of the incremental quotients:

(i) If $u \in W^{1,q}(\Omega)$, then its incremental quotient $u_{h,i}$ is defined in the set

$$\Omega_{|h|} := \{x \in \Omega \mid \text{dist}(x, \partial\Omega) > |h|\}, \quad \text{moreover} \quad u_{h,i} \in W^{1,q}(\Omega_{|h|}).$$

(ii) If $u \in W^{1,q}(\Omega)$ for $1 \leq q \leq \infty$ and $\Sigma \subset\subset \Omega$, then for any $|h| < \text{dist}(\Sigma, \Omega)/(10\sqrt{d})$ we have

$$\|u_{h,i}\|_{L^q(\Sigma)} \leq 5^{\frac{d}{q}} \|\partial_i u\|_{L^q(\Omega)}. \quad (7.3)$$

for a proof of the latter fact we refer to Lemma 8.1 of [23].

(iii) Let $u \in L^q(\Omega)$, $1 < q < \infty$, and assume that there is a constant K such that for every h small enough we have $\|u_{h,i}\|_{L^q(\Omega_{|h|})} \leq K$. Then $\partial_i u \in L^q(\Omega)$ and $\|\partial_i u\|_{L^q(\Omega)} \leq K$. Moreover $u_{h,i} \rightarrow \partial_i u$ in $L^q_{\text{loc}}(\Omega)$ as $h \rightarrow 0$. For a proof of this fact we refer to Lemma 8.2 of [23].

We can now state and prove the following theorem.

Theorem 7.2 (Local upper bounds for the gradient). *Let $\Omega \subseteq \mathbb{R}^d$ and let $\lambda > 0$. Let u be a local nonnegative weak solution to $-\Delta u = \lambda u^p$ in $B_{R_0} \subseteq \Omega$, with $0 < p < p_c = d/(d-2)$. Then for any $0 < R_\infty < R_0$ we have*

$$\|\nabla u\|_{\infty, R_\infty} \leq K[u] \|u\|_{2, R_0} \quad (7.4)$$

where

$$\begin{aligned} K[u] &= \left(\frac{15}{R_0 - R_\infty} \right)^{\frac{d}{2}} \left[\lambda b_{p, R_0}[u] + \frac{18d}{(R_0 - R_\infty)^2} \right]^{\frac{1}{2}} \left(\frac{2d^d}{2^d} \right)^{\frac{d^2}{8}} \\ &\times \left[16(d+2) + \frac{(R_0 - R_\infty)^2}{9R_\infty^2} \right] \\ &+ \left(\frac{d\mathcal{S}_2^2(p \vee 1)}{d-2} \right)^{\frac{d}{2}} \frac{4(R_0 - R_\infty)^2}{9(d-2)} |B_{R_0}|^{\frac{d-2}{d}} (\lambda b_{p, R_0}[u])^{\frac{d}{2}} \end{aligned} \quad (7.5)$$

with

$$b_{p, R_0}[u] \leq \begin{cases} 1, & \text{if } p = 1, \\ \frac{8R_0^d \mathcal{H}_p^{|p-1|}}{\lambda(R_0 - R_\infty)^2 R_\infty^d}, & \text{if } 0 \leq p < p_c \text{ and } p \neq 1, \\ \|u\|_{\infty, R_0}^{p-1}, & \text{if } p_c \leq p < p_s, \end{cases} \quad (7.6)$$

where the constant \mathcal{H}_p is given by (5.6) when $0 < p < 1$ and by (5.11) when $1 < p < p_c$.

Proof. The proof is divided into several steps. We start fixing $h_0 > 0$ small enough.

• **STEP 1.** *The equation satisfied by the incremental quotients.* First we deduce formally the equation for the positive and negative part, then we justify it rigorously at the end of this step, using Kato's inequality. If u is a solution to $-\Delta u = \lambda u^p$, then the equation satisfied by $u_{h,i}^+$ is

$$-\Delta u_{h,i}^+ = b^+(x, h) u_{h,i}^+ \leq \lambda(p \vee 1) b_p u_{h,i}^+, \quad \text{for all } |h| \leq h_0, \quad (7.7)$$

where

$$b_p = b_{p, R_0}[u] := \begin{cases} \sup_{B_{R_0}} u^{p-1} & \text{if } 1 \leq p < p_s \\ \left[\inf_{B_{R_0}} u^{1-p} \right]^{-1} & \text{if } 0 \leq p < 1 \end{cases} \quad (7.8)$$

and we observe that $b_{p,R}[u] \leq b_{p,R_0}[u]$ for any $0 < R < R_0$. Indeed, when $u_{h,i} \geq 0$:

$$\begin{aligned} -\Delta u_{h,i}^+ &= \lambda \frac{u^p(x + he_i) - u^p(x)}{h} = \lambda \frac{u^p(x + he_i) - u^p(x)}{u(x + he_i) - u(x)} \frac{u(x + he_i) - u(x)}{h} \\ &:= b^+(x, h) u_{h,i}^+ \\ &\leq \lambda(p \vee 1) \max \{ u^{p-1}(x + he_i), u^{p-1}(x) \} u_{h,i}^+ \end{aligned}$$

by using the numerical inequality (7.1), namely $(a - c)(a^p - c^p) \leq (p \vee 1) \max \{ a^{p-1}, c^{p-1} \} (a - c)^2$ valid for any $p > 0$ and all $a, c \geq 0$ to estimate

$$b^+(x, h) = \lambda \frac{u^p(x + he_i) - u^p(x)}{u(x + he_i) - u(x)} \leq \lambda(p \vee 1) \max \{ u^{p-1}(x + he_i), u^{p-1}(x) \}$$

we have used the fact that $u^p(x + he_i) - u^p(x)$ and $u(x + he_i) - u(x)$ have the same sign.

When $p \geq 1$ we have

$$-\Delta u_{h,i}^+ = b^+(x, h) u_{h,i}^+ \leq \lambda(p \vee 1) \sup_{B_{R+h_0}} (u^{p-1}) u_{h,i}^+,$$

while when $0 \leq p < 1$ we have

$$-\Delta u_{h,i}^+ = b^+(x, h) u_{h,i}^+ \leq \frac{\lambda(p \vee 1)}{\inf_{B_{R+h_0}} u^{1-p}} u_{h,i}^+$$

On the other hand, if u is a solution to $-\Delta u = \lambda u^p$, then the equation satisfied by $u_{h,i}^-$ is

$$-\Delta u_{h,i}^- = b^-(x, h) u_{h,i}^- \leq \lambda(p \vee 1) b_p u_{h,i}^-, \quad \text{for all } |h| \leq h_0, \quad (7.9)$$

where b_p is given by (7.8). Indeed when $u_{h,i} \leq 0$ we have that

$$\begin{aligned} -\Delta u_{h,i}^- &= -\lambda \frac{u^p(x + he_i) - u^p(x)}{h} = -\lambda \frac{u^p(x + he_i) - u^p(x)}{u(x + he_i) - u(x)} \frac{u(x + he_i) - u(x)}{h} \\ &:= b^-(x, h) u_{h,i}^- \\ &\leq \lambda(p \vee 1) \max \{u^{p-1}(x + he_i), u^{p-1}(x)\} u_{h,i}^- \end{aligned}$$

for the same arguments as above. Now it remains to justify the formal calculations made above. First we recall Kato's inequality: if $j : \mathbb{R} \rightarrow \mathbb{R}$ is a convex function such that $j(0) = 0$, $j'(v) > 0$ if $v > 0$, then $\Delta j(v) \geq j'(v) \Delta v$, in the weak sense, whenever $\Delta v \in L^1_{\text{loc}}(\Omega)$. Consider a sequence of convex function j_ε that approximate $j(u_{h,i}) = u_{h,i}^+$ and such that $j_\varepsilon(0) = 0$, $j'_\varepsilon(u_{h,i}) > 0$ if $u_{h,i} > 0$. Then by Kato's inequality, we have that indeed u^+ satisfy the weak formulation

$$\begin{aligned} \int_K \nabla \varphi \cdot \nabla j_\varepsilon(u_{h,i}) \, dx &= - \int_K \varphi \Delta j_\varepsilon(u_{h,i}) \, dx \leq - \int_K \varphi j'_\varepsilon(u_{h,i}) \Delta u_{h,i} \, dx \\ &= \int_K \varphi j'_\varepsilon(u_{h,i}) b^+(x) u_{h,i} \, dx \\ &\leq \int_K \varphi (j_\varepsilon(u_{h,i}) + \varepsilon) b^+(x) \, dx \end{aligned}$$

for any subdomain with compact closure $K \subset \Omega$, and all bounded $0 \leq \varphi \in C^1_0(K)$. Passing to the limit as $\varepsilon \rightarrow 0$ proves that $u_{h,i}^+$ is a weak subsolution to $-\Delta u_{h,i}^+ \leq b^+(x) u_{h,i}^+$. A similar procedure can be applied to $u_{h,i}^-$, therefore all the formal calculations made above are justified.

• **STEP 2. L^∞ -bounds for the gradients.** Since $|u_{h,i}| = u_{h,i}^+ + u_{h,i}^-$ is a weak nonnegative subsolution to $-\Delta |u_{h,i}| \leq \lambda(p \vee 1) b_p |u_{h,i}| := b(x) |u_{h,i}|$, we can apply the upper bounds of Theorem 3.5 that read

$$\|u_{h,i}\|_{\infty, R} \leq \frac{K_2^{(3)}[b]}{h_0^{\frac{d}{2}}} \|u_{h,i}\|_{2, R+h_0} \quad (7.10)$$

with $q = 2$ and the expression of the constant obtained by letting $r \rightarrow \infty$, since $b(x) \in L^\infty(B_{R+h_0})$:

$$\begin{aligned} K_2^{(3)}[b] &= \left(\frac{2d^d}{2^d}\right)^{\frac{d^2}{8}} \left[16(d+2) + \frac{h_0^2}{R^2} \right. \\ &\quad \left. + \left(\frac{d \mathcal{S}_2^2}{d-2}\right)^{\frac{d}{2}} \frac{4h_0^2}{d-2} |B_{R+h_0}|^{\frac{d-2}{d}} \|b\|_{\infty, R+h_0}^{\frac{d}{2}} \right]^{\frac{d}{4}} \\ &\leq \left(\frac{2d^d}{2^d}\right)^{\frac{d^2}{8}} \left[16(d+2) + \frac{h_0^2}{R^2} \right. \\ &\quad \left. + \left(\frac{d \mathcal{S}_2^2(p \vee 1)}{d-2}\right)^{\frac{d}{2}} \frac{4h_0^2}{d-2} |B_{R+h_0}|^{\frac{d-2}{d}} (\lambda b_p)^{\frac{d}{2}} \right]^{\frac{d}{4}} \end{aligned} \quad (7.11)$$

since

$$\|b\|_{\infty, R+h_0}^{\frac{d}{2}} = (\lambda(p \vee 1) b_p)^{\frac{d}{2}}.$$

Next we observe that by inequality (7.3) it follows that for any $\delta > 0$ and any $|h| < \delta/(10\sqrt{d})$ we have

$$\|u_{h,i}\|_{2, R+h_0} \leq 5^{\frac{d}{2}} \|\partial_i u\|_{2, R+h_0+\delta}.$$

Finally, since

$$\frac{\|u_{h,i}\|_{s,R}}{|B_R|^{\frac{1}{s}}} \leq \|u_{h,i}\|_{\infty, R} \leq K$$

holds for any $|h| \leq h_0$ with K that do not depend on s , then by remark (iii) above we have that

$$\frac{\|\partial_i u\|_{s,R}}{|B_R|^{\frac{1}{s}}} \leq K.$$

Letting now $s \rightarrow \infty$ in the above expression gives $\|\partial_i u\|_{\infty, R} \leq K$. Therefore we have proven that

$$\|\partial_i u\|_{\infty, R} \leq \frac{K_2^{(3)}[b]}{h_0^{\frac{d}{2}}} 5^{\frac{d}{2}} \|\partial_i u\|_{2, R+h_0+\delta}, \quad (7.12)$$

with $K_2^{(3)}[b]$ as in (7.11) which implies

$$\|\nabla u\|_{\infty, R} \leq \frac{K_2^{(3)}[b]}{h_0^{\frac{d}{2}}} 5^{\frac{d}{2}} \|\nabla u\|_{2, R+h_0+\delta}, \quad (7.13)$$

• **STEP 3. Energy inequalities.** We now need the energy inequalities (2.3) to estimate the L^2 norm of the gradient of u in terms of u itself. We choose $\alpha = 1$ there so that the choice $\delta = 0$ is admissible.

$$\begin{aligned} \int_{B_{R+h_0+\delta}} |\nabla u|^2 \varphi \, dx &\leq \int_{\Omega} |\nabla u|^2 \varphi \, dx \leq \lambda \int_{\Omega} u^{p+1} \varphi \, dx + \frac{1}{2} \int_{\Omega} u^2 \Delta \varphi \, dx \\ &\leq \lambda \int_{B_{R+h_0+2\delta}} u^{p+1} \, dx + \frac{2d}{\delta^2} \int_{B_{R+h_0+2\delta}} u^2 \, dx \\ &\leq \left(\lambda b_p + \frac{2d}{\delta^2} \right) \int_{B_{R+h_0+2\delta}} u^2 \, dx \\ &\leq \left(\lambda b_p + \frac{2d}{\delta^2} \right) \|u\|_{2, R+h_0+2\delta}^2 \end{aligned} \quad (7.14)$$

since we have used the fact that $u^{p-1} \leq b_p$ for any $0 \leq p < p_s$ and the test function φ of Lemma 2.2 with the choice of balls $B_{R+h_0+\delta} \subset B_{R+h_0+2\delta}$.

• STEP 4. Putting all the pieces together, we have obtained

$$\begin{aligned} \|\nabla u\|_{\infty, R} &\leq \frac{K_2^{(3)}[b]}{h_0^{\frac{d}{2}}} 5^{\frac{d}{2}} \|\nabla u\|_{2, R+h_0+\delta} \\ &\leq \frac{K_2^{(3)}[b]}{h_0^{\frac{d}{2}}} 5^{\frac{d}{2}} \left(\lambda b_p + \frac{2d}{\delta^2} \right)^{\frac{1}{2}} \|u\|_{2, R+h_0+2\delta}. \end{aligned} \quad (7.15)$$

We finally choose $h_0 = \delta > 0$ and we let $R_\infty = R$, $R_0 = R + h_0 + 2\delta = R + 3\delta$, so that $\delta = (R_0 - R_\infty)/3$ and we have obtained

$$\begin{aligned} \|\nabla u\|_{\infty, R_\infty} &\leq K_2^{(3)}[b] \left(\frac{15}{R_0 - R_\infty} \right)^{\frac{d}{2}} \left(\lambda b_p + \frac{18d}{(R_0 - R_\infty)^2} \right)^{\frac{1}{2}} \|u\|_{2, R_0} \\ &:= K[u] \|u\|_{2, R_0} \end{aligned} \quad (7.16)$$

where we recall that, with the above choices of h_0, δ we have (see (7.11))

$$\begin{aligned} K_2^{(3)}[b] &\leq \left(\frac{2d^d}{2^d} \right)^{\frac{d^2}{8}} \left[16(d+2) + \frac{(R_0 - R_\infty)^2}{9R_\infty^2} \right. \\ &\quad \left. + \left(\frac{d\mathcal{S}_2^2(p \vee 1)}{d-2} \right)^{\frac{d}{2}} \frac{4(R_0 - R_\infty)^2}{9(d-2)} |B_{R_0}|^{\frac{d-2}{d}} (\lambda b_p)^{\frac{d}{2}} \right]^{\frac{d}{4}}. \end{aligned} \quad (7.17)$$

Finally we observe that b_p can be bounded depending on the values of p as follows:

(i) If $0 \leq p < 1$ we can use the absolute bounds (6.2) to get

$$b_p = \frac{1}{\inf_{B_{R_0}} u^{1-p}} \leq \mathcal{H}_p^{1-p} \frac{8R_0^d}{\lambda(R_0 - R_\infty)^2 R_\infty^d}, \quad (7.18)$$

the constant \mathcal{H}_p being given in this case by (5.6).

(ii) If $p = 1$ then $b_p = 1$.

(iii) If $1 < p < p_c$ we can use the absolute bounds (6.1)

$$b_p = \sup_{x \in B_R(x_0)} u^{p-1}(x) \leq \mathcal{H}_p^{p-1} \frac{8R_0^d}{\lambda(R_0 - R_\infty)^2 R_\infty^d}, \quad (7.19)$$

the constant \mathcal{H}_p being given in this case by (5.11).

(iv) If $p_c \leq p < p_s$, we just leave $b_p = \|u\|_{\infty, R_0}^{p-1}$. \square

When $1 < p < p_c$ we have local absolute bounds for the gradients, which seem to be new.

Theorem 7.3 (Local absolute bounds for the gradient when $1 < p < p_c$). *Let $\Omega \subseteq \mathbb{R}^d$ and let $\lambda > 0$. Let u be a local nonnegative weak solution to $-\Delta u = \lambda u^p$ in $B_{R_0} \subseteq \Omega$, with $1 < p < p_c = d/(d-2)$. Then for any $0 < R_\infty < R_0$ we have*

$$\|\nabla u\|_{\infty, R_\infty} \leq K \quad (7.20)$$

where

$$\begin{aligned}
K &= \left(\frac{d^d}{2^{d-1}} \right)^{\frac{d^2}{8}} \frac{(15)^{\frac{d}{2}} \mathcal{H}_p \omega_d^{\frac{1}{2}} R_\infty^{\frac{d}{2}}}{(R_0 - R_\infty)^{1+\frac{d}{2}+\frac{2}{p-1}}} \left[\frac{8R_0^d \mathcal{H}_p^{p-1}}{R_\infty^d} + 18d \right]^{\frac{1}{2}} \left(\frac{8R_0^d}{\lambda R_\infty^d} \right)^{\frac{1}{p-1}} \\
&\times \left[16(d+2) + \frac{(R_0 - R_\infty)^2}{9R_\infty^2} \right. \\
&\left. + \left(\frac{d \mathcal{S}_2^2 p}{d-2} \right)^{\frac{d}{2}} \frac{2^{2+\frac{3}{2}d} \omega_d^{\frac{(d-2)}{d}} R_0^{\frac{d^2}{2}+(d-2)}}{9(d-2)(R_0 - R_\infty)^{2(d-1)} R_\infty^{\frac{d^2}{2}}} \mathcal{H}_p^{\frac{d(p-1)}{2}} \right]^{\frac{d}{4}}
\end{aligned} \tag{7.21}$$

where the constant \mathcal{H}_p is given by (5.11) and depends on R_0, R_∞ as well.

8. Table of results

Let us resume the main results of this paper: recall that $d \geq 3$ and

$$p_c = \frac{d}{d-2}, \quad p_s = \frac{d+2}{d-2}, \quad \bar{q} = \frac{d(p-1)_+}{2}, \quad q_0 = \frac{2^{\frac{d-3}{2}}}{d\omega_d^2[e(d-1)+\varepsilon]}, \quad \forall \varepsilon > 0.$$

| | Upper I | Upper II | Lower | Harnack | Absolute | Gradient |
|-----------------|--|--|---------------------------------|--------------------------------|-------------------|----------------------|
| $0 \leq p < 1$ | $0 < q \rightarrow \infty$ Thm. 3.1 | $q_0 > 0,$ $r > 0$ Thm. 3.9 | $0 < q < q_0$ Thm. 4.6 | \mathcal{H}_p Thm. 5.2 | lower Thm. 6.1 | upper Thm. 7.2 |
| $p = 1$ | $0 < q \rightarrow \infty$ Thm. 3.1 | $q_0 > 0,$ $b \in L^r,$ $r > \frac{d}{2}$ Thm. 3.8 | $0 < q < q_0$ Thm. 4.6 | \mathcal{H}_1 Thm. 5.2 | No | upper Thm. 7.2 |
| $1 < p < p_c$ | $\bar{q} < q \rightarrow \infty$ Thm. 3.1 | $q_0 > 0,$ $b = \lambda u^{p-1} \in L^r$ $r > \bar{q}$ Thm. 3.9 | $\bar{q} < q < p_c$ Thm. 4.8 | \mathcal{H}_p Thm. 5.3 | upper Thm. 6.1 | absolute Thm. 7.3 |
| $p_c < p < p_s$ | $\bar{q} < q \rightarrow \infty$ Thm. 3.1 | $q_0 > 0,$ $b = \lambda u^{p-1} \in L^r$ $r > \bar{q}$ Thm. 3.9 | $0 < q < q_0$ Thm. 4.6 | $\mathcal{H}_p[u]$ Thm. 5.1 | No | upper Thm. 7.2 |

Recall the bounds:

$$\text{Upper I} \quad \|u\|_{L^\infty(B_{R_\infty})} \|u\|_{L^{p-1}(B_{R_\infty})}^{\mu(p-1)_+} \leq I_{\infty,q} \frac{\|u\|_{L^q(B_{R_0})}^{1+\mu(p-1)_+}}{|B_{R_0}|^{\frac{1}{q}}}, \quad \mu = \frac{d}{2q-d(p-1)_+}$$

$$\text{Upper II} \quad \|u\|_{\infty, R_\infty} \leq \frac{A_{q_0}^{(1)}}{(R-R_\infty)^{\frac{d}{q_0}}} \left[A_{q_0}^{(2)} + A_{q_0}^{(3)} \|b\|_{L^r(B_{R_0})}^{\frac{rd}{2r-d}} \right]^{\frac{d}{2q_0}} \|u\|_{q_0, R}$$

$$\text{Lower} \quad \inf_{x \in B_{R_\infty}} u(x) = \|u\|_{L^{-\infty}(B_{R_\infty})} \geq I_{-\infty,q} \frac{\|u\|_{L^q(B_{R_0})}}{|B_{R_0}|^{\frac{1}{q}}}.$$

$$\text{Harnack} \quad \sup_{x \in B_{R_\infty}} u(x) \leq \mathcal{H}_p[u] \inf_{x \in B_{R_\infty}} u(x)$$

where $\mathcal{H}_p[u]$ depends on u only when $p_c \leq p < p_s$ through some local norms as follows

$$\begin{aligned} \mathcal{H}_p[u] &= \mathcal{H}_p[u](d, \bar{q}, \underline{q}, \varepsilon, R_0, R_\infty) \\ &= \frac{I_{\infty, \bar{q}}}{I_{-\infty, \underline{q}}} \left(\frac{\left(\int_{B_{R_0}} u^{\underline{q}} dx \right)^{\frac{(p-1)_+}{q}}}{\int_{B_{R_\infty}} u^{(p-1)_+} dx} \right)^{\frac{d}{2\bar{q}-d(p-1)_+}} \frac{\left(\int_{B_{R_0}} u^{\bar{q}} dx \right)^{\frac{1}{\bar{q}}}}{\left(\int_{B_{R_0}} u^{\underline{q}} dx \right)^{\frac{1}{\underline{q}}}}. \end{aligned}$$

whereas $\mathcal{H}_p[u]$ can be taken to be independent of u if $p \in [0, p_c)$, see (5.6), (5.11).

Gradient $\quad \|\nabla u\|_{\infty, R_\infty} \leq K[u] \|u\|_{2, R_0}.$

References

1. A. Ambrosetti, G. Prodi, “A primer of nonlinear analysis”, Cambridge Studies in Advanced Mathematics, **34**, Cambridge University Press, Cambridge, 1993. viii+171 pp. ISBN: 0-521-37390-5
2. M. Bonforte, G. Grillo, J. L. Vázquez, *Behaviour near extinction for the Fast Diffusion Equation on bounded domains*, J. Math. Pures Appl. **97** (2012), 1-38.
3. M. Bonforte, G. Grillo, J. L. Vázquez, *Quantitative bounds for the Dirichlet problem for semilinear elliptic equations on bounded domains*, in preparation (2012).
4. M. Bonforte, J. L. Vázquez, *Positivity, local smoothing, and Harnack inequalities for very fast diffusion equations*, Advances in Math. **223** (2010), 529–578.
5. H. Brezis, T. Kato, *Remarks on the Schrödinger operator with singular complex potentials*, J. Math. Pures Appl. (9) **58** (1979), no. 2, 137-151.
6. H. Brezis, L. Nirenberg, *Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents*, Comm. Pure Appl. Math. **36** (1983), no. 4, 437-477.
7. H. Brezis, R. E. L. Turner, *On a class of superlinear elliptic problems*, Comm. PDE **2** (1977), 601-614.
8. C. Budd, J. Norbury, *Semilinear Elliptic Equations and Supercritical Growth*, J. Diff. Eq. **68** (1987), 169–197.
9. X. Cabré, *Regularity of minimizers of semilinear elliptic problems up to dimension 4*, Comm. Pure Appl. Math. **63** (2010), no. 10, 1362-1380.
10. L. A. Caffarelli, B. Gidas, J. Spruck, *Asymptotic symmetry and local behavior of semilinear elliptic equations with critical Sobolev growth*, Comm. Pure Appl. Math. **42** (1989), no. 3, 271-297.
11. E. N. Dancer, *On the influence of domain shape on the existence of large solutions of some superlinear problems*, Math. Ann. **285** (1989), no. 4, 647-669.
12. E. N. Dancer, *Superlinear problems on domains with holes of asymptotic shape and exterior problems*, Math. Z., **229** (1998) 475–491.

13. J. Dávila, L. Dupaigne, A. Farina, *Partial regularity of finite Morse index solutions to the Lane-Emden equation*, J. Funct. Anal. **261** (2011) 218–232.
14. D. G. de Figueiredo, P.-L. Lions, R. D. Nussbaum, *A priori estimates and existence of positive solutions of semilinear elliptic equations*, J. Math. Pures Appl. **61** (1982), 41–63.
15. M. del Pino, M. Musso, F. Pacard, *Boundary singularities for weak solutions of semilinear elliptic problems*, J. Funct. Anal. **253** (2007) 241–272.
16. E. DiBenedetto, U. Gianazza, V. Vespri, *Harnack type estimates and Hölder continuity for non-negative solutions to certain sub-critically singular parabolic partial differential equations*, Manuscripta Math. **131** (2010), no. 1-2, 231–245
17. E. DiBenedetto, U. Gianazza, V. Vespri, “Harnack’s inequality for degenerate and singular parabolic equations”, Springer Monographs in Mathematics, Springer 2011.
18. L. C. Evans, “Partial differential equations. Second edition”, Graduate Studies in Mathematics, 19. American Mathematical Society, Providence, RI, 2010. xxii+749 pp. ISBN: 978-0-8218-4974-3
19. A. Farina, *On the classification of solutions of the Lane-Emden equation on unbounded domains of \mathbb{R}^N* , J. Math. Pures Appl. **87** (5) (2007) 537–561.
20. B. Gidas, Wei Ming Ni, L. Nirenberg, *Symmetry and related properties via the maximum principle*, Comm. Math. Phys. **68** (1979), 209–243.
21. B. Gidas, J. Spruck, *Global and local behavior of positive solutions of nonlinear elliptic equations*, Comm. Pure Appl. Math. **34**, 1981, 525–598.
22. D. Gilbarg, N. S. Trudinger, “Elliptic Partial Differential Equations of Second Order”, GMW **224**, 1977, Springer Verlag, Heidelberg.
23. E. Giusti, “Direct methods in the calculus of variations”, World Scientific Publishing Co., Inc., River Edge, NJ, 2003. viii+403 pp. ISBN: 981-238-043-4.
24. O. Ladyzhenskaya, N. N. Ural’tseva, “Linear and quasilinear elliptic equations”. Translated from the Russian by Scripta Technica, Inc. Translation editor: Leon Ehrenpreis Academic Press, New York-London 1968 xviii+495 pp.
25. R. Mazzeo, F. Pacard, *A construction of singular solutions for a semilinear elliptic equation using asymptotic analysis*, J. Differential Geom. **44** (1996), no. 2, 331–370.
26. W. M. Ni, R. D. Nussbaum, *Uniqueness and Nonuniqueness for Positive Radial Solutions of $\Delta u + f(u, r) = 0$* , Comm. Pure Appl. Math. **38** (1985), 67–108.
27. L. Nirenberg, *On elliptic partial differential equations*, Ann. Scuola Norm. Sup. Pisa **13** (1959), 115–162.
28. F. Pacard, *Existence de solutions faibles positives de $-\Delta u = u^\alpha$ dans des ouverts bornés de \mathbb{R}^n , $n \geq 3$* , C. R. Acad. Sci. Paris Sr. I Math. **315** (1992), no. 7, 793–798.
29. F. Pacard, *A note on the regularity of weak solutions of $-\Delta u = u^\alpha$ in \mathbb{R}^n , $n \geq 3$* , Houston J. Math. **18** (1992), no. 4, 621–632.
30. F. Pacard, *Existence and convergence of positive weak solutions of $-\Delta u = u^{n/(n-2)}$ in bounded domains of \mathbb{R}^n , $n \geq 3$* , Calc. Var. Partial Differential Equations **1** (1993), no. 3, 243–265.
31. F. Pacard, *Partial regularity for weak solutions of a nonlinear elliptic equation*, Manuscripta Math. **79** (1993), no. 2, 161–172.

32. P. Poláčik, P. Quittner, P. Souplet, *Singularity and decay estimates in super-linear problems via Liouville-type theorems. I. Elliptic equations and systems*, Duke Math. J. **139** (2007), no. 3, 555–579.
33. P. Pucci, J. Serrin, “The maximum principle”, Progress in Nonlinear Differential Equations and their Applications, **73**, Birkhäuser, Basel (2007), ISBN: 978-3-7643-8144-8.
34. J. Serrin, *Local behavior of solutions of quasi-linear equations*, Acta Math. **111**, (1964), 247-302.
35. J. Serrin, H. Zou, *Cauchy-Liouville and universal boundedness theorems for quasilinear elliptic equations and inequalities*, Acta Math. **189** (2002), 79-142.
36. N. S. Trudinger, *Harnack inequalities for nonuniformly elliptic divergence structure equations*, Invent. Math. **64**, (1981), no. 3, 517-531.
37. N. S. Trudinger, *On Harnack type inequalities and their application to quasilinear elliptic equations*, Comm. Pure Appl. Math. **20** (1967) 721-747.

Matteo Bonforte
Departamento de Matemáticas
Universidad Autónoma de Madrid
28049 Madrid, Spain.
e-mail: matteo.bonforte@uam.es

Gabriele Grillo
Dipartimento di Matematica
Politecnico di Milano
Piazza Leonardo da Vinci 32
20133 Milano, Italy.
e-mail: gabriele.grillo@polimi.it

Juan Luis Vazquez
Departamento de Matemáticas
Universidad Autónoma de Madrid
28049 Madrid, Spain.
e-mail: juanluis.vazquez@uam.es