

DIPARTIMENTO DI MATEMATICA
POLITECNICO DI MILANO

**Spiraling asymptotic profiles of
competition-diffusion systems**

Terracini, S.; Verzini, G.; Zilio, A.

Collezione dei *Quaderni di Dipartimento*, numero **QDD 227**
Inserito negli *Archivi Digitali di Dipartimento* in data 06-07-2017



Piazza Leonardo da Vinci, 32 - 20133 Milano (Italy)

Spiraling asymptotic profiles of competition-diffusion systems

Susanna Terracini, Gianmaria Verzini and Alessandro Zilio

July 4, 2017

Abstract

This paper describes the structure of the nodal set of segregation profiles arising in the singular limit of planar, stationary, reaction-diffusion systems with strongly competitive interactions of Lotka-Volterra type, when the matrix of the inter-specific competition coefficients is asymmetric and the competition parameter tends to infinity. Unlike the symmetric case, when it is known that the nodal set consists in a locally finite collection of curves meeting with equal angles at a locally finite number of singular points, the asymmetric case shows the emergence of spiraling nodal curves, still meeting at locally isolated points with finite vanishing order.

AMS-Subject Classification. 35B25 (35K51,92D25)

Keywords. Lotka-Volterra systems, Singular perturbation, Free boundary problems, Spiral Waves

1 Introduction

This paper describes the structure of the nodal set of segregation profiles arising in the singular limit of planar, stationary, reaction-diffusion systems with strongly competitive interactions of Lotka-Volterra type, as and the competition parameter tends to infinity. This structure has been widely studied when the matrix of the inter-specific competition coefficients is symmetric in connection with either the free boundary of optimal partitions involving shape energies, or the singularities of harmonic maps with values in a stratified varyfold [6, 1, 3, 15, 23]. For such problems, in the planar case, it is known that the nodal set consists in a locally finite collection of curves meeting with equal angles at a locally finite number of singular points. Our aim is to show that the effect of asymmetry of the inter-specific competition rates results in a dramatic change of the nodal pattern, now consisting of spiraling nodal curves, still meeting at locally isolated points with finite vanishing order. It has to be noticed that the asymmetry makes all the usual free boundary toolbox (Almgren and Alt-Caffarelli-Friedman monotonicity formulæ, dimension estimates) unavailable and ad-hoc arguments have been designed. Finally, we point out that spiraling waves also occur in entirely different contexts of reaction-diffusion systems (cfr e.g. [18, 21, 20]).

Lotka-Volterra type systems are the most popular mathematical models for the dynamics of many populations subject to spatial diffusion, internal reaction and either cooperative or competitive interaction. Indeed, such models are associated with reaction-diffusion systems where the reaction is the sum of an intra-specific term, often expressed by logistic type functions, and an inter-specific interaction one, usually quadratic. The study of this reaction-diffusion system has a long history and there exists a large literature on the subject. However, most of these works are concerned with the case of two species. As far as we know, the study in the case of many competing species has been much more limited, starting from two pioneering papers by Dancer and Du [10, 11] in the 1990s, where the competition of three species were considered.

We deal with a regime of strongly competing populations in the stationary case. Hence the equations of the systems take the form

$$-\Delta u_i = f_i(u_i) - \beta u_i \sum_{j \neq i} a_{ij} u_j \quad \text{in } \Omega \subset \mathbb{R}^N, \quad i = 1, \dots, k, \quad (1)$$

where $(a_{ij})_{ij}$ is the matrix of the interspecific competition coefficients, with nonnegative entries, and $\beta \geq 0$. For concreteness we require the reaction terms f_i to be locally Lipschitz, with $f_i(0) = 0$, even though specific results hold under less restrictive assumptions. One may consider also different diffusion coefficients $d_i > 0$ on the left hand side of (1). Nonetheless, in the stationary case, it is not restrictive to assume $d_i = 1$ by a change of unknowns.

We will distinguish between the *symmetric* (i.e. when $a_{ij} = a_{ji}$, for every i, j) and the *asymmetric* case (i.e. when $a_{ij} \neq a_{ji}$ for some i, j), focusing on the latter, when the parameter β diverges to $+\infty$ and we consider nonnegative solutions, that is, $u_i \geq 0$ for all $i = 1, \dots, k$. In this case, it is known that the components satisfy uniform bounds in Hölder norms and converge, up to subsequences, to some limit profiles, having disjoint supports: *the segregated states*.

Theorem 1.1 ([6]). *Let $(u_{1,\beta}, \dots, u_{k,\beta})$, for $\beta > 0$, be a family of solutions to system (1) satisfying a (uniform in β) $L_{\text{loc}}^\infty(\Omega)$ bound as $\beta \rightarrow +\infty$. Then, up to subsequences, there exists $(\bar{u}_1, \dots, \bar{u}_k)$ such that,*

$$u_{i,\beta} \rightarrow \bar{u}_i \quad \text{in } H_{\text{loc}}^1(\Omega) \cap C_{\text{loc}}^{0,\alpha}(\Omega),$$

for every $i = 1, \dots, k$ and $0 < \alpha < 1$. Moreover, the k -tuple $(\bar{u}_1, \dots, \bar{u}_k)$ is a segregated state:

$$\bar{u}_i \bar{u}_j \equiv 0 \quad \text{in } \Omega.$$

In the last decade, both the asymptotics and the qualitative properties of the limit segregated profiles have been the object of an intensive study, mostly in the symmetric case, by different teams [7, 8, 9, 1, 13, 24, 22]. Similarly, the dynamics of strongly competing species has been addressed as a singularly perturbed parabolic reaction-diffusion system in connection with spatially segregated limit profiles in [12, 14, 16, 17]. In the quoted papers, a special attention was paid to the structure of the common zero set of these limit profiles. The following theorem collects the main known facts about the geometry of the nodal set in the stationary symmetric case:

Theorem 1.2 ([6, 1, 15, 23]). *Assume that*

$$a_{ij} = a_{ji} > 0, \quad \text{for every } i \neq j.$$

Let $\bar{U} = (\bar{u}_1, \dots, \bar{u}_k)$ be a segregated limit profile as in Theorem 1.1, and let $\mathcal{Z} = \{x \in \Omega : \bar{U}(x) = 0\}$ its nodal set. Then, there exist complementary subsets \mathcal{R} and \mathcal{W} of \mathcal{Z} , respectively the regular part, relatively open in \mathcal{Z} and the singular part, relatively closed, such that:

- \mathcal{R} is a collection of hyper-surfaces of class $C^{1,\alpha}$ (for every $0 < \alpha < 1$), and for every $x_0 \in \mathcal{R}$

$$\lim_{x \rightarrow x_0^+} |\nabla \bar{U}(x)| = \lim_{x \rightarrow x_0^-} |\nabla \bar{U}(x)| \neq 0,$$

where the limits as $x \rightarrow x_0^\pm$ are taken from the opposite sides of the hyper-surface;

- $\mathcal{H}_{\text{dim}}(\mathcal{W}) \leq N - 2$, and if $x_0 \in \mathcal{W}$ then $\lim_{x \rightarrow x_0} |\nabla \bar{U}(x)| = 0$.

Furthermore, if $N = 2$, then \mathcal{Z} consists in a locally finite collection of curves meeting with definite semi-tangents with equal angles at a locally finite number of singular points.

On the contrary, the present paper is concerned with *asymmetric inter-specific competition rates*, with the purpose to highlighting the substantial differences with the symmetric case in two space dimensions. To describe our main result, we consider a simplified, yet prototypical, boundary value problem with competition terms of Lotka-Volterra type:

$$\begin{cases} -\Delta u_i = -\beta u_i \sum_{j \neq i} a_{ij} u_j & \text{in } \Omega \\ u_i = \varphi_i & \text{on } \partial\Omega, \end{cases} \quad i = 1, \dots, k. \quad (2)$$

Throughout the whole paper we will assume that:

(A1) $\Omega \subset \mathbb{R}^2$ is a simply connected, bounded domain of class $C^{1,\alpha}$;

(A2) $a_{ij} > 0$ for every $j \neq i$;

(A3) $\varphi_i \in C^{0,1}(\partial\Omega)$, $\varphi_i \geq 0$, $\varphi_i \cdot \varphi_j \equiv 0$ for every $1 \leq i \neq j \leq k$;

(A4) the trace function $\varphi = \sum_{i=1}^k \varphi_i$ has only non-degenerate zero, that is,

$$\forall x_0 \in \partial\Omega, \quad \varphi(x_0) = 0 \implies \liminf_{x \rightarrow x_0} \frac{\varphi(x)}{|x - x_0|} \geq C > 0.$$

We are interested in component-wise non-negative solutions. As we already noticed, as $\beta \rightarrow +\infty$ such solutions to (2) exhibit segregation. More precisely, let us introduce the object of our investigation, a class of *segregated states*, satisfying additional conditions, expressed as a system of differential inequalities as follows:

$$\mathcal{S} = \left\{ U = (u_1, \dots, u_k) \in (H^1(\Omega))^k : \begin{array}{l} u_i \geq 0, u_i = \varphi_i \text{ on } \partial\Omega \\ u_i \cdot u_j = 0 \text{ if } i \neq j \\ -\Delta u_i \leq 0, -\Delta \widehat{u}_i \geq 0 \end{array} \right\}, \quad (3)$$

where the *i*-th *hat operator* is defined on the generic *i*-th component of a *k*-tuple as

$$\widehat{u}_i = u_i - \sum_{j \neq i} \frac{a_{ij}}{a_{ji}} u_j, \quad (4)$$

and the differential inequalities are understood in variational sense. This is a free boundary problem, where the interfaces $\partial\{u_i > 0\} \cap \partial\{u_j > 0\}$, separate the supports of u_i and u_j . The following result has been proved in [6].

Theorem 1.3 ([6]). *For every $\beta > 0$ there exists (at least) one solution $(u_{1,\beta}, \dots, u_{k,\beta}) \in (H^1(\Omega))^k$ to system (2). For every sequence of solutions, there exists (at least) one $(\bar{u}_1, \dots, \bar{u}_k) \in \mathcal{S}$ and, up to a subsequence,*

$$u_{i,\beta_n} \rightarrow \bar{u}_i \quad \text{in } H^1(\Omega) \cap C^{0,\alpha}(\overline{\Omega}),$$

for every $i = 1, \dots, k$ and $0 < \alpha < 1$.

In order to set up our result about the nodal set in the case of asymmetric interspecific competition rates, we need some more notation. For any $U \in \mathcal{S}$ we define the *multiplicity* of a point $x \in \overline{\Omega}$, with respect to U , as

$$m(x) = \#\{i : |\omega_i \cap B_r(x)| > 0 \text{ for every } r > 0\}.$$

Our main purpose is to analyze the structure of the free boundary, i.e. the zero set of a *k*-tuple $U \in \mathcal{S}$:

$$\mathcal{Z} = \{x \in \Omega : u_i(x) = 0 \text{ for every } i = 1, \dots, k\}.$$

Such set naturally splits into the union of the *regular part* $\mathcal{R} = \mathcal{Z}_2 := \{\mathbf{x} \in \mathcal{Z} : m(\mathbf{x}) = 2\}$, and of the *singular part*

$$\mathcal{W} = \mathcal{Z} \setminus \mathcal{Z}_2.$$

We collect in the following Lemma some elementary properties about the elements of \mathcal{S} , which have already been obtained in [8].

Lemma 1.4 ([8]). *Let $U \in \mathcal{S}$. Then:*

1. $U \in C^{0,1}(\overline{\Omega})$;
2. if $m(\mathbf{x}_0) = 1$, then there exist i and $r > 0$ such that $\Delta u_i = 0$ in $B_r(\mathbf{x}_0) \cap \Omega$ (in particular, $\mathbf{x}_0 \notin \mathcal{Z}$);
3. if $m(\mathbf{x}_0) = 2$, then there exist i, j and $r > 0$ such that $\Delta(a_{ji}u_i - a_{ij}u_j) = 0$ in $B_r(\mathbf{x}_0) \cap \Omega$;
4. if $\mathbf{x}_0 \in \mathcal{W}$ then $\lim_{r \rightarrow 0} \sup_{B_r(\mathbf{x}_0)} |\nabla u_i| = 0$, for every i .

Notice that properties 2 and 3 in the above lemma are straight consequences of the definitions of \mathcal{S} and m . Now we are ready to state our main result concerning the properties of the segregation boundary.

Theorem 1.5. *Let (A1–4) hold, $U \in \mathcal{S}$, and $\mathcal{Z} = \mathcal{Z}_2 \cup \mathcal{W}$. Then:*

1. \mathcal{Z}_2 is relatively open in \mathcal{Z} , and it consists in the finite union of analytic curves;
2. \mathcal{W} is the union of a finite number of isolated points inside Ω ;
3. for every $\mathbf{x}_0 \in \partial\Omega$, either $m(\mathbf{x}_0) = 1$ or $m(\mathbf{x}_0) = 2$.

Furthermore, if $\mathbf{x}_0 \in \mathcal{W}$ then $m(\mathbf{x}_0) = h \geq 3$, and there exist an explicit constant $\alpha \in \mathbb{R}$ and an explicit bounded function $A = A(\mathbf{x})$ such that

$$\mathcal{U}(r, \vartheta) = A r^\nu \cos\left(\frac{h}{2}\vartheta - \alpha \log r\right) + o(r^\nu) \quad \text{as } r \rightarrow 0, \quad (5)$$

where (r, ϑ) denotes a (suitably rotated) system of polar coordinates about \mathbf{x}_0 , \mathcal{U} is a suitable weighted sum of the components u_i meeting at \mathbf{x}_0 ,

$$\nu = \frac{h}{2} + \frac{2\alpha^2}{h}, \quad \text{and} \quad 0 < A_0 \leq A(r, \vartheta) \leq A_1. \quad (6)$$

In particular, whenever $\alpha \neq 0$, the regular part of the free boundary is described asymptotically by h equi-distributed logarithmic spirals (locally around \mathbf{x}_0).

Remark 1.6. The value of α in (5), (6) is explicit in terms of the coefficients a_{ij} , with i and j belonging to the set of indexes associated to the $h \leq k$ densities which do not identically vanish near \mathbf{x}_0 (see equations (13), (22) below). For instance, when u_1, u_2 and u_3 meet at \mathbf{x}_0 , with $m(\mathbf{x}_0) = 3$, then (up to a change of sign)

$$\alpha = \frac{1}{2\pi} \log\left(\frac{a_{12}}{a_{21}} \cdot \frac{a_{23}}{a_{32}} \cdot \frac{a_{31}}{a_{13}}\right).$$

Consequently, when $\alpha \neq 0$, the vanishing order ν does not depend only on the number of densities involved (as in the symmetric case), but also on the competition coefficients; moreover, the vanishing order is not forced to be half-integer, in great contrast with the symmetric case. Finally, the function \mathcal{U} agrees with each density near \mathbf{x}_0 , up to a constant multiplicative factor, see equation (12).




Figure 1: numerical simulations of functions belonging to the class \mathcal{S} for different values of α . In this particular case, we have considered a system of 3 components (labeled in counterclockwise order as u_1, u_2 and u_3) in the unit ball, with boundary conditions given by suitable restrictions of $|\cos(3/2 \vartheta)|$. In picture (a), $a_{ij} = 1$ for all i, j , which yields $\alpha = 0$ (see Remark 1.6). In picture (b), $a_{ij} = 4$ if $j - i = 1 \pmod 3$ and $a_{ij} = 1$ otherwise, which yields $\alpha = 3 \log 4/2\pi (> 0$, which implies by equation (5) that the free-boundary is described asymptotically by rotations of the clockwise logarithmic spiral $\vartheta = \log 4/\pi \log r$). In picture (c), $a_{ij} = 10$ if $j - i = 1 \pmod 3$ and $a_{ij} = 1$ otherwise, which yields $\alpha = 3 \log 10/2\pi$.

Remark 1.7. In case $a_{ij} = a_{ji}$ for every $j \neq i$, then $\alpha = 0$, and the spirals reduce to straight lines; in this way we recover the equal-angles-property for multiple points already obtained in [8]. On the other hand it is easy to choose the competition coefficients to force $\alpha \neq 0$. For instance, in the case of $k = 3$ densities on the ball, one can use the construction suggested by our proof, in order to obtain the existence of an element of \mathcal{S} , with a prescribed α , for a set of traces having codimension 2 (see Fig. 1). In the same spirit, for any given real number $\nu \geq 3/2$, one can choose the competing coefficients to obtain elements of \mathcal{S} having a multiple point with vanishing order ν .

Remark 1.8. In the asymmetric case, the nodal partition determined by the supports of the components can not be optimal with respect to any Lagrangian energy. Indeed, it is known (cfr. [4, 2, 15, 23]) that boundaries of optimal partitions share the same properties of Theorem 1.2. Hence they can not exhibit logarithmic spirals. This fact is in striking contrast with the picture for symmetric inter-specific competition rates: indeed, in such a case, relatively to Theorem 1.3, we know that solutions to (2) are unique, together with their limit profiles in the class \mathcal{S} (see [9, 24, 13]). Hence, though system (2) does not possess a variational nature, it fulfills a minimization principle in the segregation limit, while this is impossible in the asymmetric setting.

Remark 1.9. Nevertheless, even in the asymmetric case, functions in the class \mathcal{S} still share with the solutions of variational problems, including harmonic functions, the following fundamental features:

- singular points are isolated and have a finite vanishing order;
- the possible vanishing orders are quantized;
- the regular part is smooth.

It is natural to wonder whether similar analogies still hold dimensions higher than two. As already remarked, however, new strategies and unconventional techniques have to be designed to treat the asymmetric case, since a number of standard tools in free boundary problems have to fail in such situation: for instance, as the planar case shows, the Almgren monotonicity formula can not hold, and the singularities do not admit, in general, homogeneous blow-ups.

The proof of Theorem 1.5 proceeds as follows. In Section 2, exploiting some topological properties of the zero set of harmonic functions, we show that it suffices to consider the case in which $\Omega = B$,

and a unique connected component of $\mathcal{W} \cap B$ is joined to the boundary by a finite number of smooth curves that describe $\mathcal{Z}_2 \cap B$. In Section 3, assuming that such connected component is given by a point, we give a description of the set $\mathcal{Z}_2 \cap B$. To do this, by a conformal mapping, we translate the original problem in the ball B to that of describing the zero set of an harmonic function defined on the half plane. Finally, in Section 4 we prove that any connected component of \mathcal{W} is actually just a point. We achieve this by noticing that in any ball contained in B and whose boundary intersects \mathcal{W} , the set \mathcal{Z} coincides necessarily with that of a harmonic function.

Notation

Unless otherwise specified, we adopt the following conventions:

- points in Ω are denoted with \mathbf{x}, \mathbf{y} , and so on; points in \mathbb{R}_+^2 have coordinates (x, y) ;
- the null set of a k -tuple is $\mathcal{Z} = \{\mathbf{x} \in \Omega : u_i(\mathbf{x}) = 0 \forall i = 1, \dots, k\}$;
- the multiplicity of a point is $m(\mathbf{x}) = \#\{i : |\omega_i \cap B_r(\mathbf{x})| > 0 \text{ for every } r > 0\}$;
- $\mathcal{Z}_h = \{\mathbf{x} \in \mathcal{Z} : m(\mathbf{x}) = h\}$, for $h = 0, 1, \dots, k$;
- the singular set is $\mathcal{W} = \mathcal{Z} \setminus \mathcal{Z}_2 = \mathcal{Z}_0 \cup \mathcal{Z}_3 \cup \dots \cup \mathcal{Z}_k$ (we will see that no zero of multiplicity 1 is allowed inside Ω);
- $\omega_i = \{\mathbf{x} \in \Omega : u_i(\mathbf{x}) > 0\}$ (open), $\text{supp}(u_i) = \bar{\omega}_i$ (closed);
- $\Gamma_{ij} = \bar{\omega}_i \cap \bar{\omega}_j \cap \mathcal{Z}_2$;
- an open curve, or simply a curve when no confusion may arise, is a 1-dimensional manifold (without boundary), i.e. the image of a (open) interval through a regular map; in particular, it is locally diffeomorphic to an interval, but it may not be rectifiable. In particular, we will show that Γ_{ij} is such a curve, as far as it is non-empty.

2 Preliminary reduction

In this section we show that, without loss of generality, we can reduce to the model case scenario described in the following assumption (MCS).

Assumption (MCS). Without loss of generality, beyond (A1–4), we can assume that:

- $k \geq 3$;
- for each $i = 1, \dots, k$ both ω_i and $\bar{\omega}_i \cap \partial\Omega$ are connected, simply connected sets;
- the traces φ_i are labelled in counterclockwise sense;
- Γ_{ij} is a non-empty connected open curve whenever $i - j = \pm 1 \pmod k$, and it is empty otherwise;
- \mathcal{W} has a unique connected component, which lies away from $\partial\Omega$
- Ω is the unit ball $B = B_1(\mathbf{0})$, and, for some $|e| = 1$,

$$\mathbf{0} \in \mathcal{W} \subset \{\mathbf{x} \in B : \mathbf{x} \cdot \mathbf{e} \geq 0\}. \quad (7)$$

More precisely, we will show the following result.

Proposition 2.1. Ω can be decomposed in the finite union of some domains, each of which satisfy (MCS), up to a relabelling of the restricted densities and to some conformal deformations.

The rest of this section is devoted to the proof of the above proposition. We refer the reader also to [5, Section 7], where a similar preliminary analysis was conducted under the assumption that $a_{ij} = 1$ for every i and j .

Remark 2.2. In order to reduce to assumption (MCS), we will perform a number of operations like dividing Ω into subsets, adding and/or relabeling densities, and so on. With some abuse of notation, we will always write Ω for the domain and k for the number of densities. Notice that, once the proposition is proved, the proof of Theorem 1.5 will be reduced to show that, under (MCS), \mathcal{W} consists in a single point, around which the asymptotic expansion (5) holds true.

As a first step we use the maximum principle to reduce to connected, simply connected sub-domains.

Lemma 2.3. Each u_i is positive and harmonic in ω_i , and there are no interior 1-multiplicity zeroes: $\mathcal{Z}_1 \cap \Omega = \emptyset$. Furthermore, possibly by introducing a new family of densities, we have that each ω_i is connected and simply connected, and

$$\omega_i \cup \{x : \varphi_i(x) > 0\} \text{ is pathwise connected.}$$

Proof. The first part of the statement follows by Lemma 1.4, and from the strong maximum principle. Let us assume that there exists $x_0 \in \mathcal{Z}_1 \cap \Omega$, that is, let us assume that there exists $x_0 \in \Omega$, $r > 0$ and i such that $u_i(x_0) = 0$ and $|\omega_i \cap B_r(x_0)| > 0$, while $|\omega_j \cap B_r(x_0)| = 0$ for $j \neq i$. But then u_i is harmonic in $B_r(x_0)$ and, since its boundary data on ∂B_r is non-trivial, by the maximum principle we have that $u_i(x_0) > 0$, a contradiction. Finally, since $\partial\omega_i \subset \mathcal{Z} \cup \partial\Omega$, again the maximum principle implies that each connected component σ of some ω_i must satisfy $\partial\sigma \cap \partial\Omega \neq \emptyset$. By assumption (A4) we know that $\partial\Omega \setminus \{\varphi = 0\}$ has a finite number of connected components; we deduce that the same holds also for each ω_i . Introducing, if necessary, further formal densities, we can then assume that each set ω_i is connected. Since (by continuity of the densities u_i) these sets are open, they are also path-connected. Next, it is easily shown that each of these components is simply connected. Indeed, let $\gamma \subset \omega_i$ be any Jordan curve, and let Σ denote the bounded region of \mathbb{R}^2 such that $\partial\Sigma = \gamma$. By the maximum principle, all the other components u_j , $j \neq i$, when restricted to Σ , are trivial. But then the function u_i is harmonic in the mentioned set, and again the maximum principle forces $u_i > 0$ in the interior of this set, implying that the curve γ is contractible in ω_i . Finally, the pathwise connectedness of $\omega_i \cup \{x : \varphi_i(x) > 0\}$ easily follows from the fact that $\partial\Omega$ is of class $C^{1,1}$. \square

Remark 2.4. In principle it may happen that some point $x_0 \in \partial\Omega$ is a zero of multiplicity 1. Nonetheless, in such case, there exists an index i for which φ_i is positive on both sides of x_0 , and such point is separated from \mathcal{Z} .

Lemma 2.5. The set $\mathcal{Z} \cup \partial\Omega$ is connected.

Proof. Let $U, V \subset \mathbb{R}^2$ be two open sets such that $U \cap V = \emptyset$, $(\mathcal{Z} \cup \partial\Omega) \subset U \cup V$. Since $\partial\Omega$ is connected, we deduce that one of the sets, say U , is a subset of Ω and the other, V , contains its boundary: $U \subset \Omega$ and $\partial\Omega \subset V$. Moreover $\partial U \cap (\mathcal{Z} \cup \partial\Omega) = \emptyset$ by definition. We deduce that each connected component of ∂U must be a subset of some ω_i , and by simple connectedness we find that necessarily $U \cap (\mathcal{Z} \cup \partial\Omega) = \emptyset$. \square

Next we turn to analyze the regular part of the segregation boundary.

Lemma 2.6. Each Γ_{ij} is either empty or a C^1 connected, open curve. In the latter case, $\omega_i \cup \Gamma_{ij} \cup \omega_j$ is an open and simply connected subset of Ω . In particular, $\mathcal{Z}_2 = \cup_{i \neq j} \Gamma_{ij}$ is the disjoint union of a finite number of regular curves.

Proof. We first show that if $\mathbf{x}_0 \in \mathcal{Z}_2$, then $\mathbf{x}_0 \in \Gamma_{ij}$, which is locally defined by a smooth curve near \mathbf{x}_0 . Indeed, let $\mathbf{x}_0 \in \mathcal{Z}_2$ and, by definition, let $i \neq j$ and $r > 0$ be such that $\omega_h \cap B_r(\mathbf{x}_0)$ is not empty if and only if $h = i, j$. It follows that the function $a_{ji}u_i - a_{ij}u_j$, restricted to $B_r(\mathbf{x}_0)$, is harmonic and vanishes in \mathbf{x}_0 : since ω_i and ω_j are path-connected, \mathbf{x}_0 is a simple zero, and the implicit function theorem implies that the set Γ_{ij} is represented by a smooth curve, locally near $\mathbf{x}_0 \in \Gamma_{ij}$.

We now show the connectedness of Γ_{ij} : let us consider $\mathbf{x}_0 \neq \mathbf{x}_1 \in \Gamma_{ij}$. Locally at both \mathbf{x}_0 and \mathbf{x}_1 , Γ_{ij} is a smooth curve, and since ω_i and ω_j are open and (path-)connected, we can easily construct two non self-intersecting curves $\gamma_i : [0, 1] \rightarrow \text{supp}(u_i)$ and $\gamma_j : [0, 1] \rightarrow \text{supp}(u_j)$ such that $\gamma_i(0, 1) \subset \omega_i$ and $\gamma_j(0, 1) \subset \omega_j$ and, moreover, $\gamma_i(0) = \gamma_j(1) = \mathbf{x}_0$ and $\gamma_i(1) = \gamma_j(0) = \mathbf{x}_1$. It follows by construction that on the Jordan's curve $\gamma := \{\mathbf{x}_0\} \cup \gamma_i \cup \{\mathbf{x}_1\} \cup \gamma_j$ only u_i and u_j are non trivial. Calling Σ the open, simply connected region enclosed by γ , then, by the maximum principle, all the other functions are trivial when restricted to Σ . It follows in particular that $\Delta(a_{ji}u_i - a_{ij}u_j) = 0$ in Σ , and it vanishes on $\partial\Sigma$ exactly at \mathbf{x}_0 and \mathbf{x}_1 . Standard properties of harmonic functions imply that $\Gamma_{ij} \cap \Sigma$ is connected.

We are left to show that if $\Gamma_{ij} \neq \emptyset$, then $\omega_i \cup \Gamma_{ij} \cup \omega_j$ is an open and simply connected subset of Ω , but this is an immediate consequence of the above construction. \square

The Hopf lemma implies that, for any i , some Γ_{ij} must be non-empty.

Lemma 2.7. *Let $B_r(\mathbf{x}_0) \subset \omega_i$ and $\mathbf{p} \in \partial B_r(\mathbf{x}_0) \cap \partial\omega_i$. Then $m(\mathbf{p}) = 2$.*

Proof. Hopf's lemma forces $\nabla u_i(\mathbf{p}) \neq 0$, and Lemma 1.4 yields $\mathbf{p} \in \mathcal{Z}_2$. \square

Turning to the analysis of higher multiplicity points, we first show that the non-degeneracy assumption (A4) implies that the singular set \mathcal{W} lies in the interior of Ω . We need a preliminary result.

Lemma 2.8. *Let $\mathbf{x}_0 \in \partial\Omega$ be such that, for some $i \neq j$, $\mathbf{x}_0 \in \overline{\{x : \varphi_i(x) > 0\}} \cap \overline{\{x : \varphi_j(x) > 0\}}$, and let $\boldsymbol{\nu}$ denote the exterior normal unit vector to $\partial\Omega$ at \mathbf{x}_0 . Then:*

1. *there exist a unit vector \mathbf{e} , with $-1/2 < \mathbf{e} \cdot \boldsymbol{\nu} < 0$, and positive constants α, L , such that $\alpha t \leq u_i(\mathbf{x}_0 + t\mathbf{e}) \leq Lt$, for $t > 0$ sufficiently small;*
2. *there exist constants $M > 0, \gamma > 1$ such that $|u_h(\mathbf{x})| \leq M|\mathbf{x} - \mathbf{x}_0|^\gamma$ for every $h \neq i, j$;*
3. *if $\Gamma_{ij} = \emptyset$ then there exists $\rho > 0$ sufficiently small such that $(B_\rho(\mathbf{x}_0 + \rho\mathbf{e}) \cap \{\mathbf{x} : (\mathbf{x} - \mathbf{x}_0) \cdot \mathbf{e}^\perp > 0\}) \subset \omega_i$, where the unit vector \mathbf{e}^\perp , orthogonal to \mathbf{e} , is chosen such that $\mathbf{e}^\perp \cdot \boldsymbol{\nu} < 0$ (see Fig. 2).*




Figure 2: the half-disk $B_\rho(\mathbf{x}_0 + \rho\mathbf{e}) \cap \{\mathbf{x} : (\mathbf{x} - \mathbf{x}_0) \cdot \mathbf{e}^\perp > 0\}$, contained in ω_i when Γ_{ij} is empty and ρ is small (Lemma 2.8).

Proof. We assume w.l.o.g. $\mathbf{x}_0 = (0, 0)$. All the following arguments are understood as local (near 0).

We start by proving 1. Recall that, by Lemma 1.4, U is Lipschitz with constant, say, $L > 0$. Since $U(\mathbf{x}_0) = 0$, this immediately yields $u_h(\mathbf{x}) \leq L|\mathbf{x} - \mathbf{x}_0|$, for every h and \mathbf{x} . On the other hand, since $\partial\Omega$ is of class $C^{1,1}$, we can assume that $\Omega = \{(x, y) \in \mathbb{R}^2 : y > f(x)\}$, with $f \in C^1$ and $f(0) = f'(0) = 0$. Assumption (A4) implies, for instance, $\varphi_i(x, f(x)) > Cx$ (resp. $\varphi_j(x, f(x)) > -Cx$) for $x > 0$ (resp. $x < 0$). Then, for any $m = \tan \theta > 0$ sufficiently small, there exists $\alpha > 0$ such that

$$u_i(x, mx) \geq u_i(x, f(x)) - L|mx - f(x)| \geq (C - Lm)x + o(x) \geq \alpha x, \quad \text{for } x > 0 \text{ small,} \quad (8)$$

and 1 follows, with $\mathbf{e} = (\cos \theta, \sin \theta)$. Note that, at least for a smaller m , a similar estimate holds true also for u_j :

$$u_j(x, -mx) \geq -\alpha x, \quad \text{for } x < 0 \text{ small.}$$

From the previous point we have that (locally near 0) ω_h is contained in the angle $\{(x, y) : y \geq m|x|\}$ whenever $h \neq i, j$. Standard comparison arguments with the positive harmonic function of a cone yield point 2, with $C = \|U\|_{L^\infty}$ and $\gamma = \pi/(\pi - 2\theta)$.

Finally, let $\rho > 0$ small to be fixed. Notice that, choosing $\delta = \alpha \cos \theta / (2L)$, we have

$$u_i|_{\partial B_{\delta\rho}(\rho\mathbf{e}) \cap \Omega} \geq u_i(\rho\mathbf{e}) - L\delta\rho \geq \frac{\alpha}{2}\rho \cos \theta \quad (9)$$

(in particular, $0 < \delta < 1$). Let us consider the number

$$\bar{r} := \sup\{r > 0 : (B_r(\rho\mathbf{e}) \cap \{(x, y) : y > mx\}) \subset \omega_i\}.$$

On the one hand, by (9), we have that $\bar{r} > \delta\rho$. On the other hand, since $\mathbf{x}_0 = 0 \notin \omega_i$, it holds $\bar{r} \leq \rho$. Since 3 is equivalent to $\bar{r} = \rho$, to conclude we assume by contradiction that $\bar{r} < \rho$. This implies the existence of an index $h \neq i$ and of a point $\mathbf{p} \in \bar{\omega}_i \cap \bar{\omega}_h$, with $\mathbf{p} \in \Omega$ and $|\rho\mathbf{e} - \mathbf{p}| = \bar{r}$. We have that $B_{\bar{r}}(\rho\mathbf{e})$ is both an interior (half-)ball touching at \mathbf{p} for ω_i , and an exterior one for ω_h . In particular, by Lemma 2.7 we infer that $m(\mathbf{p}) = 2$; then Lemma 1.4 implies

$$\lim_{\substack{\mathbf{x} \rightarrow \mathbf{x}_0 \\ \mathbf{x} \in \{u_i > 0\}}} \nabla u_i(\mathbf{x}) = -\frac{a_{ih}}{a_{hi}} \lim_{\substack{\mathbf{x} \rightarrow \mathbf{x}_0 \\ \mathbf{x} \in \{u_h > 0\}}} \nabla u_h(\mathbf{x}), \quad (10)$$

where, since $\Gamma_{ij} = \emptyset$, $h \neq j$.

Now, by construction

$$A_1 := ((B_{\bar{r}}(\rho\mathbf{e}) \setminus B_{\delta\rho}(\rho\mathbf{e})) \cap \{(x, y) : y > mx\}) \subset \omega_i;$$

recalling (8) and (9), the function

$$\eta_1(\mathbf{x}) = \frac{\log \bar{r} - \log |\mathbf{x} - \rho\mathbf{e}|}{\log \bar{r} - \log \delta\rho} \frac{\alpha}{2} \rho \cos \theta \quad \text{satisfies} \quad \begin{cases} -\Delta \eta_1 = 0 & \text{in } A_1 \\ \eta_1 \leq u_i & \text{on } \partial A_1, \end{cases}$$

so that

$$\lim_{\substack{\mathbf{x} \rightarrow \mathbf{p} \\ \mathbf{x} \in \omega_i}} |\nabla u_i(\mathbf{x})| \geq |\nabla \eta_1(\mathbf{p})| = \frac{1}{\log \bar{r} / (\delta\rho)} \cdot \frac{1}{\bar{r}} \cdot \frac{\alpha}{2} \rho \cos \theta \geq \frac{\alpha \cos \theta}{-2 \log \delta} > 0, \quad (11)$$

independently of ρ . On the other hand, recalling point 2, we can easily construct a barrier from above for u_h . Indeed, let

$$A_2 := (B_{2\bar{r}}(\rho\mathbf{e}) \setminus B_{\bar{r}}(\rho\mathbf{e})) \cap \Omega, \quad \eta_2(\mathbf{x}) = \frac{\log 2\bar{r} - \log |\mathbf{x} - \rho\mathbf{e}|}{\log 2} M(2\rho)^\gamma;$$

then

$$\begin{cases} -\Delta u_h \leq 0 = -\Delta \eta_2 & \text{in } A_2 \\ u_h \leq \eta_2 & \text{on } \partial A_2. \end{cases}$$

We deduce that

$$\lim_{\substack{\mathbf{x} \rightarrow \mathbf{p} \\ \mathbf{x} \in \omega_h}} |\nabla u_h(\mathbf{x})| \leq |\nabla \eta_2(\mathbf{p})| = \frac{1}{\log 2} \cdot \frac{1}{\bar{r}} \cdot M(2\rho)^\gamma \leq \frac{M2^\gamma}{\delta \log 2} \rho^{\gamma-1},$$

which is in contradiction, when ρ is sufficiently small, with (11) and (10). \square

Corollary 2.9. *Let $\mathbf{x}_0 \in \overline{\{x : \varphi_i(x) > 0\}} \cap \overline{\{x : \varphi_j(x) > 0\}}$. Then $\Gamma_{ij} \neq \emptyset$.*

Proof. Applying the above lemma twice (the second time exchanging the role of i and j), we obtain the existence of e, e', ρ, ρ' such that $-1 < e \cdot e' < 0$ and

$$(B_\rho(\mathbf{x}_0 + \rho e) \cap \{\mathbf{x} : (\mathbf{x} - \mathbf{x}_0) \cdot e^\perp > 0\}) \subset \omega_i, \quad (B_{\rho'}(\mathbf{x}_0 + \rho' e') \cap \{\mathbf{x} : (\mathbf{x} - \mathbf{x}_0) \cdot (e')^\perp > 0\}) \subset \omega_j,$$

a contradiction since

$$(B_\rho(\mathbf{x}_0 + \rho e) \cap \{\mathbf{x} : (\mathbf{x} - \mathbf{x}_0) \cdot e^\perp > 0\}) \cap (B_{\rho'}(\mathbf{x}_0 + \rho' e') \cap \{\mathbf{x} : (\mathbf{x} - \mathbf{x}_0) \cdot (e')^\perp > 0\}) \neq \emptyset. \quad \square$$

Lemma 2.10. *Let $\mathbf{x}_0 \in \partial\Omega$. Then either $m(\mathbf{x}_0) = 1$ or $m(\mathbf{x}_0) = 2$. In particular, $\overline{\mathcal{W}} \subset \Omega$.*

Proof. If $\varphi_i(\mathbf{x}_0) > 0$ for some i , then $m(\mathbf{x}_0) = 1$. On the other hand, let $\varphi_i(\mathbf{x}_0) = 0$ for every i . Note that assumption (A4) implies that \mathbf{x}_0 is an isolated zero of the trace function φ . We deduce the existence of two points $\mathbf{x}_\pm \in \partial\Omega$ with the properties that φ is strictly positive in \mathbf{x}_\pm and also on the curve $\overline{\mathbf{x}_- \mathbf{x}_0}, \overline{\mathbf{x}_0 \mathbf{x}_+}$ (here we denote with $\overline{\mathbf{x}_1 \mathbf{x}_2}$ the relatively open portion of $\partial\Omega$ having counterclockwise ordered endpoints \mathbf{x}_1 and \mathbf{x}_2 , respectively). Two cases may occur.

Case 1: both $\mathbf{x}_\pm \in \{\varphi_i > 0\}$, for some i . For $r > 0$ sufficiently small we have $B_r(\mathbf{x}_-) \cap \Omega \subset \omega_i$, $B_r(\mathbf{x}_+) \cap \Omega \subset \omega_i$. Recalling that ω_i is pathwise connected (Lemma 2.3), we can find a path γ such that $\gamma(0) = \mathbf{x}_+$, $\gamma(1) = \mathbf{x}_-$ and $\gamma(0, 1) \subset \omega_i$. Let us denote with Σ the bounded connected component of $\mathbb{R}^2 \setminus (\gamma([0, 1]) \cup \overline{\mathbf{x}_- \mathbf{x}_+})$. Then we have that the only nontrivial density on $\partial\Sigma$, and hence on Σ is u_i , so that $m(\mathbf{x}_0) = 1$.

Case 2: $\mathbf{x}_- \in \{\varphi_i > 0\}$, $\mathbf{x}_+ \in \{\varphi_j > 0\}$, with $j \neq i$. Then Corollary 2.9 applies, and $\Gamma_{ij} \neq \emptyset$. Consequently, this case can be treated in a similar way than the previous one, with the only difference that now γ can be constructed in such a way that $\gamma(0, 1) \subset \omega_i \cup \Gamma_{ij} \cup \omega_j$ (which is pathwise connected by Lemma 2.6). Then u_h vanishes in $\overline{\Sigma}$, for every $h \neq i, j$, and $m(\mathbf{x}_0) = 2$. \square

Notice that the interior of \mathcal{W} , which may or may not be empty, coincides with \mathcal{Z}_0 (by definition of multiplicity). On the other hand, by the Hopf lemma points of higher multiplicity are limit of regular zeroes. We make this more precise in the next two lemmas.

Lemma 2.11. *The boundary of \mathcal{W} is the accumulation set of \mathcal{Z}_2 :*

$$\partial^* \mathcal{W} \subset \bigcup_{i \neq j} \overline{\Gamma}_{ij}.$$

Proof. Let $\mathbf{x}_0 \in \partial^* \mathcal{W}$, we need to show that for all $r > 0$ there exists $\mathbf{p} \in B_r(\mathbf{x}_0) \cap \mathcal{Z}_2$. Fixing $r > 0$, since $\mathbf{x}_0 \in \partial \mathcal{W}$ there exists $\mathbf{x}' \in B_r(\mathbf{x}_0) \setminus \mathcal{W}$. Then either $\mathbf{x}' \in \mathcal{Z}_2$ and we are done, or $u_i(\mathbf{x}') > 0$ for some index i . In the latter case, let $R > 0$ be such that $B_R(\mathbf{x}') \subset \{u_i > 0\} \cap B_r(\mathbf{x}_0)$ and let us consider the segment $t \in [0, 1] \mapsto (1-t)\mathbf{x}' + t\mathbf{x}_0$. Since $\{u_i > 0\} \cap B_r(\mathbf{x}_0)$ is an open set, there exists a maximum value $\bar{t} \in [0, 1]$ such that

$$B_R((1-t)\mathbf{x}' + t\mathbf{x}_0) \subset \{u_i > 0\} \quad \text{for all } t \in [0, \bar{t}].$$

Consequently $B_R((1-\bar{t})\mathbf{x}' + \bar{t}\mathbf{x}_0)$ satisfies the assumptions of Lemma 2.7. \square

Lemma 2.12. For any non empty Γ_{ij} , each of its limit sets is either a point of $\partial\Omega$ or a connected subset of \mathcal{W} .

Proof. Being Γ_{ij} a locally smooth (non self-intersecting) curve contained in Ω , we immediately obtain that it admits a global one-to-one parametrization $\phi \in C^1(I; \Omega)$, for some interval $I \subset \mathbb{R}$: for instance, we can take ϕ as a solution to the system

$$\begin{cases} \dot{\phi}(t) = J\nabla(a_{ji}u_i - a_{ij}u_j)(\phi(t)) \\ \phi(0) = \mathbf{x}_0 \end{cases}$$

where J is the symplectic matrix and $\mathbf{x}_0 \in \Gamma_{ij}$ is any point. Since in the set $\omega_i \cup \Gamma_{ij} \cup \omega_j$ the function $a_{ji}u_i - a_{ij}u_j$ is locally smooth and has non vanishing gradient, we obtain the existence of $-\infty \leq a < 0 < b \leq +\infty$, the maximal times of definition of ϕ , and of the α - and ω -limit sets of ϕ :

$$\alpha(\phi) = \bigcap_{a < t < 0} \overline{\phi(a, t)}, \quad \omega(\phi) = \bigcap_{b > t > 0} \overline{\phi(t, b)}.$$

It is easy to check that both limits are non-empty, closed, connected subsets of $\overline{\Omega}$. We consider the set $\omega(\phi)$, the other is analogous. Let $\mathbf{x}_0 \in \omega(\phi)$. By construction we have that $m(\mathbf{x}_0) \geq 2$. Hence, either $\omega(\phi) \subset \mathcal{W}$, or $m(\mathbf{x}_0) = 2$. In the latter case, we see that $\mathbf{x}_0 \notin \Omega$, otherwise we may solve the above Cauchy problem in $B_r(\mathbf{x}_0)$, contradicting the maximality of b . Therefore $\mathbf{x}_0 \in \partial\Omega$, and Lemma 2.10 forces $\omega(\phi) \equiv \mathbf{x}_0$. \square

Finally, we are in a position to introduce the cut procedures which will yield Proposition 2.1.

Lemma 2.13. We can reduce the problem to the case in which, for each $i = 1, \dots, k$, $\text{supp}(u_i) \cap \partial\Omega$ is a connected curve. Furthermore, we can assume that any curve Γ_{ij} reaches the boundary at most once, there are at least three non trivial densities in Ω and \mathcal{W} is not empty.




Figure 3: splitting for Lemma 2.13: on the left, the original domain; on the right, the split components.

Proof. By assumption (A4) and Lemma 2.10, $\text{supp}(u_i) \cap \partial\Omega$ has a finite number of connected components, on each of which u_i does not identically vanish. If, say, $\text{supp}(u_1) \cap \partial\Omega$ contains more than one connected component, let \mathbf{x}_0 and \mathbf{x}_1 be any two points that belong to different connected components, with $u_1(x_i) > 0$. Let $\gamma : [0, 1] \rightarrow \text{supp}(u_1)$ be a smooth simple curve such that $\gamma(0, 1) \subset \omega_1$, $\gamma(0) = \mathbf{x}_0$ and $\gamma(1) = \mathbf{x}_1$: γ cuts the domain Ω in two subdomains, on which the number of connected components of $\text{supp}(u_1) \cap \partial\Omega$ has reduced by at least one. Moreover, the two subdomains are regular except for two corner points, in \mathbf{x}_1 and \mathbf{x}_2 . Since $u_1(x_i) > 0$, one can easily cut neighborhoods of x_i in such a way that

the new subdomains are smooth, and all the assumptions hold, in particular assumption (A4) because U is Lipschitz (see Fig. 3). Iterating the previous construction a finite number of times we can assume that each $\text{supp}(u_i) \cap \partial\Omega$ is a connected curve.

Next, let us assume that $\bar{\Gamma}_{12} \cap \partial\Omega = \{\mathbf{y}_0, \mathbf{y}_1\}$. Note that $\mathbf{y}_0 \neq \mathbf{y}_1$. By construction, each of the two connected components of $\partial\Omega \setminus \{\mathbf{y}_0, \mathbf{y}_1\}$ must coincide with $\text{supp}(u_i) \cap \partial\Omega$ for either $i = 1$ or $i = 2$. We deduce that $k = 2$, $\mathcal{W} = \emptyset$ and $\mathcal{Z} \cap \Omega = \mathcal{Z}_2 \cap \Omega = \Gamma_{12} \cap \Omega$, thus the regularity of the free boundary follows from the previous discussion (actually, it is the zero set of the harmonic function $a_{ji}u_i - a_{ij}u_j$). Therefore we are left to deal with the case $k \geq 3$, in which each Γ_{ij} has at least one limit in \mathcal{W} . \square

Lemma 2.14. *Up to a further reduction, we can label the densities u_i in a counterclockwise sense, in such a way that $\Gamma_{ij} \neq \emptyset$ if and only if $j - i = \pm 1 \pmod k$, and both $\bar{\Gamma}_{ij} \cap \partial\Omega$ and $\bar{\Gamma}_{ij} \cap \mathcal{W}$ are non-empty.*




Figure 4: splitting for Lemma 2.14: on the left, the original domain; on the right, the split components.

Proof. Taking into account Lemma 2.13, the lemma will follow once we show that Ω can be decomposed in such a way that (in each subdomain) $\Gamma_{ij} \neq \emptyset$ if and only if $\text{supp } \varphi_i \cap \text{supp } \varphi_j \neq \emptyset$. Let $\Gamma_{ij} \neq \emptyset$ and $\text{supp } \varphi_i \cap \text{supp } \varphi_j = \emptyset$; Lemmas 2.3 and 2.6 imply that $\{\varphi_i > 0\} \cap \omega_i \cap \Gamma_{ij} \cap \omega_j \cap \{\varphi_j > 0\}$ is pathwise connected. As a consequence, reasoning as in the previous lemma, we can construct a smooth simple curve $\gamma : [0, 1] \rightarrow \text{supp}(u_i) \cup \text{supp}(u_j)$ with $\varphi_i(\gamma(0)) > 0$, $\varphi_j(\gamma(1)) > 0$, and which intersects Γ_{ij} just once (and transversally). Again, γ cuts the domain Ω in two subdomains, on both of which $\text{supp } u_i$ intersects $\text{supp } u_j$ also on the boundary. As before, the two subdomains are regular except for two corner points, which can be treated cutting some neighborhoods (see Fig. 4). Iterating the procedure a finite number of times, and relabelling the components, the lemma follows. \square

Lemma 2.15. *Under the previous reductions, the set \mathcal{Z} is connected.*

Proof. Let us assume that \mathcal{Z} is not connected. Then there exists a Jordan curve γ that separates \mathcal{Z} in two components, and in particular $\gamma \cap \mathcal{Z} = \emptyset$. Since $\mathcal{Z} \cup \partial\Omega$ is connected (Lemma 2.5), γ must cross $\partial\Omega$ in at least two points, and we can assume that γ has a finite (even) number of transverse intersections with $\partial\Omega$. Let $t_0 < t_1 \in [0, 1]$ be such that

$$\gamma(t_0), \gamma(t_1) \in \partial\Omega, \quad \gamma(t) \in \Omega \quad \text{for } t \in (t_0, t_1);$$

from the discussion above, there exists i such that $\gamma([t_0, t_1]) \in \text{supp}(u_i)$ (otherwise γ would intersect \mathcal{Z}). Let σ be the component of $\partial\Omega \setminus \{\gamma(t_0), \gamma(t_1)\}$ that is contained in $\text{supp}(u_i)$ (which exists by Lemma 2.13): the simple closed curve $\gamma([t_0, t_1]) \cup \sigma$ does not intersect \mathcal{Z} , and thus we can deform continuously γ into $\gamma([0, t_0]) \cup \sigma \cup \gamma([t_1, 1])$ without ever crossing \mathcal{Z} . Iterating these steps a finite number of times, we end up with a new Jordan curve γ' that is by construction homotopic to γ in $\mathbb{R}^2 \setminus \mathcal{Z}$ and such that $\gamma' \cap \Omega = \emptyset$. In particular, no point of \mathcal{Z} is contained in the unbounded portion of $\mathbb{R}^2 \setminus \gamma'$, a contradiction. \square

Lemma 2.16. *Under the previous reductions, the set \mathcal{W} is connected.*

Proof. Assume by contradiction that \mathcal{W} is the disjoint union of the two closed sets \mathcal{W}_1 and \mathcal{W}_2 . By Lemmas 2.14, 2.12 and 2.10 we have that, for each i , either $\bar{\Gamma}_{(i-1)i} \cap \mathcal{W}_1 = \emptyset$ or $\bar{\Gamma}_{(i-1)i} \cap \mathcal{W}_2 = \emptyset$ (here Γ_{01} stands for Γ_{1k}). We deduce that \mathcal{Z} is the disjoint union of

$$\mathcal{W}_1 \cup \left\{ \bigcup_i \Gamma_{(i-1)i} : \bar{\Gamma}_{(i-1)i} \cap \mathcal{W}_2 = \emptyset \right\} \quad \text{and} \quad \mathcal{W}_2 \cup \left\{ \bigcup_i \Gamma_{(i-1)i} : \bar{\Gamma}_{(i-1)i} \cap \mathcal{W}_1 = \emptyset \right\},$$

in contradiction with Lemma 2.15. \square

Proof of Proposition 2.1. Using Lemmas 2.3, 2.10, 2.13, 2.14 and 2.16 we obtain that Ω splits, up to some residual set of $\Omega \setminus \mathcal{Z}$ (recall Figs. 3, 4) into the finite union of domains satisfying all the required properties in (MCS), but the last. By the Riemann Mapping Theorem we can assume that, up to a conformal change of coordinates, the domain is the unit ball $B = B_1(\mathbf{0}) \subset \mathbb{R}^2$. With an abuse of notation, we keep writing $u_i, \omega_i, \mathcal{Z}, \mathcal{W}$, and so on, for the corresponding transformed objects, defined in B instead of Ω . Since Ω is of class $C^{1,\alpha}$, we have that the conformal transformation can be extended to a $C^{1,\alpha}(\bar{B})$ map onto $\bar{\Omega}$ (see, e.g., [19, Thm. 3.6]). We deduce that assumption (A4) holds true also for the transformed densities. Now, if $\mathcal{W} \equiv \{\mathbf{0}\}$ then the proposition follows; in the other case, let $r > 0$ be such that

$$\mathcal{W} \subset \bar{B}_r(\mathbf{0}), \quad \mathcal{W} \cap \partial B_r(\mathbf{0}) \ni \mathbf{x}_0,$$

where $r < 1$ by Lemma 2.10. Using as a second conformal change of variables the Möbius transform which sends $\mathbf{0}$ to \mathbf{x}_0 (and B onto itself) we have that the pre-image of $B_r(\mathbf{0})$ is contained in a ball touching the origin, concluding the proof. \square

Remark 2.17. We stress that, under the conformal changes of variables we introduced in the proof above, the local expansion of any transformed function near the origin is the same than that of the original one near \mathbf{x}_0 , up to a similarity transformation (this will provide the asymptotic expansion (5)).

Once we reduced to work on a (finite number of domains) Ω satisfying (MCS), we conclude this section by introducing a suitable conformal map, which provides an equivalent formulation of our problem for a function defined in the half-plane. By construction, we have that $\omega_i \cap \partial B$ is a connected curve on ∂B for every index i , while each non-empty Γ_{ij} consists of a smooth curve going from ∂B to \mathcal{W} . Furthermore, the densities u_1, \dots, u_k are ordered in counterclockwise order around the origin.

Let

$$\mathcal{U}(\mathbf{x}) = \begin{cases} u_1(\mathbf{x}) & \text{if } \mathbf{x} \in \text{supp}(u_1), \\ (-1)^{i-1} \left(\prod_{j=2}^i \frac{a_{(j-1)j}}{a_{j(j-1)}} \right) u_i(\mathbf{x}) & \text{if } \mathbf{x} \in \text{supp}(u_i), i = 2, \dots, k. \end{cases} \quad (12)$$

Using Lemma 1.4, we obtain that

$$\mathcal{U} \in C^1(B \setminus (\Gamma_{1k} \cup \mathcal{W})), \quad \Delta \mathcal{U} = 0 \text{ in } B \setminus (\Gamma_{1k} \cup \mathcal{W}).$$

For concreteness, from now on we assume that k is even, the odd case following with minor changes, see Remark 2.18 below. Let

$$\lambda := \frac{a_{k1}}{a_{1k}} \cdot \prod_{j=2}^k \frac{a_{(j-1)j}}{a_{j(j-1)}} > 0. \quad (13)$$

Using again Lemma 1.4 we infer that, for every $x \in \Gamma_{1k}$,

$$\lim_{\substack{\mathbf{y} \rightarrow \mathbf{x} \\ \mathbf{y} \in \omega_k}} \nabla \mathcal{U}(\mathbf{y}) = \prod_{j=1}^k \frac{a_{(j-1)j}}{a_{j(j-1)}} \lim_{\substack{\mathbf{y} \rightarrow \mathbf{x} \\ \mathbf{y} \in \omega_k}} \nabla u_k(\mathbf{y}) = \lambda \lim_{\substack{\mathbf{y} \rightarrow \mathbf{x} \\ \mathbf{y} \in \omega_1}} \nabla u_1(\mathbf{y}) = \lambda \lim_{\substack{\mathbf{y} \rightarrow \mathbf{x} \\ \mathbf{y} \in \omega_1}} \nabla \mathcal{U}(\mathbf{y}). \quad (14)$$

Now, if $\lambda \neq 1$ then \mathcal{U} is not harmonic on $B \setminus \mathcal{W}$. On the other hand, for any λ , we can construct a harmonic function by lifting \mathcal{U} to the universal covering of $B \setminus \mathcal{W}$. More precisely, we consider the conformal map

$$\mathcal{T} : \mathbb{R}_+^2 := \{(x, y) \in \mathbb{R}^2 : y > 0\} \rightarrow B \setminus \{\mathbf{0}\}, \quad \mathcal{T} : (x, y) \mapsto \mathbf{x} = (e^{-y} \cos x, e^{-y} \sin x). \quad (15)$$

Notice that \mathcal{T} is a universal covering. Up to a rotation, let us assume that the endpoint on ∂B of Γ_{1k} is the point of coordinates $(1, 0)$. The Lifting Theorem provides the existence of a unique regular curve $\Gamma \subset \overline{\mathbb{R}_+^2}$, containing $(0, 0)$, which lifts Γ_{1k} . Also $\Gamma + (2\pi, 0)$ (the horizontal translated of Γ) is a regular curve, it starts from $(2\pi, 0)$, and it has empty intersection with Γ . Noticing that $B \setminus (\Gamma_{1k} \cup \mathcal{W})$ is simply connected (this can be shown as in Lemma 2.6), we can define the set S as the unique lifting of $B \setminus (\Gamma_{1k} \cup \mathcal{W})$ such that

$$\partial S \supset [\Gamma \cup (\Gamma + (2\pi, 0)) \cup ([0, 2\pi] \times \{0\})]. \quad (16)$$

Since \mathcal{T} is conformal, the function

$$v(x, y) := \mathcal{U}(\mathcal{T}(x, y)) \quad (17)$$

is harmonic on S , and

$$\int_S |\nabla v|^2 dx dy < +\infty.$$

Using (14), we can extend v in such a way that

$$v(x + 2\pi, y) = \lambda v(x, y),$$

so that v is harmonic in the whole $\mathbb{R}_+^2 \setminus \mathcal{T}^{-1}(\mathcal{W} \setminus \{\mathbf{0}\})$, and continuous in $\overline{\mathbb{R}_+^2}$. Resuming, v is a solution of the problem

$$\begin{cases} \Delta v = 0 & \text{in } \mathbb{R}_+^2 \setminus \mathcal{T}^{-1}(\mathcal{W} \setminus \{\mathbf{0}\}) \\ v \equiv 0 & \text{in } \mathcal{T}^{-1}(\mathcal{W} \setminus \{\mathbf{0}\}) \\ v(x, 0) = v_0(x) & x \in \mathbb{R} \\ v(x + 2\pi, y) = \lambda v(x, y) \\ \int_S |\nabla v|^2 dx dy < +\infty, \end{cases} \quad (18)$$

where v_0 is a suitable combination of the trace functions φ_i .

Remark 2.18. Notice that when k is odd one can double the angle, using the map

$$\mathcal{T}_2 : (x, y) \mapsto \mathbf{x} = (e^{-2y} \cos 2x, e^{-2y} \sin 2x).$$

In such a way, we can reduce to (18), with λ^2 instead of λ .

Remark 2.19. The (multi-valued) inverse \mathcal{T}^{-1} satisfies

$$\mathcal{T}^{-1}(r \cos \vartheta, r \sin \vartheta) = (\vartheta, -\log r),$$

where we use the polar system around $\mathbf{0}$, writing $\mathbf{x} = (r \cos \vartheta, r \sin \vartheta)$. Therefore

$$\mathcal{U}(\mathbf{x}) = v(\vartheta, -\log r) \quad (19)$$

(this will provide the asymptotic expansion (5)).

3 A representation formula in \mathbb{R}_+^2

In this section we deal with problem (18), in the particular case in which $\mathcal{W} = \{0\}$ and the number of densities is even, say $2n^*$ (recall Remark 2.18). In such a case, we can assume that the curve Γ (the lifting of Γ_{1k} to the half plane $\mathbb{R}_+^2 := \{(x, y) \in \mathbb{R}^2 : y > 0\}$) has $C^{1,\alpha}$, regular parametrization

$$\Gamma := \{(x(t), y(t)) : t \in [0, +\infty)\}, \quad \text{satisfying} \quad \begin{cases} t \mapsto (x(t), y(t)) \text{ is injective} \\ (x(0), y(0)) = (0, 0), \\ y(t) > 0 \text{ for } t > 0, \\ \lim_{t \rightarrow +\infty} y(t) = +\infty, \\ \Gamma \cap (\Gamma + (2\pi, 0)) = \emptyset \end{cases}$$

By connectedness, we have that $\Gamma \cap (\Gamma + (2h\pi, 0)) = \emptyset$ too, as long as $h \in \mathbb{Z} \setminus \{0\}$. Under this perspective, the set $S \subset \mathbb{R}_+^2$ is a “strip”, i.e. the unique unbounded connected open set having boundary

$$\partial S := \Gamma \cup (\Gamma + (2\pi, 0)) \cup ([0, 2\pi] \times \{0\}).$$

Definition 3.1. We denote by

$$S_y := \{(x, y) \in S\}$$

the horizontal sections of S , having endpoints

$$X_{\min}(y) := \inf\{x : (x, y) \in S\} > -\infty, \quad X_{\max}(y) := \sup\{x : (x, y) \in S\} < +\infty,$$

and diameter

$$\text{diam } S_y = X_{\max}(y) - X_{\min}(y),$$

respectively (see Figure 5). Of course, $(X_{\min}(y), y) \in \Gamma$ and $(X_{\max}(y), y) \in \Gamma + (2\pi, 0)$.




Figure 5: possible behavior of S . The thick horizontal segments correspond to the connected components of $S_{\bar{y}}$.

We observe that, though in general the diameter of S_y may be arbitrarily large, as well as its number of connected components, in any case its length is bounded by 2π . This can be easily deduced recalling that, according to Section 2, S is a covering of the disk minus a simple curve connecting the boundary to the center. We provide a short, self-contained proof, which will be useful in the following.

Lemma 3.2. For every y , $|S_y| \leq 2\pi$.

Proof. For any $y > 0$, we have that S_y is open in the topology of the horizontal line having height y , and therefore it is the disjoint union of at most countable open intervals:

$$S_y = \bigcup_{i=1}^k (s_{2i-1}, s_{2i}) \times \{\bar{y}\}, \quad k \leq +\infty. \quad (20)$$

Note that if $x_1, x_2 \in \bar{S}_y$ and $x_2 - x_1 = 2h\pi$, with $h \in \mathbb{Z}$, then necessarily either $h = 0$, or both $(x_i, y) \in \partial S$, $h = \pm 1$; indeed, by translation, we can assume w.l.o.g. that, say, $(x_1, y) \in \partial S$, $(x_2, y) \in \bar{S}$, so that $(x_1, y) \in \Gamma + (2j\pi, 0)$, with $j \in \{0, 1\}$, and

$$(x_2, y) = (x_1 + 2h\pi, y) \in \Gamma + (2(j+h)\pi, 0),$$

and this curve does not intersect \bar{S} if $j+h \notin \{0, 1\}$.

We deduce that, for every i , there exists $h(i) \in \mathbb{N}$ such that

$$(\hat{s}_{2i-1}, \hat{s}_{2i}) := (s_{2i-1} + 2h(i)\pi, s_{2i} + 2h(i)\pi) \subset (X_{\min}(y), X_{\min}(y) + 2\pi),$$

and $(\hat{s}_{2i-1}, \hat{s}_{2i}) \cap (\hat{s}_{2j-1}, \hat{s}_{2j}) = \emptyset$ for $i \neq j$. Thus

$$|S_y| = \left| \bigcup_{i=1}^k (\hat{s}_{2i-1}, \hat{s}_{2i}) \right| \leq |(X_{\min}(y), X_{\min}(y) + 2\pi)| = 2\pi. \quad \square$$

As we mentioned, since $\mathcal{W} \setminus \{0\}$ is empty, (18) reduces to

$$\begin{cases} \Delta v = 0 & \text{in } \mathbb{R}_+^2 \\ v = 0 & \text{on } \Gamma \\ v(x + 2\pi, y) = \lambda v(x, y) \\ \int_S |\nabla v|^2 dx dy < +\infty, \end{cases} \quad (21)$$

where $\lambda > 0$ is fixed constant, and we can assume w.l.o.g. that v is of class C^2 up to $\{y = 0\}$ (possibly replacing it with the restriction on \mathbb{R}_+^2 of $v(x, y + \varepsilon)$, $\varepsilon > 0$ small).

Remark 3.3. Since $|S_y| \leq 2\pi$, and v vanishes at the endpoints of any connected component of $|S_y|$, we readily infer the validity of a Poincaré inequality for v in S :

$$\int_{S_y} v^2 dx \leq 4 \int_{S_y} |\nabla v|^2 dx \text{ for every } y, \quad \int_S v^2 dx dy \leq 4 \int_S |\nabla v|^2 dx dy < +\infty.$$

Furthermore, we can apply the standard trace inequality on the half plane $\{y \geq \bar{y}\}$ to the function $v|_S$ (with null extension outside S) in order to obtain

$$\int_{S_{\bar{y}}} v^2 dx \leq 2 \|v|_S\|_{H^1(\mathbb{R} \times (\bar{y}, +\infty))}^2 \leq 10 \int_{S \cap \{y > \bar{y}\}} |\nabla v|^2 dx dy.$$

The aim of this section is to prove the following result.

Proposition 3.4. Under the above notation, assume that $v|_S$ has exactly $2n^*$ nodal regions (as well as $v(x, 0)|_{(0, 2\pi)}$), and let us define

$$\alpha := \frac{\log \lambda}{2\pi}. \quad (22)$$

Then there exist constants q, a, b such that

$$S_y \subset \left\{ (x, y) : -\frac{\alpha}{n^*}y + q \leq x \leq -\frac{\alpha}{n^*}y + q + 2\pi + o(1) \right\},$$

and

$$v(x, y) = [a \cos(n^*x + \alpha y) + b \sin(n^*x + \alpha y) + o(1)] \exp(\alpha x - n^*y)$$

as $y \rightarrow +\infty$, uniformly in S .

To prove the proposition, the basic idea is to reduce the third condition in (21) to a periodic one, and then to use separation of variables to write the solution in Fourier series. To this aim, for concreteness from now on we assume $\lambda > 1$, so that

$$\alpha > 0.$$

The case $0 < \lambda < 1$ can be treated in the same way, while the case $\lambda = 1$ is actually easier (indeed, before lifting to \mathbb{R}_+^2 , the function \tilde{v} defined in (12) is already harmonic and bounded in $B_1 \setminus \{0\}$).

Lemma 3.5. *Let*

$$w(x, y) := e^{-\alpha x} v(x, y).$$

Then w is C^2 in $\overline{\mathbb{R}_+^2}$, 2π -periodic with respect to the x variable, and there exist (real) numbers $a_k, b_k, k \in \mathbb{Z}$, such that

$$w(x, y) = \sum_{k \in \mathbb{Z}} [a_k \cos(kx + \alpha y) + b_k \sin(kx + \alpha y)] e^{-ky}.$$

Proof. A direct calculation shows that, for every (x, y) ,

$$w(x + 2\pi, y) = e^{-\alpha(x+2\pi)} v(x + 2\pi, y) = \lambda e^{-2\alpha\pi} e^{-\alpha x} v(x, y) = w(x, y).$$

Since $\Delta(e^{\alpha x} w(x, y)) = 0$ on the half-plane, then w satisfies

$$\Delta w + 2\alpha w_x + \alpha^2 w = 0.$$

Using the periodicity of w we can write $w(x, y) = \sum_k W_k(y) e^{ikx}$; substituting we obtain

$$\sum \left(-k^2 W_k + W_k'' + 2ik\alpha W_k + \alpha^2 W_k \right) e^{ikx} = 0$$

for all (x, y) in the half-space. This implies, for every k ,

$$W_k'' - (k - i\alpha)^2 W_k = 0, \quad \text{that is} \quad W_k(y) = A_k e^{(k-i\alpha)y} + B_k e^{(-k+i\alpha)y},$$

where A_k, B_k are suitable complex constants. As a consequence

$$\begin{aligned} w(x, y) &= \sum \left[A_k e^{ky} e^{i(kx-\alpha y)} + B_k e^{-ky} e^{i(kx+\alpha y)} \right] \\ &= \sum \left[A_{-k} e^{-i(kx+\alpha y)} + B_k e^{i(kx+\alpha y)} \right] e^{-ky}. \end{aligned}$$

Since w is real, we have that A_{-k} and B_k are conjugated, and the lemma follows by choosing $a_k = A_{-k} + B_k, b_k = i(-A_{-k} + B_k)$. \square

Corollary 3.6. *If v satisfies (21) then*

$$v = v_{\text{nice}} + v_{\text{bad}},$$

where

$$\begin{aligned} v_{\text{nice}}(x, y) &= \sum_{k=0}^{+\infty} [a_k \cos(kx + \alpha y) + b_k \sin(kx + \alpha y)] e^{\alpha x - ky}, \\ v_{\text{bad}}(x, y) &= \sum_{k=1}^{+\infty} [a_{-k} \cos(kx - \alpha y) - b_{-k} \sin(kx - \alpha y)] e^{\alpha x + ky}, \end{aligned} \quad (23)$$

where the constants a_k, b_k are those in Lemma 3.5 (of course, the terminology is due to the fact that there exist points in S having vertical coordinate y arbitrarily large).

Now we want to exploit the further conditions about v to determine the constants in (23). Roughly speaking, the idea is that the condition

$$\int_S |\nabla v|^2 dx dy < +\infty$$

should annihilate the “bad” part (namely, it should imply that $a_k = b_k = 0$ for every $k < 0$), whereas the number of nodal regions of v should determine the dominant part (that is, the first nonzero a_k and b_k). Actually, the first step is not so straight: indeed, since we do not know the actual position of S , it is not trivial to exclude the integrability on S of quantities of order $e^{2(\alpha x + ky)}$, $k > 0$, even for arbitrarily large k .

To start with, we collect in the following lemma some routine consequences of the theory of Fourier series, when applied to w .

Lemma 3.7. *There exist a constant C (only depending on $\|w\|_{C^2([0, 2\pi] \times [0, \pi/(2\alpha)])}$) such that, for every $k \in \mathbb{Z}$, $k \neq 0$,*

1. $|a_k| + |b_k| \leq \frac{C}{k^2}$;
2. $\pi(a_k^2 + b_k^2)e^{-2ky} < \int_0^{2\pi} w^2(x, y) dx + C$.

Proof. Choosing $y = 0$ in the expression of w we obtain, for every $k \geq 1$,

$$|a_k + a_{-k}| = \frac{1}{\pi} \left| \int_0^{2\pi} w(x, 0) \cos kx dx \right| = \frac{1}{\pi k^2} \left| \int_0^{2\pi} w_{xx}(x, 0) \cos kx dx \right| \leq \frac{C_1}{k^2}.$$

Analogously, the choice $y = \pi/(2\alpha)$ yields

$$\left| e^{k\pi/(2\alpha)} a_{-k} - e^{-k\pi/(2\alpha)} a_k \right| = \frac{1}{\pi} \left| \int_0^{2\pi} w\left(x, \frac{\pi}{2\alpha}\right) \sin kx dx \right| \leq \frac{C_2}{k^2}.$$

We deduce

$$\begin{aligned} |a_k| &\leq \left| a_k + a_{-k} - a_{-k} + e^{-2k\pi/\alpha} a_k \right| \leq \frac{C_1 + C_2 e^{-k\pi/(2\alpha)}}{k^2} \leq \frac{C_1 + C_2}{k^2}, \\ |a_{-k}| &\leq |a_k + a_{-k} - a_k| \leq \frac{2C_1 + C_2}{k^2}. \end{aligned}$$

One can obtain analogous estimates for $|b_k|, |b_{-k}|$ similarly, thus concluding estimate 1.

For the second part, we resume the notation of the proof of Lemma 3.5. Applying Parseval's identity we obtain, for every $y \geq 0$,

$$\begin{aligned} \frac{1}{2\pi} \int_0^{2\pi} w^2(x, y) dx &= \sum_{k \in \mathbb{Z}} W_k(y) \overline{W_k(y)} \\ &= \sum_{k \in \mathbb{Z}} A_k B_{-k} e^{2ky} + A_{-k} B_k e^{-2ky} + \underbrace{A_k A_{-k} e^{-2i\alpha y} + B_k B_{-k} e^{2i\alpha y}}_{=2 \operatorname{Re} A_k A_{-k} e^{-2i\alpha y}} \\ &= \sum_{k \in \mathbb{Z}} \frac{a_{-k}^2 + b_{-k}^2}{4} e^{2ky} + \frac{a_k^2 + b_k^2}{4} e^{-2ky} \\ &\quad + \frac{a_k a_{-k} - b_k b_{-k}}{2} \cos(2\alpha y) + \frac{a_k b_{-k} + a_{-k} b_k}{2} \sin(2\alpha y), \end{aligned}$$

that is

$$\int_0^{2\pi} w^2(x, y) dx = \pi \sum_{k \in \mathbb{Z}} (a_k^2 + b_k^2) e^{-2ky} + \mathcal{R}(y),$$

where

$$|\mathcal{R}(y)| \leq 2 \sum_{k \in \mathbb{Z}} (|a_k| + |b_k|) (|a_{-k}| + |b_{-k}|) \leq C, \quad \text{for every } y,$$

by the first part of the lemma. \square

The above estimates allow to show that, in case $v_{\text{bad}} \neq 0$, then infinitely many coefficients c_k, d_k must be different from zero.

Lemma 3.8. *Let us assume that*

$$\bar{k} := \inf \{k \in \mathbb{Z} : a_k^2 + b_k^2 \neq 0\} > -\infty.$$

Then Proposition 3.4 follows, and $\bar{k} = n^* > 0$ (recall that $2n^*$ is the number of nodal regions of $v(x, 0)$ in $[0, 2\pi]$).

Proof. From (23), we infer that

$$\begin{aligned} &\left| e^{\bar{k}y} w(x, y) - [c_{\bar{k}} \cos(\bar{k}x + \alpha y) + d_{\bar{k}} \sin(\bar{k}x + \alpha y)] \right| \\ &= \left| \sum_{k \geq \bar{k}+1} [a_k \cos(kx + \alpha y) + b_k \sin(kx + \alpha y)] e^{-(k-\bar{k})y} \right|. \end{aligned}$$

In particular, since the zero sets of v and w coincide,

$$(x, y) \in \Gamma \quad \implies \quad \left| c_{\bar{k}} \cos(\bar{k}x + \alpha y) + d_{\bar{k}} \sin(\bar{k}x + \alpha y) \right| \leq C e^{-y}.$$

Since Γ is connected, we obtain that there exists $q \in \mathbb{R}$ such that for any $\varepsilon > 0$ there exists \bar{y} large such that

$$(\Gamma \cap \{y \geq \bar{y}\}) \subset \{(x, y) : q \leq \bar{k}x + \alpha y \leq q + \varepsilon\}.$$

The same holds for any nodal line of v , in particular for $\Gamma + (2\pi, 0)$. Thus we have shown that, for $y \geq \bar{y}$, S lies between the two straight lines of equations $\bar{k}x + \alpha y = q$, $\bar{k}x + \alpha y = q + 2\pi + \varepsilon$.

Let us assume by contradiction $\bar{k} < 0$. We infer that, for any $(x, y) \in S$ with $y \geq \bar{y}$ sufficiently large, x has to be positive. Since α is positive too, recalling Remark 3.3 we can write

$$\begin{aligned} +\infty &> \int_S v^2 dx dy \geq \int_{\bar{y}}^{+\infty} dy \int_{S_y} e^{2\alpha x} w^2 dx \geq \int_{\bar{y}}^{+\infty} dy \int_{S_y} w^2 dx \\ &\geq \int_{\bar{y}}^{+\infty} [(a_{-\bar{k}}^2 + b_{-\bar{k}}^2)e^{-2\bar{k}y} - C] dy, \end{aligned}$$

forcing $a_{-\bar{k}}^2 + b_{-\bar{k}}^2 = 0$, in contradiction with the definition of \bar{k} .

Therefore $\bar{k} \geq 0$, and $v = v_{\text{nice}}$ (with summation starting from $k = \bar{k}$). Recalling again that S is controlled above and below by straight lines, we finally deduce that v and the first non-zero term in v_{nice} are close each other, for y large, in the C^2 norm. But then the number of nodal zones of v in S_y , i.e. $2n^*$, must be equal to that of $\cos(kx + \alpha y)$, i.e. $2\bar{k}$, concluding the proof. \square

The previous lemma, together with Lemma 3.7, suggests that if the bad part is non-zero then the quantity $\int_0^{2\pi} w^2(x, y) dx$, as a function of y , must increase more than exponentially.

Lemma 3.9. *If there exist constants A, β and a sequence $y_n \rightarrow +\infty$ such that*

$$\int_0^{2\pi} w^2(x, y_n) dx \leq Ae^{\beta y_n},$$

then Proposition 3.4 follows.

Proof. Combining such assumption with the second estimate in Lemma 3.7 we have, for every k and n ,

$$\pi(a_k^2 + b_k^2)e^{-2ky_n} \leq Ae^{\beta y_n} + C.$$

Choosing n sufficiently large, this inequality forces $a_k = b_k = 0$ whenever $2k < -\beta$, hence Lemma 3.8 applies. \square

Since $v^2 = e^{2\alpha x} w^2$ is integrable on S , it is easy to see that the assumption of Lemma 3.9 is fulfilled when the strip S lies on the right of a fixed straight line (it is even trivial if it lies in the sector $\{x \geq 0\}$). In fact, by the trace inequality it is sufficient to assume this properties only for a sequence $(S_{y_n})_n$.

Lemma 3.10. *If there exist a constant $m \in \mathbb{R}$ and a sequence $y_n \rightarrow +\infty$ such that*

$$\frac{X_{\min}(y_n)}{y_n} \geq m,$$

then Proposition 3.4 follows.

Proof. Remark 3.3 yields, for every n ,

$$\begin{aligned} \int_0^{2\pi} w^2(x, y_n) dx &= \int_{S_{y_n}} e^{-2\alpha x} v^2 dx \leq e^{-2\alpha X_{\min}(y_n)} \int_{S_{y_n}} v^2 dx \\ &\leq e^{-2\alpha m y_n} \cdot 10 \int_{S \cap \{y > y_n\}} |\nabla v|^2 dx dy \end{aligned}$$

(recall that $\alpha \geq 0$). But then we can conclude by applying Lemma 3.9. \square

At this point, we have ruled out the case when S stays frequently on the right of some straight line. To face the complementary case, we need the following result, which can be seen as a one-phase version of the Alt-Caffarelli-Friedman monotonicity lemma, adapted to this situation.

Lemma 3.11. Let $t \mapsto (x(t), y(t))$ denote a regular parameterization of Γ . For $\bar{y} > 0$ let

$$t_1(\bar{y}) := \min\{t : y(t) = \bar{y}\}, \quad X_1(\bar{y}) := x(t_1(\bar{y})) + 2\pi,$$

so that $(X_1(\bar{y}), \bar{y})$ belongs to $\Gamma + (2\pi, 0)$ and $X_{\min}(\bar{y}) \leq X_1(\bar{y}) \leq X_{\max}(\bar{y})$.

If $X_1(\bar{y}) < 0$ then

$$\int_{S \setminus ([X_1(\bar{y}), 2\pi] \times [0, \bar{y}])} |\nabla v|^2 dx dy \leq e^{-X_1^2(\bar{y})/\bar{y}} \int_S |\nabla v|^2 dx dy.$$

Proof. For easier notation, throughout the proof of the lemma we will assume that v is truncated to 0 in $\mathbb{R}_+^2 \setminus S$. Furthermore, for $X_1(\bar{y}) < \xi < 2\pi$, we write (see Figure 6)




Figure 6: notation for the proof of Lemma 3.11 (the thick vertical segments correspond to the connected components of Γ_ξ).

$$\eta(\xi) := \max\{y(t) : x(t) + 2\pi = \xi, t < t_1(\bar{y})\} < \bar{y},$$

$$\Gamma_\xi := \{(\xi, y) \in S : y \leq \eta(\xi)\},$$

$$\Xi_\xi := \bigcup_{\xi \leq x \leq 2\pi} \Gamma_x,$$

and

$$\Phi(\xi) := \int_{S \setminus \Xi_\xi} |\nabla v|^2 dx dy = \int_{\Gamma_\xi} v v_x dy$$

(recall that v is harmonic where it is not zero). Note that

$$\Phi(\xi) = \Phi(X_1(\bar{y})) + \int_{X_1(\bar{y})}^{\xi} dx \int_{\Gamma_x} |\nabla v|^2 dy.$$

Thus Φ is absolutely continuous and, since $v = 0$ on $\partial\Gamma_\xi$, Poincaré inequality implies (for a.e. ξ)

$$\begin{aligned} \Phi'(\xi) &= \int_{\Gamma_\xi} |\nabla v|^2 dy = \int_{\Gamma_\xi} v_x^2 dy + \int_{\Gamma_\xi} v_y^2 dy \geq \int_{\Gamma_\xi} v_x^2 dy + \frac{\pi^2}{|\Gamma_\xi|^2} \int_{\Gamma_\xi} v^2 dy \\ &\geq \frac{2\pi}{|\Gamma_\xi|} \left(\int_{\Gamma_\xi} v^2 dy \right)^{1/2} \left(\int_{\Gamma_\xi} v_x^2 dy \right)^{1/2} \geq \frac{2\pi}{|\Gamma_\xi|} \int_{\Gamma_\xi} v v_x dy = \frac{2\pi}{|\Gamma_\xi|} \Phi(\xi), \end{aligned}$$

that is

$$\frac{\Phi'(\xi)}{\Phi(\xi)} \geq \frac{2\pi}{|\Gamma_\xi|}. \quad (24)$$

In view of integrating this equation, we can use Jensen's inequality to write

$$-\frac{1}{\xi} \int_\xi^0 \frac{1}{|\Gamma_x|} dx \geq \frac{1}{-\frac{1}{\xi} \int_\xi^0 |\Gamma_x| dx} = \frac{-\xi}{\text{area}(\Xi_\xi)} \geq \frac{-\xi}{2\pi\bar{y}}$$

(recall that $\Xi_\xi \subset S \cap \{y < \bar{y}\}$ and $|S_y| = 2\pi$), and therefore

$$\int_{\bar{x}}^0 \frac{2\pi}{|\Gamma_x|} dx \geq \frac{\xi^2}{\bar{y}}.$$

From (24) we deduce

$$\Phi(\xi) \leq e^{-\xi^2/\bar{y}} \Phi(0) \leq e^{-\xi^2/\bar{y}} \int_S |\nabla v|^2 dx dy,$$

and the lemma follows by choosing $\xi = X_1(\bar{y})$ and recalling that

$$S \setminus ([X_1(\bar{y}), 2\pi] \times [0, \bar{y}]) \subset S \setminus \Xi_{X_1(\bar{y})}. \quad \square$$

The previous lemma allows to treat the case when the diameter of S_y does not grow too much, for some subsequence.

Lemma 3.12. *If there exist a constant $\delta > 0$ and a sequence $y_n \rightarrow +\infty$ such that*

$$\frac{X_1(y_n) - X_{\min}(y_n)}{y_n} \leq \delta,$$

then Proposition 3.4 follows (of course, the same holds if $X_1(y_n)$ is replaced with $X_{\max}(y_n)$).

Proof. Recalling Lemma 3.10 we can assume $X_{\min}(y_n)/y_n \rightarrow -\infty$ so that, in particular, $X_1(y_n)$ must be negative for n large. But then Lemma 3.11 applies, and we have

$$\begin{aligned} \int_0^{2\pi} w^2(x, y_n) dx &\leq e^{-2\alpha X_{\min}(y_n)} \int_{S_{y_n}} v^2 dx \leq 10e^{-2\alpha X_{\min}(y_n)} \int_{S \cap \{y > y_n\}} |\nabla v|^2 dx dy \\ &\leq C \exp\left(-2\alpha X_{\min}(y_n) - \frac{X_1^2(y_n)}{y_n}\right) \leq C \exp\left(2\alpha\delta - 2\alpha \frac{X_1(y_n)}{y_n} - \frac{X_1^2(y_n)}{y_n^2}\right) y_n \\ &\leq C \exp(2\alpha\delta + \alpha^2) y_n, \end{aligned}$$

and again we can conclude by Lemma 3.9. \square

Summarizing, we are left to consider the case when, for any sequence $y_n \rightarrow +\infty$, it holds

$$\frac{X_{\min}(y_n)}{y_n} \rightarrow -\infty, \quad \frac{X_1(y_n) - X_{\min}(y_n)}{y_n} \rightarrow +\infty.$$

To this aim, a deeper understanding of the structure of $S_{\bar{y}}$ is in order, when \bar{y} is large. Even though this analysis can be performed for every \bar{y} , to avoid technicalities we prefer to consider only those values for which the line $y = \bar{y}$ has a finite number of transverse intersections with Γ (recall that, by Sard's Lemma, such condition holds for a.e. \bar{y}).

Lemma 3.13. Let $t \mapsto (x(t), y(t))$ denote a regular parameterization of Γ , and let $\bar{y} > 0$ be a regular value for $t \mapsto y(t)$. Then:

1. there exist $t_1 < t_2 < \dots < t_k$, k odd, such that $y(t) = \bar{y}$ if and only if $t = t_i$ for some i ;
2. $(-1)^i y'(t_i) < 0$;
3. writing $S_{\bar{y}} = \bigcup_{j=1}^k (s_{2j-1}, s_{2j}) \times \{\bar{y}\}$, the intervals being disjoint and ordered, it holds
$$\{s_{2j-1}\}_j = \{x(t_{2i-1}), x(t_{2i}) + 2\pi\}_i, \quad \{s_{2j}\}_j = \{x(t_{2i}), x(t_{2i+1}) + 2\pi\}_i;$$
4. $|x(t_{i+1}) - x(t_i)| < 2\pi$ for every $i = 1, \dots, k-1$;
5. for every j , (s_{2j}, \bar{y}) and (s_{2j+1}, \bar{y}) both belong either to Γ or to $\Gamma + (2\pi, 0)$.

Proof. Since \bar{y} is regular, we immediately deduce that $\Gamma \cap \{y = \bar{y}\}$ is finite; thus properties 1 and 2 are trivial consequences of the fact that $y(t) < \bar{y}$ for $t < t_1$ and $y(t) > \bar{y}$ for $t > t_k$.

Observe that $t \mapsto (x(t) + 2\pi, y(t))$ parameterize $\Gamma + (2\pi, 0)$: we deduce that the inward normal to ∂S is given by $(y'(t), -x'(t))$ for points of Γ and by $(-y'(t), x'(t))$ for points of $\Gamma + (2\pi, 0)$. By 2, we deduce that if i is odd (resp. even), then $S_{\bar{y}}$ lies on the right (resp. on the left) of $x(t_i)$ and on the left (resp. on the right) of $x(t_i) + 2\pi$, and also 3 follows.

Let us assume by contradiction that property 4 is false and, say,

$$x(t_i) < x(t_i) + 2\pi < x(t_{i+1}) < x(t_{i+1}) + 2\pi, \quad y(t_i) = y(t_{i+1}) = \bar{y}, \quad y(t) \neq \bar{y} \text{ in } [a, b].$$

Let A be the bounded region of \mathbb{R}^2 , surrounded by the Jordan curve joining $\Gamma|_{(t_i, t_{i+1})}$ and the segment $[x(t_i), x(t_{i+1})] \times \{\bar{y}\}$. We deduce that the connected curve $(\Gamma + (2\pi, 0))|_{(t_i, t_{i+1})}$ belongs to A for $t \rightarrow t_i^+$, and to $\mathbb{R}^2 \setminus A$ for $t \rightarrow t_{i+1}^+$, so that for some $t^* \in (t_i, t_{i+1})$ it intersects ∂A , i.e. Γ , a contradiction.

In a similar fashion, let 5 be false and, for concreteness, let i', i'' be such that $s_{2j} = x(t_{2i'})$ and $s_{2j+1} = x(t_{2i''}) + 2\pi$. Let us consider the Jordan curve γ obtained by joining $\Gamma|_{(0, t_{2i'})}$, $(\Gamma + (2\pi, 0))|_{(0, t_{2i''})}$ and the two segments $[s_{2j}, s_{2j+1}] \times \{\bar{y}\}$, $[0, 2\pi] \times \{0\}$. Calling A the bounded region delimited by γ we have that $\Gamma|_{(t_{2i'}, +\infty)}$ belongs to A for $t \rightarrow t_{2i'}^+$, while it belongs to its complement for $t \rightarrow +\infty$. Thus $\Gamma|_{(t_{2i'}, +\infty)}$ must intersect γ , a contradiction since $(s_{2j}, s_{2j+1}) \times \{\bar{y}\} \cap S_{\bar{y}} = \emptyset$. \square

Remark 3.14. Property 4 and 5 in the above lemma imply that, if $I \subset [X_{\min}(\bar{y}), X_{\max}(\bar{y})]$ is a closed interval, with $|I| \geq 2\pi$, then it contains an interval $(s_{2j-1}, s_{2j}) \subset S_{\bar{y}}$ which has one endpoint on Γ and the other on $\Gamma + (2\pi, 0)$: indeed, by 4, I contains at least one point of Γ and one point of $\Gamma + (2\pi, 0)$, and the claim follows by 5. Moreover, if $x(t_{i_1})$ and $x(t_{i_2})$ belong to different connected components of $\mathbb{R} \setminus I$, then either s_{2j-1} or s_{2j} belong to $\Gamma|_{(t_{i_1}, t_{i_2})}$.

Definition 3.15. If $S_{\bar{y}}$ is not connected we call its connected components *bottlenecks*. A bottleneck

$$\Sigma = (s', s'') \times \{\bar{y}\}$$

disconnects S in two (open, connected) components, only one of which is adjacent to $\{y = 0\}$ (where the non-homogeneous Dirichlet datum is assigned for v):

$$S \setminus \Sigma = \mathcal{N}_{\Sigma} \cup \mathcal{Z}_{\Sigma}, \quad \text{with} \quad \begin{cases} \mathcal{N}_{\Sigma} \cap \mathcal{Z}_{\Sigma} = \emptyset \\ \partial \mathcal{N}_{\Sigma} \supset [0, 2\pi] \times \{0\} \\ v = 0 \text{ on } \partial \mathcal{Z}_{\Sigma} \setminus \Sigma. \end{cases}$$

In the following we will only deal with the component \mathcal{Z}_{Σ} (\mathcal{Z} stays for ‘‘Zero Dirichlet data’’), which may be either unbounded (when the endpoints of Σ belong to different components of ∂S) or not (when they both belong either to Γ or to $\Gamma + (2\pi, 0)$).

Finally, we denote with $B_r(\Sigma)$ the disk of radius r centered at the middle point of Σ , and with

$$B'_r(\Sigma) := \text{the connected component of } S \cap B_r(\Sigma) \text{ which contains } \Sigma.$$

In fact, any small bottleneck provides a strong decay of the gradient of v , as shown in the following lemma.

Lemma 3.16. *Let $\pi < R_1 < R_2$. Under the notation of Definition 3.15 it holds*

$$\int_{\mathcal{Z} \setminus B'_{R_1}(\Sigma)} |\nabla v|^2 dx dy \leq \frac{C(R_1, R_2)}{\log(2R_1/|\Sigma|)} \int_{B'_{R_2}(\Sigma)} |\nabla v|^2 dx dy.$$

Proof. Let $2\ell = |\Sigma|$ and

$$\eta(x) = \begin{cases} \frac{\log|x| - \log\ell}{\log R_1 - \log\ell} & x \in \mathcal{Z}(\Sigma) \cap B'_{R_1}(\Sigma), \ell \leq |x| \leq R_1 \\ 1 & x \in \mathcal{Z}(\Sigma) \setminus B'_{R_1}(\Sigma) \\ 0 & \text{elsewhere.} \end{cases}$$

Note that $\eta^2 v$ is in $H_0^1(\mathcal{Z}(\Sigma))$ and $\nabla \eta \neq 0$ only in $B'_{R_1}(\Sigma)$. Since v is harmonic and $v^2|_{B'_{R_2}(\Sigma)}$ is subharmonic on $B_{R_2}(\Sigma)$ (when extended to zero outside $B'_{R_2}(\Sigma)$), we obtain

$$\begin{aligned} \int_{\mathcal{Z} \setminus B'_{R_1}(\Sigma)} |\nabla v|^2 dx dy &\leq \int_{\mathcal{Z}} |\nabla(\eta v)|^2 dx dy = \int_{\mathcal{Z}} \nabla v \cdot \nabla(\eta^2 v) dx dy + \int_{\mathcal{Z}} |\nabla \eta|^2 v^2 dx dy \\ &\leq 2\pi \int_{\ell}^1 \frac{r dr}{r^2 (\log R_1 - \log \ell)^2} \cdot \left(\max_{\mathcal{Z} \cap B'_{R_1}(\Sigma)} v^2 \right) \\ &\leq \frac{1}{\log R_1 - \log \ell} \cdot \frac{2}{(R_2 - R_1)^2} \max_{x \in \mathcal{Z} \cap B'_{R_1}(\Sigma)} \int_{B_{R_2 - R_1}(x)} (v|_{B'_{R_2}(\Sigma)})^2 dx dy \\ &\leq \frac{1}{\log(2R_1/|\Sigma|)} \cdot \frac{2}{(R_2 - R_1)^2} \int_{B'_{R_2}(\Sigma)} v^2 dx dy. \end{aligned}$$

To conclude, we recall that there exists $C_P = C_P(R_1, R_2)$ such that the Poincaré inequality

$$\int_{B_{R_2}} u^2 dx dy \leq C_P \int_{B_{R_2}} |\nabla u|^2 dx dy$$

holds, for every $u \in H^1(B_{R_2})$ such that $u|_{\gamma} = 0$, for some connected curve γ having endpoints on ∂B_{R_1} and ∂B_{R_2} , respectively (indeed, this implies a Poincaré inequality in $B_{R_2} \setminus B_{R_1}$, with constant $4/R_2^2$ independent of γ ; then the inequality on B_{R_2} follows using the trace inequality both on B_{R_1} and $B_{R_2} \setminus B_{R_1}$). \square

The above estimate can be used to deal with groups of bottlenecks belonging to localized intervals of $S_{\bar{y}}$.

Lemma 3.17. *Let \bar{y} be a regular value of $t \mapsto y(t)$ and $X_1(\bar{y})$ be defined as in Lemma 3.11. For $h \in \mathbb{N}$ we define*

$$\sigma_h := X_1(\bar{y}) - 3(2h + 1)\pi,$$

$$I_h := (\sigma_h - 3\pi, \sigma_h + 3\pi) \times \{\bar{y}\},$$

$$\delta_h := \max\{|\Sigma| : \Sigma \text{ is a bottleneck in } S_{\bar{y}}, \Sigma \subset I_h\},$$

$$B'_h := \left\{ \bigcup_i A_i : A_i \text{ is a connected component of } S \cap B_{3\pi}(\sigma_h, \bar{y}) \text{ which intersects } S_{\bar{y}} \cap I_h \right\}.$$

Then there exists a constant $C > 0$ such that

$$\int_{B'_{h+1}} |\nabla v|^2 dx dy \leq \frac{C}{\log(3\pi/\delta_h)} \int_{B'_h} |\nabla v|^2 dx dy.$$

Proof. We recall that, as shown in the proof of Lemma 3.2, $\overline{S_{\bar{y}}} \cap \{X_1 + 2j\pi : j \in \mathbb{Z}\} = \{X_1\}$ (for easier notation we drop the dependence on \bar{y}). As a consequence, each connected component of $S_{\bar{y}}$ is compactly contained in $(X_1 + 2j\pi, X_1 + 2(j+1)\pi) \times \{\bar{y}\}$, for some $j \in \mathbb{Z}$.

If $X_1 - 6h\pi < X_{\min}$ then there is nothing to prove, thus we can assume $h < (X_1 - X_{\min})/(6\pi)$. Recalling the definition of X_1 , and using Remark 3.14, we have that every bottleneck $\Sigma \in I_{h+1}$ belongs to $\mathcal{Z}_{\Sigma'}$, for some $\Sigma' \in (\sigma_h - \pi, \sigma_h + \pi) \times \{\bar{y}\}$. More formally, writing

$$S_{\bar{y}} \cap \{\sigma_h - \pi < x < \sigma_h + \pi\} = \bigcup_{i=1}^n \Sigma_i,$$

we obtain that

$$B'_{h+1} \subset B_{3\pi}(\sigma_h, \bar{y}) \cap \bigcup_{i=1}^n \mathcal{Z}_{\Sigma_i}.$$

In order to apply Lemma 3.16 we choose $R_1 = 3\pi/2$, $R_2 = 2\pi$; we obtain that, for every i ,

$$B'_{R_2}(\Sigma_i) \subset B_{3\pi}(\sigma_h, \bar{y}), \quad B'_{R_1}(\Sigma_i) \cap B_{3\pi}(\sigma_{h+1}, \bar{y}) = \emptyset,$$

and thus

$$\begin{aligned} \int_{B'_{h+1}} |\nabla v|^2 dx dy &\leq \sum_{i=1}^n \int_{\mathcal{Z}_{\Sigma_i} \setminus B'_{R_1}(\Sigma_i)} |\nabla v|^2 dx dy \\ &\leq \sum_{i=1}^n \frac{C}{\log(3\pi/|\Sigma_i|)} \int_{B'_{R_2}(\Sigma_i)} |\nabla v|^2 dx dy \leq \frac{C}{\log(3\pi/\delta_h)} \int_{B'_h} |\nabla v|^2 dx dy, \end{aligned}$$

for a suitable constant $C > 0$. □

Such decay estimate can be easily iterated.

Lemma 3.18. *There exists $\gamma > 0$ such that, for $1 \leq k < (X_1 - X_{\min})/(6\pi)$,*

$$\int_{S_{\bar{y}} \cap I_k} v^2 dx \leq \left(\frac{\gamma}{\log(3k/2)} \right)^k \int_{S \cap \{x < X_1(\bar{y})\}} |\nabla v|^2 dx dy.$$

Proof. Iterating Lemma 3.17 we obtain

$$\int_{B'_{k+1}} |\nabla v|^2 dx dy \leq \left(\prod_{h=1}^k \frac{C}{\log(3\pi/\delta_h)} \right) \int_{B'_0} |\nabla v|^2 dx dy.$$

Now, using a Poincaré inequality analogous to that mentioned at the end of the proof of Lemma 3.16, together with the trace inequality in $B'_{k+1} \cap \{y > \bar{y}\}$, we obtain the existence of a constant C_T such that

$$\int_{S_{\bar{y}} \cap I_k} v^2 dx \leq C_T \int_{B'_{k+1}} |\nabla v|^2 dx dy.$$

Since $B'_0 \subset \{x < X_1(\bar{y})\}$, this implies

$$\int_{S_{\bar{y}} \cap I_k} v^2 dx \leq \left(\prod_{h=1}^k \frac{\gamma}{\log(3\pi/\delta_h)} \right) \int_{S \cap \{x < X_1(\bar{y})\}} |\nabla v|^2 dx dy,$$

where $\gamma = \max\{C_T, 1\} \cdot C$. But then the lemma follows by the elementary estimate

$$\max \left\{ \prod_{h=1}^k \frac{1}{\log(3\pi/\delta_h)} : \sum_{h=1}^k \delta_h \leq 2\pi, \delta_h > 0 \text{ for every } h \right\} \leq \left(\frac{1}{\log(3k/2)} \right)^k. \quad \square$$

At this point, we have all the ingredients to conclude.

End of the proof of Proposition 3.4. As we already mentioned, after Lemmas 3.8, 3.10 and 3.12, we can assume that, along the sequence $y_n \rightarrow +\infty$ of regular values of $t \mapsto y(t)$, it holds

$$\frac{X_{\min}(y_n)}{y_n} \rightarrow -\infty, \quad \frac{X_1(y_n) - X_{\min}(y_n)}{y_n} \rightarrow +\infty.$$

Let $k_n \in \mathbb{N}$, $k_n \rightarrow +\infty$, be defined by

$$\frac{X_1(y_n) - X_{\min}(y_n)}{6\pi} - 1 < k_n < \frac{X_1(y_n) - X_{\min}(y_n)}{6\pi}.$$

We have ($X_{1,n} := X_1(y_n)$)

$$\begin{aligned} \int_0^{2\pi} w^2(x, y_n) dx &= \int_{S_{y_n} \cap \{x > X_{1,n}\}} e^{-2\alpha x} v^2 dx + \sum_{h=1}^{k_n} \int_{S_{y_n} \cap I_h} e^{-2\alpha x} v^2 dx \\ &\leq \underbrace{e^{-2\alpha X_{1,n}} \int_{S_{y_n} \cap \{x > X_{1,n}\}} v^2 dx}_{(A)} + \underbrace{\sum_{h=1}^{k_n} e^{-2\alpha(X_{1,n} - 6(h+1)\pi)} \int_{S_{y_n} \cap I_h} v^2 dx}_{(B)}. \end{aligned}$$

The first term can be easily estimated, as usual, using Remark 3.3:

$$(A) \leq e^{-2\alpha X_{1,n}} \int_{S_{y_n}} v^2 dx \leq 10e^{-2\alpha X_{1,n}} \int_{S \cap \{y > y_n\}} |\nabla v|^2 dx dy.$$

On the other hand, we can bound the second term using Lemma 3.18:

$$\begin{aligned} (B) &\leq \sum_{h=1}^{k_n} e^{-2\alpha(X_{1,n} - 6(h+1)\pi)} \left(\frac{\gamma}{\log(3h/2)} \right)^h \int_{S \cap \{x < X_{1,n}\}} |\nabla v|^2 dx dy \\ &\leq e^{-2\alpha(X_{1,n} - 6\pi)} \int_{S \cap \{x < X_{1,n}\}} |\nabla v|^2 dx dy \cdot \sum_{h=1}^{k_n} \exp(12h\pi + h(\log \gamma - \log \log(3h/2))) \\ &\leq e^{-2\alpha(X_{1,n} - 6\pi)} \int_{S \cap \{x < X_{1,n}\}} |\nabla v|^2 dx dy \cdot \underbrace{\sum_{h=1}^{\infty} \exp(12\pi + \log \gamma - \log \log(3h/2)) h}_{\leq C} \\ &\leq C e^{-2\alpha X_{1,n}} \int_{S \cap \{x < X_{1,n}\}} |\nabla v|^2 dx dy. \end{aligned}$$

Now, there are two possibilities: either (up to subsequences) $X_{1,n} \geq 0$, in which case

$$\int_0^{2\pi} w^2(x, y_n) dx \leq (A) + (B) \leq \int_S |\nabla v|^2 dx dy < +\infty,$$

and we can conclude using Lemma 3.9; otherwise, if $X_{1,n} < 0$ we can use Lemma 3.11, obtaining

$$\begin{aligned} \int_0^{2\pi} w^2(x, y_n) dx &\leq (A) + (B) \leq C e^{-2\alpha X_{1,n}} \int_{S \setminus ([X_1(\bar{y}), 2\pi] \times [0, \bar{y}])} |\nabla v|^2 dx dy \\ &\leq C \exp\left(-2\alpha X_{1,n} - \frac{X_{1,n}^2}{y_n}\right) \leq C \exp(\alpha^2 y_n), \end{aligned}$$

and again, by Lemma 3.9, the proposition follows. \square

4 High multiplicity points are isolated

In this section we go back to the general case of system (18), aiming at excluding that \mathcal{W} contains more than one point (one can stick to the case of an even number of nodal regions, even though all the arguments hold also in the odd case, taking into account Remark 2.18). This will allow to complete the proof of Theorem 1.5

Proposition 4.1. *Under the reduction of Section 2, we have that $\mathcal{W} = \{0\}$.*

By contradiction, throughout the section we assume that the above statement is false. Nonetheless, recall that by construction we can bound the set \mathcal{W} into a strip of width π .

Lemma 4.2. *Assume that $\mathcal{W} \supsetneq \{0\}$. Then $\mathcal{T}^{-1}(\mathcal{W} \setminus \{0\})$ is the union of infinitely many connected, simply connected, components. Moreover there exists $a > 0$ such that, up to a translation, we can assume that any of such components is contained in $[-\pi/2 + 2m\pi, \pi/2 + 2m\pi] \times [a, +\infty)$, for some $m \in \mathbb{Z}$.*

Proof. The lemma follows by the definition of the map \mathcal{T} (see equation (15)), together with Proposition 2.1, and in particular property (7) of the model case scenario (MCS). \square

Definition 4.3. We denote the unique lifting of Γ_{ij} (i.e. the connected component of $\mathcal{T}^{-1}(\Gamma_{ij})$), with an endpoint in $[2m\pi, 2(m+1)\pi) \times \{0\}$, $m \in \mathbb{Z}$, as

$$\tilde{\Gamma}_{ij}^m,$$

as long as it is non-empty (i.e., if i and j are consecutive). Analogously, for any $m \in \mathbb{Z}$, we introduce the set

$$S^m = S + (2m\pi, 0)$$

(recall that the strip S has been defined in equation (16)).

To prove the proposition we have to distinguish two cases, according to the horizontal behaviour of the smooth curves $\tilde{\Gamma}_{ij}^m$.

Lemma 4.4. *Consider the families of infima, depending on i, j, m ,*

$$\inf\{x : (x, y) \in \tilde{\Gamma}_{ij}^m\}, \quad \inf\{x : (x, y) \in S^m\}.$$

Then, if one of them is finite, each of them is. An analogous statement holds for the suprema $\sup\{x : (x, y) \in \tilde{\Gamma}_{ij}^m\}, \sup\{x : (x, y) \in S^m\}$.

Proof. The lemma easily follows from the trivial property

$$\begin{aligned} \inf\{x : (x, y) \in \tilde{\Gamma}_{k_1}^m\} &= \inf\{x : (x, y) \in S^m\} \leq \inf\{x : (x, y) \in \tilde{\Gamma}_{ij}^m\} \\ &\leq \inf\{x : (x, y) \in \tilde{\Gamma}_{k_1}^{m+1}\} = \inf\{x : (x, y) \in \tilde{\Gamma}_{k_1}^m\} + 2\pi. \end{aligned} \quad \square$$

Case 1) Both $\inf\{x : (x, y) \in \tilde{\Gamma}_{ij}^m\}$ and $\sup\{x : (x, y) \in \tilde{\Gamma}_{ij}^m\}$ are finite, for every i, j, m .

Let $Q = (-\pi, \pi) \times \mathbb{R}_+$ (notice that $\partial Q \cap \mathcal{W} = \emptyset$, by Lemma 4.2). We can cover the set $Q \setminus \mathcal{Z}$ with a finite number of copies of S , that is, there exists $M \in \mathbb{N}$ such that

$$Q \setminus \mathcal{T}^{-1}(\mathcal{Z} \setminus \{0\}) \subset \bigcup_{m=-M}^M S^m.$$

On Q we introduce the $k(2M + 1)$ functions $v_i^m \in Lip(Q)$, $i = 1, \dots, k$, $m = -M, \dots, M$ defined as

$$v_i^m(x, y) = \begin{cases} |v(x, y)| & \text{if } (x, y) \in S^m, \mathcal{T}(x, y) \in \omega_i \\ 0 & \text{otherwise.} \end{cases}$$

Using the notation of [23], we want to show that the vector of such densities belongs to the class $\mathcal{G}(Q)$ defined in that paper. More precisely, in the present context this is equivalent to the following lemma.

Lemma 4.5. *Let v_i^m be defined as above. Then*

- each v_i^m is a non-negative, locally Lipschitz, subharmonic function and there exists $\mu_i^m \in \mathcal{M}(Q)$ non-negative Radon measure, supported on $Q \cap \partial\{v_i^m > 0\}$ such that

$$-\Delta v_i^m = -\mu_i^m \quad \text{in the sense of distributions } \mathcal{D}'(Q);$$

- for every $\mathbf{x}_0 \in Q$ and $0 < r < \text{dist}(\mathbf{x}_0, \partial Q)$ the following identity holds

$$\frac{d}{dr} \int_{B_r(\mathbf{x}_0)} \sum_{i,m} |\nabla v_i^m|^2 = 2 \int_{\partial B_r(\mathbf{x}_0)} \sum_{i,m} (\partial_\nu v_i^m)^2 d\sigma.$$

Proof. The first point follows directly from the definition of the involved functions, while for the second one we adopt the same strategy of [1, Theorem 15] and [23, Theorem 8.4]. We consider $\delta > 0$ as a small quantity and we integrate the equation in v_i^m against $(\mathbf{x} - \mathbf{x}_0) \cdot \nabla v_i^m$ on the set $B_r(\mathbf{x}_0) \cap \{v_i^m > \delta\}$. Some integrations by parts (the same one exploits in order to prove the Pohozaev identity) yield

$$\int_{\partial B_r(\mathbf{x}_0) \cap \{v_i^m > \delta\}} |\nabla v_i^m|^2 = 2 \int_{\partial B_r(\mathbf{x}_0) \cap \{v_i^m > \delta\}} \sum_{i,m} (\partial_\nu v_i^m)^2 d\sigma + \frac{1}{r} \int_{B_r(\mathbf{x}_0) \cap \partial\{v_i^m > \delta\}} |\nabla v_i^m|^2 (\mathbf{x} - \mathbf{x}_0) \cdot \nu d\sigma$$

and thus

$$\frac{d}{dr} \int_{B_r(\mathbf{x}_0)} \sum_{i,m} |\nabla v_i^m|^2 = 2 \int_{\partial B_r(\mathbf{x}_0)} \sum_{i,m} (\partial_\nu v_i^m)^2 d\sigma + \frac{1}{r} \lim_{\delta \rightarrow 0^+} \sum_{i,m} \int_{B_r(\mathbf{x}_0) \cap \partial\{v_i^m > \delta\}} |\nabla v_i^m|^2 (\mathbf{x} - \mathbf{x}_0) \cdot \nu d\sigma.$$

To evaluate the last limit, let $\varepsilon > 0$ and let us consider the set $S_\varepsilon = \{x \in Q : \sum_{i,m} |\nabla v_i^m| \leq \varepsilon\}$. Thanks to Lemma 1.4 we have that

$$Q \cap \mathcal{T}^{-1}(\mathcal{W} \setminus \{0\}) \subset Q \cap S_\varepsilon \quad \text{for all } \varepsilon > 0.$$

We also observe that, since each v_i^m is harmonic when positive,

$$\int_{B_r(\mathbf{x}_0) \cap \partial\{v_i^m > \delta\}} |\nabla v_i^m|^2 d\sigma = - \int_{B_r(\mathbf{x}_0) \cap \partial\{v_i^m > \delta\}} \partial_\nu v_i^m = \int_{\partial B_r(\mathbf{x}_0) \cap \{v_i^m > \delta\}} \partial_\nu v_i^m \leq C$$

for a constant C independent of δ .

We split the remainder into two parts. Outside of S_ε we can exploit the definition of v_i^m and the regularity of the zero set (which is given by the union of the curves $\tilde{\Gamma}_{ij}^m$), and find

$$\begin{aligned} \lim_{\delta \rightarrow 0^+} \sum_{i,m} \int_{B_r(\mathbf{x}_0) \cap \partial\{v_i^m > \delta\} \setminus S_\varepsilon} |\nabla v_i^m|^2 (\mathbf{x} - \mathbf{x}_0) \cdot \nu d\sigma &= \sum_{i,m} \int_{B_r(\mathbf{x}_0) \cap \partial\{v_i^m > 0\} \setminus S_\varepsilon} |\nabla v_i^m|^2 (\mathbf{x} - \mathbf{x}_0) \cdot \nu d\sigma \\ &= \sum_{i,j,m} \int_{B_r(\mathbf{x}_0) \cap \tilde{\Gamma}_{ij}^m \setminus S_\varepsilon} |\nabla v|^2 (\mathbf{x} - \mathbf{x}_0) \cdot J ds - \sum_{i,j,m} \int_{B_r(\mathbf{x}_0) \cap \tilde{\Gamma}_{ij}^m \setminus S_\varepsilon} |\nabla v|^2 (\mathbf{x} - \mathbf{x}_0) \cdot J ds = 0. \end{aligned}$$

Here J is the symplectic matrix and the two opposite contributions follows by the fact that each $\tilde{\Gamma}_{ij}^m$ appears as the boundary of the positivity set of two functions v_i^m . On the other hand, inside of S_ε , there exists $C > 0$ independent of δ such that

$$\int_{B_r(\mathbf{x}_0) \cap \partial\{v_i^m > \delta\} \cap S_\varepsilon} |\nabla v_i^m|^2 (\mathbf{x} - \mathbf{x}_0) \cdot \nu d\sigma \leq C\varepsilon \int_{B_r(\mathbf{x}_0) \cap \partial\{v_i^m > \delta\} \cap S_\varepsilon} |\nabla v_i^m|^2 d\sigma \leq C\varepsilon.$$

From the arbitrariness of ε we conclude

$$\lim_{\delta \rightarrow 0^+} \sum_{i,m} \int_{B_r(\mathbf{x}_0) \cap \partial\{v_i^m > \delta\}} |\nabla v_i^m|^2 (\mathbf{x} - \mathbf{x}_0) \cdot \nu d\sigma = 0. \quad \square$$

Proof of Proposition 4.1 (Case 1). By Lemma 4.5, we are in a position to apply the results of [1, Theorem 16], [2, Theorem 4.7] and [23, Theorem 1.1]. We find in particular that the set $\mathcal{T}^{-1}(\mathcal{W} \setminus \{0\}) \cap Q$ is made of (at most countable many) isolated points. Since \mathcal{W} is a connected set, it must be that $\mathcal{T}^{-1}(\mathcal{W} \setminus \{0\}) \cap Q$ contains at most a point $\tilde{\mathbf{x}}_0 \in Q$. But then, going back to the original coordinates, we find that

$$\mathcal{W} = \{0\} \cup \{\mathcal{T}(\tilde{\mathbf{x}}_0)\}.$$

Again by the connectedness of \mathcal{W} , we conclude that $\mathcal{W} = \{0\}$. □

Case 2) Either $\inf\{x : (x, y) \in \tilde{\Gamma}_{ij}^m\}$ or $\sup\{x : (x, y) \in \tilde{\Gamma}_{ij}^m\}$ are not finite, for every i, j, m .

Also in this case we write $Q = (-\pi, \pi) \times (0, +\infty)$ (we recall again that $\partial Q \cap \mathcal{W} = \emptyset$, by Lemma 4.2). Possibly up to a further translation and using Sard's Lemma, we can assume that any intersection between each $\tilde{\Gamma}_{ij}^m$ and ∂Q is transverse. Furthermore, we denote by D_i , $i \in \mathbb{N}$, the (open) connected components of $Q \setminus \{v = 0\}$.

Lemma 4.6. *If v solves (18) then $v \in H^1(Q)$.*

Proof. Notice that we already know that $v \in H^1(S^m)$, for every $m \in \mathbb{Z}$, and $v \in H^1(D_i)$, for every $i \in \mathbb{N}$. More precisely, there exists a sequence $\{m_i\}_{i \in \mathbb{N}} \subset \mathbb{Z}$ such that

$$D_i \subset S^{m_i} \iff m = m_i$$

and, consequently, we have

$$v(x, y)|_{D_i} = v(x + 2m_i\pi, y)|_{D_i - (2m_i\pi, 0)} = \lambda^{m_i} v(x, y)|_{D_i - (2m_i\pi, 0)},$$

where now each $D_i - (2m_i\pi, 0)$ is a subset of S^0 and, moreover, for each $i \neq j$, $D_i - (2m_i\pi, 0) \cap D_j - (2m_j\pi, 0) = \emptyset$. Consequently we have, for each $i \in \mathbb{N}$,

$$\int_{D_i} |\nabla v|^2 = \lambda^{2m_i} \int_{D_i - (2m_i\pi, 0)} |\nabla v|^2.$$

A priori, the function $i \mapsto m_i$ is not one-to-one: indeed, there may exist different connected components that belong to the same translated strip. With an abuse of notation, in the rest of the proof we shall assume that D_i stands for the union of connected components of the same translation of the strip, so that the function $i \mapsto m_i$ is one-to-one. With this notation, for any $n \in \mathbb{N}$ we define the sequence of functions

$$v_n = \sum_{i=0}^n v|_{D_i} = \sum_{i=0}^n \lambda^{m_i} v|_{D_i - (2m_i\pi, 0)}.$$

By definition, clearly we have that $v_n \in H^1(Q)$. We want to show that the limit of the sequence, that is the function v itself, is a $H^1(Q)$ function. To this end, it is sufficient to show that the series of the norm

$$\int_Q |\nabla v|^2 + v^2 = \lim_{n \rightarrow +\infty} \int_Q |\nabla v_n|^2 + v_n^2 = \sum_{i=0}^{+\infty} \int_{D_i} |\nabla v|^2 + v^2$$

converges. Here we shall only address the contribution regarding the L^2 -norm of gradient, as the L^2 -norm of the functions themselves can be estimated by means of the Poincaré inequality (see Remark 3.3, or Lemma 3.16).

Let $I \subset \mathbb{N}$ be the subset of indices such that if $i \in I$, then $m_i > 1$; since the general term in the series is positive, we can proceed as follows

$$\sum_{i=0}^{+\infty} \int_{D_i} |\nabla v|^2 = \sum_{i \in I} \int_{D_i} |\nabla v|^2 + \sum_{i \notin I} \int_{D_i} |\nabla v|^2.$$

In the following we deal with the case $\lambda \geq 1$, the opposite case following with minor changes. Consequently, we can estimate the second term as follows

$$\sum_{i \notin I} \lambda^{2m_i} \int_{D_i - (2m_i\pi, 0)} |\nabla v|^2 \leq \lambda^2 \sum_{i \notin I} \int_{D_i - (2m_i\pi, 0)} |\nabla v|^2 \leq \lambda^2 \int_S |\nabla v|^2 < +\infty,$$

in such a way that we are left to consider only the terms with $i \in I$. We now have two alternatives. If the set I is finite then $v \in H^1(Q)$ since $v \in H^1(S^0)$. Otherwise, let $i \in I$, so that $m_i > 0$. By definition, it means that the set $D_i - (2m_i\pi, 0)$ belongs to $Q - (2m_i\pi, 0)$. Up to a further translation, we can assume that $\{x = 0\} \cap \mathcal{T}^{-1}(\mathcal{W} \setminus \{0\}) \neq \emptyset$ and we denote

- $y_0 := \min\{y : (0, y) \in \mathcal{T}^{-1}(\mathcal{W} \setminus \{0\})\}$,
- $\sigma^m := \{(2m\pi, y) : 0 \leq y \leq y_0\}$ for $m \in \mathbb{Z}$.

In this setting we have that, while $\mathbb{R}_+^2 \setminus \mathcal{T}^{-1}(\mathcal{W} \setminus \{0\})$ is connected,

$$\mathbb{R}_+^2 \setminus (\sigma^m \cup \mathcal{T}^{-1}(\mathcal{W} \setminus \{0\}))$$

is disconnected, for every m . We deduce that, for any $0 \leq m < m_i$, the segment σ^m disconnects S^0 in at least two connected components, one containing $S^0 \cap Q$, the other $D_i - (2m_i\pi, 0)$. We are then in a similar position to that of Definition 3.15, with the only difference that now we have to consider

vertical *bottlenecks*. With this language, S^0 goes through at least $m_i - 1$ bottlenecks before reaching $D_i - (2m_i\pi, 0)$. Using the same proof of Lemma 3.18 we have that, for m_i large enough,

$$\int_{D_i - (2m_i\pi, 0)} |\nabla v|^2 \leq \left(\frac{C}{\log C(m_i - 1)} \right)^{m_i - 1} \int_S |\nabla v|^2$$

where the constant $C > 1$ is universal. Consequently we have

$$\begin{aligned} \sum_{i \in I} \int_{D_i} |\nabla v|^2 &\leq \sum_{i \in I} \lambda^{2m_i} \int_{D_i - (2m_i\pi, 0)} |\nabla v|^2 \leq \sum_{i \in I} \lambda^{2m_i} \left(\frac{C}{\log C(m_i - 1)} \right)^{m_i - 1} \int_S |\nabla v|^2 \\ &\leq \lambda \sum_m \left(\frac{C\lambda^2}{\log Cm} \right)^m \int_S |\nabla v|^2 < +\infty. \end{aligned} \quad \square$$

To conclude, we want to show that v is harmonic across the singular set. The key step in this direction is that v can be integrated by parts on each of its nodal connected components.

Lemma 4.7. *For every i ,*

$$\partial D_i \cap \mathcal{T}^{-1}(\mathcal{W} \setminus \{0\}) = \emptyset.$$

Proof. Since we are dealing with Case 2, by Lemma 4.4 we know that $\bar{Q} \cap \mathcal{T}^{-1}(\mathcal{Z}_2)$ consists in the countable union of smooth paths of finite lengths, pairwise disjoint. Moreover, infinitely many of them have one endpoint on the line $\{x = -\pi\}$ and the other on $\{x = \pi\}$. Let us denote with γ_j , $j \in \mathbb{N}$ such paths. We recall that \mathcal{W} is connected and $0 \in \mathcal{W}$, so that any connected component of $\mathcal{T}^{-1}(\mathcal{W} \setminus \{0\})$ is unbounded. We deduce that, for every j , there exist two open connected sets Q_j^\pm satisfying

$$Q \setminus \gamma_i = Q_j^- \cup Q_j^+, \quad Q_j^- \cap Q_j^+ = \emptyset, \quad \bar{Q}_j^- \cap \mathcal{T}^{-1}(\mathcal{W} \setminus \{0\}) = \emptyset.$$

With this notation, we are left to prove that any D_i is contained in some Q_j^- . Notice that, up to a relabelling, one can assume that $Q_j^- \subset Q_{j+1}^-$, $Q_j^+ \supset Q_{j+1}^+$. Let us consider the open set $Q^- = \bigcup_j Q_j^-$. Since $\mathcal{W} \setminus \{0\}$ is non-empty, we have that $Q^- \neq Q$, and by construction

$$\partial Q^- \setminus \partial Q = \lim_j \gamma_j \subset \mathcal{T}^{-1}(\mathcal{W} \setminus \{0\}).$$

Recalling Lemma 4.2 we infer that

$$Q^+ := \bigcap_j Q_j^+ \subset \left[-\frac{\pi}{2}, \frac{\pi}{2} \right] \times [a, +\infty).$$

Now, take any D_i . Of course, for every j , either $D_i \subset Q_j^-$ or $D_i \subset Q_j^+$; on the other hand, $\partial D_i \cap \partial Q \neq \emptyset$. We deduce that $D_i \not\subset Q^+$, or equivalently that $D_i \subset Q_j^-$ for some j . \square

Lemma 4.8. *The function v is harmonic in Q (i.e., it is harmonic in \mathbb{R}_+^2).*

Proof. Let $\varphi \in C_0^\infty(Q)$, $\varphi \geq 0$, and let us write

$$\text{supp } \varphi \cap \mathcal{T}^{-1}(\mathcal{Z}_2) = \bigcup_{j=0}^{+\infty} \gamma_j.$$

By Lemmas 4.6 and 4.7, we have that

$$\int_Q \nabla v \cdot \nabla \varphi = \sum_{i=0}^{+\infty} \int_{D_i} \nabla v \cdot \nabla \varphi = \sum_{i=0}^{+\infty} \int_{\partial D_i} \varphi \partial_\nu v = \sum_{i=0}^{+\infty} \sum_{j: \gamma_j \subset \partial D_i} \int_{\gamma_j} \varphi \partial_{\nu_i} v, \quad (25)$$

where ν_i denotes the normal direction to γ_j which points outwards with respect to D_i . Of course, for every j there exist exactly two indexes i_1, i_2 such that $\gamma_j = \overline{D}_{i_1} \cap \overline{D}_{i_2} \cap \text{supp } \varphi$, and $\nu_{i_1} = -\nu_{i_2}$; as a consequence, for every j ,

$$\sum_{i: \partial D_i \supset \gamma_j} \int_{\gamma_j} \varphi \partial_{\nu_i} v = \int_{\gamma_j} \varphi \partial_{\nu_{i_1}} v + \int_{\gamma_j} \varphi \partial_{\nu_{i_2}} v = 0. \quad (26)$$

In order to plug this relation into (25) and conclude the proof, we need to show that the right hand side of (25) converges absolutely, so that we are allowed to rearrange its terms. To this aim let us notice that, since each D_i is a nodal region of v , and φ is non-negative (and compactly supported in \mathcal{Q}), we have that $\varphi \partial_{\nu} v$ does not change its sign on ∂D_i . This yields, for every i ,

$$\sum_{j: \gamma_j \subset \partial D_i} \left| \int_{\gamma_j} \varphi \partial_{\nu_i} v \right| = \left| \sum_{j: \gamma_j \subset \partial D_i} \int_{\gamma_j} \varphi \partial_{\nu_i} v \right| = \left| \int_{\partial D_i} \varphi \partial_{\nu} v \right| = \left| \int_{D_i} \nabla v \cdot \nabla \varphi \right| \leq \int_{D_i} |\nabla v| |\nabla \varphi|,$$

and finally

$$\sum_{i, j: \gamma_j \subset \partial D_i} \left| \int_{\gamma_j} \varphi \partial_{\nu_i} v \right| \leq \|v\|_{H^1(\mathcal{Q})} \|\varphi\|_{H^1(\mathcal{Q})} < +\infty$$

by Lemma 4.6. Therefore, combining (25) and (26) we have, for every $\varphi \in C_0^\infty(\mathcal{Q})$, $\varphi \geq 0$,

$$\int_{\mathcal{Q}} \nabla v \nabla \varphi = 0. \quad \square$$

Proof of Proposition 4.1 (Case 2). Since the zeroes of higher multiplicity of a (non-trivial) harmonic function in the plane are isolated, using Lemma 4.8 we find again that the set $\mathcal{T}^{-1}(\mathcal{W} \setminus \{0\}) \cap \mathcal{Q}$ is made of (at most countable many) isolated points, and we can conclude as in Case 1. \square

Proof of Theorem 1.5. By Proposition 2.1 we have that the singular set \mathcal{W} consists in the finite union of connected components, which are single points by Proposition 4.1, therefore the only thing that is left to prove is the asymptotic expansion (5). Let $m(\mathbf{x}_0) = h \geq 3$, and let \mathcal{U}, v be defined as in (12), (17), respectively. Taking into account Proposition 3.4, together with Remarks 2.17 and 2.19, we obtain

$$\begin{aligned} v(x, y) &= \exp\left(\alpha x - \frac{h}{2}y\right) \cdot \left[a \cos\left(\frac{h}{2}x + \alpha y\right) + b \sin\left(\frac{h}{2}x + \alpha y\right) + o(1) \right] \\ &= C \exp\left(\alpha x + \frac{2\alpha^2}{h}y\right) \cdot \exp\left(-\frac{2\alpha^2}{h}y - \frac{h}{2}y\right) \cdot \left[\cos\left(\frac{h}{2}(x - x_0) + \alpha y\right) + o(1) \right]. \end{aligned}$$

Recalling (15) we infer (up to a rotation) (5), where

$$A(r, \vartheta) := C \exp\left(\alpha(\vartheta + x_0) - \frac{2\alpha^2}{h} \log r\right).$$

Finally, using again Proposition 3.4 we infer that, for some fixed q ,

$$q \leq \vartheta - \frac{2\alpha}{h} \log r \leq q + 2\pi + o(1) \quad \text{as } r \rightarrow 0^+,$$

and also the second condition in (6) follows. \square

Acknowledgments

Work partially supported by the PRIN-2015KB9WPT Grant: “Variational methods, with applications to problems in mathematical physics and geometry”, by the ERC Advanced Grant 2013 n. 339958: “Complex Patterns for Strongly Interacting Dynamical Systems – COMPAT”, by the ERC Advanced Grant 2013 n. 321186 “ReaDi – Reaction-Diffusion Equations, Propagation and Modelling”, and by the INDAM-GNAMPA group.

References

- [1] L. A. Caffarelli, A. L. Karakhanyan, and F.-H. Lin. The geometry of solutions to a segregation problem for nondivergence systems. *J. Fixed Point Theory Appl.*, 5(2):319–351, 2009.
- [2] L. A. Caffarelli and F.-H. Lin. Singularly perturbed elliptic systems and multi-valued harmonic functions with free boundaries. *J. Amer. Math. Soc.*, 21(3):847–862, 2008.
- [3] L. A. Caffarelli and F. H. Lin. Analysis on the junctions of domain walls. *Discrete Contin. Dyn. Syst.*, 28(3):915–929, 2010.
- [4] M. Conti, S. Terracini, and G. Verzini. Nehari’s problem and competing species systems. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 19(6):871–888, 2002.
- [5] M. Conti, S. Terracini, and G. Verzini. An optimal partition problem related to nonlinear eigenvalues. *J. Funct. Anal.*, 198(1):160–196, 2003.
- [6] M. Conti, S. Terracini, and G. Verzini. Asymptotic estimates for the spatial segregation of competitive systems. *Adv. Math.*, 195(2):524–560, 2005.
- [7] M. Conti, S. Terracini, and G. Verzini. On a class of optimal partition problems related to the Fučík spectrum and to the monotonicity formulae. *Calc. Var. Partial Differential Equations*, 22(1):45–72, 2005.
- [8] M. Conti, S. Terracini, and G. Verzini. A variational problem for the spatial segregation of reaction-diffusion systems. *Indiana Univ. Math. J.*, 54(3):779–815, 2005.
- [9] M. Conti, S. Terracini, and G. Verzini. Uniqueness and least energy property for solutions to strongly competing systems. *Interfaces Free Bound.*, 8(4):437–446, 2006.
- [10] E. N. Dancer and Y. H. Du. Positive solutions for a three-species competition system with diffusion. I. General existence results. *Nonlinear Anal.*, 24(3):337–357, 1995.
- [11] E. N. Dancer and Y. H. Du. Positive solutions for a three-species competition system with diffusion. II. The case of equal birth rates. *Nonlinear Anal.*, 24(3):359–373, 1995.
- [12] E. N. Dancer, K. Wang, and Z. Zhang. Dynamics of strongly competing systems with many species. *Trans. Amer. Math. Soc.*, 364(2):961–1005, 2012.
- [13] E. N. Dancer, K. Wang, and Z. Zhang. The limit equation for the Gross-Pitaevskii equations and S. Terracini’s conjecture. *J. Funct. Anal.*, 262(3):1087–1131, 2012.
- [14] E. N. Dancer and Z. Zhang. Dynamics of Lotka-Volterra competition systems with large interaction. *J. Differential Equations*, 182(2):470–489, 2002.

- [15] B. Helffer, T. Hoffmann-Ostenhof, and S. Terracini. Nodal domains and spectral minimal partitions. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 26(1):101–138, 2009.
- [16] D. Hilhorst, S. Martin, and M. Mimura. Singular limit of a competition-diffusion system with large interspecific interaction. *J. Math. Anal. Appl.*, 390(2):488–513, 2012.
- [17] H. Murakawa and H. Ninomiya. Fast reaction limit of a three-component reaction-diffusion system. *J. Math. Anal. Appl.*, 379(1):150–170, 2011.
- [18] J. D. Murray. *Mathematical biology. II*, volume 18 of *Interdisciplinary Applied Mathematics*. Springer-Verlag, New York, third edition, 2003. Spatial models and biomedical applications.
- [19] C. Pommerenke. *Boundary behaviour of conformal maps*, volume 299 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1992.
- [20] B. Sandstede and A. Scheel. Period-doubling of spiral waves and defects. *SIAM J. Appl. Dyn. Syst.*, 6(2):494–547, 2007.
- [21] B. Sandstede, A. Scheel, and C. Wulff. Center-manifold reduction for spiral waves. *C. R. Acad. Sci. Paris Sér. I Math.*, 324(2):153–158, 1997.
- [22] N. Soave and A. Zilio. Uniform bounds for strongly competing systems: the optimal Lipschitz case. *Arch. Ration. Mech. Anal.*, 218(2):647–697, 2015.
- [23] H. Tavares and S. Terracini. Regularity of the nodal set of segregated critical configurations under a weak reflection law. *Calc. Var. Partial Differential Equations*, 45(3-4):273–317, 2012.
- [24] K. Wang and Z. Zhang. Some new results in competing systems with many species. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 27(2):739–761, 2010.

susanna.terracini@unito.it

Dipartimento di Matematica “Giuseppe Peano”, Università di Torino,
Via Carlo Alberto 10, 10123 Torino, Italy

gianmaria.verzini@polimi.it

Dipartimento di Matematica, Politecnico di Milano
piazza Leonardo da Vinci 32, 20133 Milano, Italy

azilio@ehess.fr

Centre d’analyse et de mathématique sociales - UMR 8557 (CNRS / EHESS)
École des hautes études en sciences sociales
54, boulevard Raspail, 75006 Paris, France