

10/6/2009

A DIRICHLET PROBLEM WITH FREE GRADIENT DISCONTINUITY

MICHELE CARRIERO, ANTONIO LEACI & FRANCO TOMARELLI

ABSTRACT. We prove the existence of strong solution for Blake & Zisserman functional under Dirichlet boundary condition. The result is obtained by showing partial regularity of weak solutions up to the boundary through blow-up technique and a decay property for bi-harmonic functions in half disk.

1. INTRODUCTION

The main result of this paper is the existence of strong minimizer of Blake & Zisserman functional [5] with Dirichlet boundary datum in 2-dimensional image segmentation; the boundary datum is prescribed by penalization.

We refer to [5],[8],[10],[16],[27],[28] for motivation and background analysis of variational approach to image segmentation and digital image processing.

Precisely we focus the functional

$$(1.1) \quad E(K_0, K_1, v) = \int_{\tilde{\Omega} \setminus (K_0 \cup K_1)} |D^2 v|^2 \, d\mathbf{x} + \alpha \mathcal{H}^1(K_0 \cap \tilde{\Omega}) + \beta \mathcal{H}^1((K_1 \setminus K_0) \cap \tilde{\Omega}),$$

with the aim of minimizing it among admissible triplets (K_0, K_1, v) , say triplets fulfilling

$$(1.2) \quad \begin{cases} K_0, K_1 \text{ Borel subsets of } \mathbb{R}^2, & K_0 \cup K_1 \text{ closed,} \\ v \in C^2(\tilde{\Omega} \setminus (K_0 \cup K_1)), & v \text{ approximately continuous in } \tilde{\Omega} \setminus K_0, \\ v = w \text{ a.e. in } \tilde{\Omega} \setminus \Omega. \end{cases}$$

Theorem 1.1. (Strong solution of Dirichlet problem for BZ functional)

Let $\alpha, \beta, \Omega, \tilde{\Omega}, M, T_0, T_1$ and w be s.t.

$$(1.3) \quad 0 < \beta \leq \alpha \leq 2\beta,$$

$$(1.4) \quad \Omega \subset\subset \tilde{\Omega} \subset\subset \mathbb{R}^2,$$

$$(1.5) \quad \Omega \text{ is an open set with Lipschitz boundary}$$

$$(1.6) \quad \exists M \text{ finite set : } (\partial\Omega \setminus M) \in C^2 \text{ uniformly,}$$

$$(1.7) \quad (T_0 \cup T_1) \cap \partial\Omega \text{ is a finite set,}$$

$$(1.8) \quad \mathcal{H}^1((T_0 \cup T_1) \cap \tilde{\Omega}) < +\infty, \quad T_0 \cup T_1 \text{ closed subset of } \mathbb{R}^2,$$

$$(1.9) \quad w \in C^2(\tilde{\Omega} \setminus (T_0 \cup T_1)), \quad w \text{ approximately continuous in } \tilde{\Omega} \setminus T_0,$$

1991 *Mathematics Subject Classification.* 49J45-49K10.

Key words and phrases. Calculus of variations, free discontinuity problems, regularity of weak solutions.

$$(1.10) \quad \begin{cases} D^2w \in L^2(\tilde{\Omega} \setminus (T_0 \cup T_1)), \quad D^2w \in L^\infty(\partial\tilde{\Omega} \setminus (T_0 \cup T_1)) \\ \text{with } \partial\tilde{\Omega} \text{ open set s.t. } \partial\Omega \subset\subset \partial\tilde{\Omega} \subset \tilde{\Omega}, \\ \exists C > 0 : \|w\|_{L^\infty}, \|\nabla w\|_{L^\infty}, \|\nabla^2 w\|_{L^\infty} \leq C \text{ in } \partial\tilde{\Omega}, \\ \text{Lip}(\gamma') \leq C \text{ with } \gamma \text{ arc-length parametrization of } \partial\Omega, \\ \exists \bar{\rho} > 0 : \mathcal{H}^1(\partial\Omega \cap B_{\bar{\rho}}(\mathbf{x})) < C\bar{\rho} \quad \forall \mathbf{x} \in \partial\Omega, \forall \bar{\rho} \leq \bar{\rho}, \end{cases}$$

$$(1.11) \quad \exists (\mathfrak{T}_0, \mathfrak{T}_1, \omega) \text{ fulfilling (1.8), (1.9), } \omega = \text{aplim } w \quad \tilde{\Omega}, (\mathfrak{T}_0 \cup \mathfrak{T}_1) \subsetneq (T_0 \cup T_1).$$

Then there is at least one triplet (C_0, C_1, u) minimizing the functional E defined by (1.1) with finite energy, among admissible triplets (K_0, K_1, v) fulfilling (1.2). Moreover any minimizing triplet (K_0, K_1, v) fulfills:

$$(1.12) \quad K_0 \cap \tilde{\Omega} \text{ and } K_1 \cap \tilde{\Omega} \text{ are } (\mathcal{H}^1, 1) \text{ rectifiable sets,}$$

$$(1.13) \quad \mathcal{H}^1(K_0 \cap \tilde{\Omega}) = \mathcal{H}^1(\overline{S_v}), \quad \mathcal{H}^1(K_1 \cap \tilde{\Omega}) = \mathcal{H}^1(\overline{S_{\nabla v}} \setminus S_v),$$

$$(1.14) \quad \begin{cases} v \in \text{GSBV}^2(\tilde{\Omega}), \text{ hence } v \text{ and } \nabla v \\ \text{have well defined two-sided traces, } \mathcal{H}^1 \text{ a.e. finite on } K_0 \cup K_1, \end{cases}$$

$$(1.15) \quad v \text{ minimizes functional } \mathcal{E} \text{ defined by (2.1) among } v \text{ s.t. } v = w \text{ a.e. } \tilde{\Omega} \setminus \Omega,$$

$$(1.16) \quad \mathcal{E}(v) = E(K_0, K_1, \tilde{v}).$$

Theorem 1.2. Let $\alpha, \beta, \mu, q, g, \Omega, \tilde{\Omega}, M, T_0, T_1$ and w be s.t. (1.3), (1.4), (1.5), (1.6), (1.7), (1.8), (1.9), (1.10), (1.11) and

$$(1.17) \quad \mu > 0, q > 1, g \in L^q(\tilde{\Omega}) \cap L_{loc}^{2q}(\tilde{\Omega}), w \in L^q(\tilde{\Omega})$$

hold true.

Then there is at least one triplet (C_0, C_1, u) minimizing the Blake & Zisserman functional F :

$$(1.18) \quad F(K_0, K_1, v) = E(K_0, K_1, v) + \mu \int_{\tilde{\Omega}} |v - g|^q \, d\mathbf{x},$$

with finite energy, among triplets (K_0, K_1, v) fulfilling (1.2).

Moreover (1.12), (1.13) and (1.14) hold true for any minimizing triplet (K_0, K_1, v) , such v is also a minimizer of the weak functional \mathcal{F} defined by (2.7) under constraint $v = w$ a.e. $\tilde{\Omega} \setminus \Omega$ and

$$(1.19) \quad \mathcal{F}(v) = F(K_0, K_1, \tilde{v}).$$

Theorem 1.3. Assume $\alpha, \beta, \mu, q, g, \Omega, \tilde{\Omega}, M, T_0, T_1$ and w fulfil (1.3)-(1.11), and

$$(1.20) \quad \alpha = \beta.$$

Then there is at least one pair (K, v) minimizing the functional

$$(1.21) \quad \int_{\tilde{\Omega} \setminus K} |D^2v|^2 \, d\mathbf{x} + \alpha \mathcal{H}^1(K \cap \tilde{\Omega})$$

among pairs (K, v) with closed $K \subset \mathbb{R}^2$, $v \in C^2(\tilde{\Omega} \setminus K)$ and $v = w$ a.e. in $\tilde{\Omega} \setminus \Omega$.

If in addition (1.17) holds true then there is at least one pair (K, v) minimizing the functional

$$(1.22) \quad \int_{\tilde{\Omega} \setminus K} \left(|D^2 v|^2 + \mu |v - g|^q \right) dx + \alpha \mathcal{H}^1 \left(K \cap \tilde{\Omega} \right)$$

among pairs (K, v) with closed $K \subset \mathbb{R}^2$, $v \in C^2(\tilde{\Omega} \setminus K)$ and $v = w$ a.e. in $\tilde{\Omega} \setminus \Omega$.

In both cases $K \cap \tilde{\Omega}$ is $(\mathcal{H}^1, 1)$ rectifiable for optimal K .

Moreover:

if the pair (K, v) minimizes (1.22) then v minimizes \mathcal{F} among $v \in GSBV^2(\tilde{\Omega})$ s.t. $v = w$ a.e. $\tilde{\Omega} \setminus \Omega$;

if the pair (K, v) minimizes (1.21) then v minimizes \mathcal{E} among $v \in GSBV^2(\tilde{\Omega})$ s.t. $v = w$ a.e. $\tilde{\Omega} \setminus \Omega$.

Remark 1.4. Thanks to (1.7),(1.8),(1.9) the Dirichlet datum for the Blake & Zisserman functional (1.1) is given as an essential triplet (T_0, T_1, w) (see [9],[16],[18]). The substantial meaning of the boundary condition amounts to impose a penalization whenever the competing function and its gradient do not coincide with the exterior traces of w and ∇w at $\partial\Omega$.

Remark 1.5. About hypothesis (1.7) we notice that it is obviously fulfilled when $(T_0 \cup T_1) \cap \partial\Omega$ is a single point; in such case Theorems 1.1-1.3 entail existence of locally minimizing triplets of E, F with nontrivial Dirichlet data (see [17]): in fact, if T_0 is the negative real axis, $T_1 = \emptyset$ and $\Omega = \mathbb{R}^2$, then Definition 2.6 (locally minimizing triplet) of [17] can be equivalently formulated with $\forall B_\rho(\mathbf{0})$ in place of $\forall A \subset \subset \mathbb{R}^2$.

Remark 1.6. Finiteness of M and $(T_0 \cup T_1) \cap \partial\Omega$ in hypotheses (1.6), (1.7) can be weakened as follows (in Theorems 1.1,1.2,1.3)

$$(1.6') \quad M \text{ closed subset of } \mathbb{R}^2 : \mathcal{H}^1(M) = 0, \quad (1.7') \quad \mathcal{H}^1((T_0 \cup T_1) \cap \partial\Omega) = 0.$$

Remark 1.7. Requiring $D^2 w$ in L^∞ only around the boundary and far away from singular set $(T_0 \cup T_1)$ of Dirichlet datum in (1.10) allows to apply Theorem 1.1 with the choices $T_0 =$ negative real axis, $T_1 = \emptyset$ and $w = W$, where in polar coordinates

$$W(r, \theta) = \pm \sqrt{\frac{\alpha}{193\pi}} r^{3/2} \left(\sqrt{21} \left(\sin \frac{\theta}{2} - \frac{5}{3} \sin \left(\frac{3}{2}\theta \right) \right) \pm \left(\cos \frac{\theta}{2} - \frac{7}{3} \cos \left(\frac{3}{2}\theta \right) \right) \right)$$

is the candidate nontrivial local minimizer for \mathcal{E} in \mathbb{R}^2 (see [17]). Notice that W belongs to $H^2(B_1(\mathbf{0}) \setminus T_0)$ but $D^2 W$ is not bounded around the origin: $D^2 W$ has a singularity of order $r^{-1/2}$.

Remark 1.8. Hypothesis (1.11) is only a technical assumption: actually (1.11) is fulfilled by any Dirichlet datum (T_0, T_1, w) provided the datum is reasonably expressed! More precisely assumption (1.11) entails that (T_0, T_1, w) is an essential triplet (in the sense of Definition 2.11 in [16]).

Theorems 1.1, 1.2, 1.3 are achieved by showing partial regularity of a suitably defined weak solution with penalized Dirichlet datum (Theorem 2.1). The novelty here consists in the regularization at the boundary for a free gradient discontinuity problem; the regularity is proven at points with 2-dimensional energy density by: blow-up (Theorems 6.1,6.2), suitable joining along lunulae filling half-disk (Lemma 3.1) and a decay estimate of weak functionals \mathcal{F} and \mathcal{E} evaluated at local minimizers

(Theorems 6.3, 6.4). When performing such analysis, an essential tool is provided by an L^2 decay estimate of hessian for a bi-harmonic function in a half-disk vanishing together with its normal derivative on the diameter (Theorem 5.1): proving this decay requires a careful application of Duffin extension formula [21] and Almansi decomposition [1], since the bi-harmonic extension to the whole disk may increase a lot the L^2 norm of the hessian in the complementary half-disk. The extension of bi-harmonic functions is quite different from extension of an harmonic function vanishing at the diameter which is based on to classical Schwarz reflection principle (see Remark 5.4) that doubles L^2 norm of the gradient in the whole disk: this doubling property was exploited in [6] to prove decay property for local minimizers of Mumford & Shah functional with Dirichlet boundary condition (see [26]); unfortunately bi-harmonic extension lacks this doubling property (see Remark 5.5).

The present paper focuses the two dimensional case, nevertheless all the results proven here are valid in the n dimensional case except the compactness property (Theorem 4.4) and hessian decay (Theorem 5.1).

About minimization of functionals (1.18), (1.1) under Neumann boundary condition we refer to [8],[9]. About the description of the rich list of (differential, integral and geometric) extremality conditions for (1.18), (1.1) we refer to [16]. The framework and results of the present paper will allow to prove existence of strong local minimizers for Blake & Zisserman functional and several extremality conditions fulfilled by strong local minimizers in forthcoming paper [17]. In general uniqueness of minimizers of functionals F and E fails due to lack of convexity: we refer to [4] for explicit examples of multiplicity and property of generic uniqueness with respect to data α, β, g .

Outline of the paper

1. Introduction
2. Weak Dirichlet problem for Blake & Zisserman functional
3. Joining and matching between lunulae
4. Truncation, Poincaré inequalities and compactness properties in $GSBV$ and $GSBV^2$
5. Hessian decay for bi-harmonic functions in half disk
6. Blow-up and Decay at boundary points
7. Proof of main results
8. References.

2. WEAK DIRICHLET PROBLEM FOR BLAKE & ZISSERMAN FUNCTIONAL

We denote by $B_\varrho(\mathbf{x})$ the open ball $\{\mathbf{y} \in \mathbb{R}^2; |\mathbf{y} - \mathbf{x}| < \varrho\}$, and set $B_\varrho = B_\varrho(\mathbf{0})$, $B_\varrho^+ = B_\varrho \cap \{y > 0\}$, $B_\varrho^- = B_\varrho \cap \{y < 0\}$. We denote by χ_U the characteristic function of U for any $U \subset \mathbb{R}^2$. If x, y are real numbers we denote by $[x]$ the integer part of x and set $x \vee y = \max(x, y)$, $x \wedge y = \min(x, y)$. For any pair of vectors \mathbf{a}, \mathbf{b} we denote the tensor product by $\mathbf{a} \otimes \mathbf{b}$ and set $\mathbf{a} \odot \mathbf{b} = (1/2)(\mathbf{a} \otimes \mathbf{b} + \mathbf{b} \otimes \mathbf{a})$.

For any pair of 2×2 matrices A, B we set $A : B = \sum_{i,j=1}^2 A_{ij} B_{ij}$. For any real $s > 1$ we denote by s' the conjugate exponent $s' = s/(s-1)$.

For any Borel function $v : \Omega \rightarrow \mathbb{R}$ and $\mathbf{x} \in \Omega, z \in \overline{\mathbb{R}} := \mathbb{R} \cup \{-\infty, +\infty\}$, we set $z = \text{ap lim}_{\mathbf{y} \rightarrow \mathbf{x}} v(\mathbf{y})$ (approximate limit of v at \mathbf{x}) if, for every $g \in C^0(\overline{\mathbb{R}})$,

$$g(z) = \lim_{\rho \rightarrow 0} \int_{B_\rho(\mathbf{0})} g(v(\mathbf{x} + \boldsymbol{\xi})) d\boldsymbol{\xi};$$

the function $\tilde{v}(\mathbf{x}) = \text{ap lim}_{\mathbf{y} \rightarrow \mathbf{x}} v(\mathbf{y})$ is called *representative* of v ; the *singular set* of v is $S_v = \{\mathbf{x} \in \Omega : \exists z \text{ s.t. } \text{ap lim}_{\mathbf{y} \rightarrow \mathbf{x}} v(\mathbf{y}) = z\}$.

By referring to [2] and [16]: Dv denotes the distributional gradient of v , $\nabla v(\mathbf{x})$ denotes the approximate gradient of v , $SBV(\Omega)$ denotes the De Giorgi class of functions $v \in BV(\Omega)$ such that

$$\int_{\Omega} |Dv| = \int_{\Omega} |\nabla v| \, d\mathbf{x} + \int_{S_v} |v^+ - v^-| \, d\mathcal{H}^1.$$

$$SBV_{loc}(\Omega) := \{v \in SBV(\Omega') : \forall \Omega' \subset\subset \Omega\},$$

$$GSBV(\Omega) := \{v : \Omega \rightarrow \mathbb{R} \text{ Borel function; } -k \vee v \wedge k \in SBV_{loc}(\Omega) \forall k \in \mathbb{N}\}.$$

$$GSBV^2(\Omega) := \{v \in GSBV(\Omega), \nabla v \in (GSBV(\Omega))^2\}.$$

We will exploit the weak formulation \mathcal{E} of functional E introduced in [8]:

$$(2.1) \quad \mathcal{E}(v) = \int_{\tilde{\Omega} \setminus (S_v \cup S_{\nabla v})} |\nabla^2 v|^2 \, d\mathbf{x} + \alpha \mathcal{H}^1(S_v) + \beta \mathcal{H}^1(S_{\nabla v} \setminus S_v).$$

Theorem 2.1. (Dirichlet problem for weak form of Blake & Zisserman: functionals \mathcal{F} and \mathcal{E})

Assume (1.3), (1.4), (1.5) and

$$(2.2) \quad w \in C^2(\tilde{\Omega} \setminus \overline{(S_w \cup S_{\nabla w})}), \quad w \text{ approximately continuous in } \tilde{\Omega} \setminus S_w,$$

$$(2.3) \quad \mathcal{E}(w) < +\infty$$

$$(2.4) \quad \mathcal{H}^1(\overline{(S_w \cup S_{\nabla w})} \setminus (S_w \cup S_{\nabla w})) = 0,$$

$$(2.5) \quad \mathcal{H}^1(\overline{(S_w \cup S_{\nabla w})} \cap \partial\Omega) = 0 \quad (\text{or } \overline{(S_w \cup S_{\nabla w})} \cap \partial\Omega \text{ finite}).$$

Set

$$(2.6) \quad X(\tilde{\Omega}) \stackrel{\text{def}}{=} \left\{ v \in GSBV^2(\tilde{\Omega}) \text{ s.t. } v = w \text{ a.e. in } \tilde{\Omega} \setminus \Omega \right\}.$$

Then there is at least one u minimizing functional \mathcal{E} in $X(\tilde{\Omega})$ with finite energy.

Moreover, if $\mu > 0$, $g \in L^q(\tilde{\Omega})$ and $w \in L^q(\tilde{\Omega})$, then there is at least one u minimizing functional \mathcal{F} in $X(\tilde{\Omega})$ with finite energy:

$$(2.7) \quad \mathcal{F}(v) = \int_{\tilde{\Omega} \setminus (S_v \cup S_{\nabla v})} |\nabla^2 v|^2 \, d\mathbf{x} + \alpha \mathcal{H}^1(S_v) + \beta \mathcal{H}^1(S_{\nabla v} \setminus S_v) + \mu \int_{\tilde{\Omega}} |v - g|^q \, d\mathbf{x}$$

Proof. Obviously $\mathcal{F}(v) \geq 0 \forall v \in X(\tilde{\Omega})$.

Assumptions (2.2), (2.3), (2.4), (2.5) entail $w \in X(\tilde{\Omega})$ and $\mathcal{F}(w) < +\infty$.

Let $v_h \in X$ be a minimizing sequence for \mathcal{F} . By Theorem 8 in [8] there is v_∞ in $X(\tilde{\Omega})$ and a subsequence s.t., without relabeling, $v_h \rightarrow v_\infty$ a.e. in $\tilde{\Omega}$.

$v_h = w$ in $\tilde{\Omega} \setminus \Omega$ entails $v_\infty = w$ in $\tilde{\Omega} \setminus \Omega$. By Theorem 10 in [8]:

$$\mathcal{F}(v_\infty) \leq \liminf_h \mathcal{F}(v_h),$$

hence $\mathcal{F}(v_\infty) = \inf_{v \in X(\tilde{\Omega})} \mathcal{F}(v)$. The same argument applies to \mathcal{E} . \square

Remark 2.2. *Assumptions of Theorems 1.1, 1.2 and 1.3 (about Dirichlet datum for strong formulation) on triplets (T_0, T_1, w) entails the assumptions (about Dirichlet datum for weak formulation) on w of Theorem 2.1.*

Remark 2.3. *So far we know that both \mathcal{E} and \mathcal{F} achieve finite minimum under Dirichlet boundary condition provided the structural assumptions ((1.3)-(1.11) and (1.17)) of the paper hold true. We want to show that also E, F have the same property.*

Definition 2.4. About functionals defined by (1.1), (1.18), (2.1), (2.7) we will often use the short notation $E, F, \mathcal{E}, \mathcal{F}$; nevertheless, whenever required by clearness of exposition about interchange of various ingredients (functions, parameters, sets, Dirichlet datum) we will use several different (self-explaining) notation:

$$\mathcal{F}(v), \mathcal{F}_g(v), \mathcal{F}_{gw}(v), \mathcal{F}(v, A), \mathcal{F}_{gw}(v, \mu, \alpha, \beta, A); \quad F(K_0, K_1, v), F_g(K_0, K_1, v), \\ \mathcal{E}(v), \mathcal{E}(v, A), \mathcal{E}(v, \alpha, \beta, A); \quad E(K_0, K_1, v).$$

Lemma 2.5. (Scaling) *Let $v \in GSBV^2(B_r(\mathbf{x}_0))$ where $\mathbf{x}_0 = (x_0, y_0)$. For $\lambda > 0$ and for every $\mathbf{x} \in B_1$ set*

$$(2.8) \quad v_r(\mathbf{x}) = \frac{v(\mathbf{x}_0 + r\mathbf{x})}{\lambda^{1/2} r^{3/2}}, \quad g_r(\mathbf{x}) = \frac{g(\mathbf{x}_0 + r\mathbf{x})}{\lambda^{1/2} r^{3/2}}.$$

Then $v_r \in GSBV^2(B_1)$ and

$$(2.9) \quad \mathcal{F}_g(v, \mu, \alpha, \beta, B_r(\mathbf{x}_0)) = \lambda r \mathcal{F}_{g_r}(v_r, \mu \lambda^{\frac{q}{2}-1} r^{1+\frac{3}{2}q}, \frac{\alpha}{\lambda}, \frac{\beta}{\lambda}, B_1)$$

and, by setting $K_{0r} = (K_0 - \mathbf{x}_0)/r$, $K_{1r} = (K_1 - \mathbf{x}_0)/r$,

$$(2.10) \quad F_g(K_0, K_1, v, \mu, \alpha, \beta, B_r(\mathbf{x}_0)) = \lambda r F_g(K_{0r}, K_{1r}, v_r, \mu \lambda^{\frac{q}{2}-1} r^{1+\frac{3}{2}q}, \frac{\alpha}{\lambda}, \frac{\beta}{\lambda}, B_1).$$

Proof. The thesis follows by change of variables. \square

Definition 2.6. For any $\mathbf{x} \in \tilde{\Omega}$ and r s.t. $0 < r < \text{dist}(\mathbf{x}, \partial\tilde{\Omega})$, we say that u is an Ω local minimizer of $\mathcal{F}_{gw}(\cdot, \mu, \alpha, \beta, A)$ if

$$(2.11) \quad \left\{ \begin{array}{l} u \in GSBV^2(B_r(\mathbf{x})): u = w \text{ a.e. } B_r(\mathbf{x}) \setminus \Omega, \mathcal{F}_{gw}(u, \mu, \alpha, \beta, A) < +\infty, \\ \mathcal{F}_{gw}(u, \mu, \alpha, \beta, A) \leq \mathcal{F}_{gw}(u + \eta, \mu, \alpha, \beta, A) \\ \forall A \subset\subset B_r(\mathbf{x}), \forall \eta \in GSBV^2(B_r(\mathbf{x})) : \text{spt } \eta \subset A, \eta = 0 \text{ a.e. } B_r(\mathbf{x}) \setminus \Omega. \end{array} \right.$$

Definition 2.7. For any $\mathbf{x} \in \tilde{\Omega}$ and r s.t. $0 < r < \text{dist}(\mathbf{x}, \partial\tilde{\Omega})$, we say that u is an Ω local minimizer of $\mathcal{E}_w(\cdot, \alpha, \beta, A)$ if

$$(2.12) \quad \left\{ \begin{array}{l} u \in GSBV^2(B_r(\mathbf{x})): u = w \text{ a.e. } B_r(\mathbf{x}) \setminus \Omega, \mathcal{E}_w(u, \alpha, \beta, A) < +\infty, \\ \mathcal{E}_w(u, \alpha, \beta, A) \leq \mathcal{E}_w(u + \eta, \alpha, \beta, A) \\ \forall A \subset\subset B_r(\mathbf{x}), \forall \eta \in GSBV^2(B_r(\mathbf{x})) : \text{spt } \eta \subset A, \eta = 0 \text{ a.e. } B_r(\mathbf{x}) \setminus \Omega. \end{array} \right.$$

Remark 2.8. *Definitions 2.6 and 2.7 will be used also with open sets different from Ω provided suitable data g, w are defined in the contest.*

Definitions 2.6 and 2.7 though different from the ones given in [9] are in fact equivalent to them since sublevels of energy \mathcal{F} (or \mathcal{E}) are linear subspaces of GSBV² (due to Corollary 4.5 in [3]).

Remark 2.9. Due to Lemma 2.5, if u is a $B_\varrho(\mathbf{x})$ local minimizer of $\mathcal{F}_{g,w}(\cdot, \mu, \alpha, \beta, B_\varrho(\mathbf{x}))$ then for any \mathbf{a}, λ, c

$$\mathbf{y} \rightarrow u(\mathbf{y}) = \lambda^{-1/2} \varrho^{-3/2} (u(\mathbf{x} + \varrho \mathbf{y}) - \varrho \mathbf{a} \cdot \mathbf{y} - c)$$

is a $B_1(\mathbf{0})$ local minimizer of $\mathcal{F}_{\gamma,\omega}(\cdot, \mu \lambda^{\frac{q}{2}-1} r^{1+\frac{3}{2}q}, \alpha/\lambda, \beta/\lambda, B_1(\mathbf{0}))$ where

$$\begin{aligned} \gamma &= \lambda^{-1/2} \varrho^{-3/2} (g(\mathbf{x} + \varrho \mathbf{y}) - \varrho \mathbf{a} \cdot \mathbf{y} - c) \\ \omega &= \lambda^{-1/2} \varrho^{-3/2} (w(\mathbf{x} + \varrho \mathbf{y}) - \varrho \mathbf{a} \cdot \mathbf{y} - c), \end{aligned}$$

The same property holds true for \mathcal{E}_w .

Now we prove a density upper bound for the functional \mathcal{F} near the points $\mathbf{x} \in \partial\Omega$ analogous to the estimate in [16]: Thm 2.12 and Rmk 2.13.

Theorem 2.10. (Density upper bound for the functional \mathcal{F} at the boundary) Let u be a minimizer in $X(\tilde{\Omega})$ for the functional \mathcal{F} with (1.3)–(1.5), (1.17), (2.2)–(2.6) and

$$(2.13) \quad \exists \bar{\varrho} > 0 : \mathcal{H}^1(\partial\Omega \cap B_{\bar{\varrho}}(\mathbf{x})) < C\bar{\varrho} \quad \forall \mathbf{x} \in \partial\Omega, \quad \forall \bar{\varrho} \leq \bar{\varrho}.$$

Then for every $0 < \varrho \leq \bar{\varrho} \wedge 1$ and for every $\mathbf{x} \in \partial\Omega \setminus (\overline{S_w} \cup \overline{S_{\nabla w}})$ such that $\overline{B_\varrho}(\mathbf{x}) \subset \tilde{\Omega}$ we have

$$(2.14) \quad \mathcal{F}(u, \overline{B_\varrho}(\mathbf{x})) \leq c_0 \varrho$$

where $c_0 = C^2 \pi + 2^{q-1} \pi^{\frac{1}{2}} \mu (\|w\|_{L^{2q}(B_\varrho(\mathbf{x}))}^q + \|g\|_{L^{2q}(B_\varrho(\mathbf{x}))}^q) + (2\pi + C)\alpha$.

If $q = 2$ and $g, w \in L^\infty(\tilde{\Omega})$, then we can choose

$$c_0 = C^2 \pi + 2\pi \mu (\|w\|_{L^\infty}^2 + \|g\|_{L^\infty}^2) + (2\pi + C)\alpha.$$

Proof. By minimality of u for \mathcal{F} we get

$$\mathcal{F}(u) \leq \mathcal{F}(v),$$

where

$$v = u \chi_{\tilde{\Omega} \setminus (B_\varrho(\mathbf{x}) \cap \Omega)}.$$

Taking into account $\beta \leq \alpha$, since $\mathcal{F}(u, \tilde{\Omega} \setminus \overline{B_\varrho}(\mathbf{x})) = \mathcal{F}(v, \tilde{\Omega} \setminus \overline{B_\varrho}(\mathbf{x}))$ then

$$\begin{aligned} \mathcal{F}(u, \overline{B_\varrho}(\mathbf{x})) &\leq \mathcal{F}(v, \overline{B_\varrho}(\mathbf{x})) \leq \int_{B_\varrho(\mathbf{x}) \setminus \tilde{\Omega}} (|\nabla^2 w|^2 + \mu |w - g|^q) d\mathbf{y} + \mu \int_{B_\varrho(\mathbf{x}) \cap \Omega} |g|^q d\mathbf{y} \\ &\quad + \alpha \mathcal{H}^1(\partial B_\varrho(\mathbf{x}) \cap \Omega) + \alpha \mathcal{H}^1(\partial\Omega \cap B_\varrho(\mathbf{x})) \\ &\leq C^2 \pi \varrho^2 + 2^{q-1} \mu \int_{B_\varrho(\mathbf{x}) \setminus \tilde{\Omega}} (|w|^q + |g|^q) d\mathbf{y} \\ &\quad + \mu \int_{B_\varrho(\mathbf{x}) \cap \Omega} |g|^q d\mathbf{y} + 2\pi \alpha \varrho + \alpha \mathcal{H}^1(\partial\Omega \cap B_\varrho(\mathbf{x})) \\ &\leq C^2 \pi \varrho^2 + 2^{q-1} \mu (\|w\|_{L^{2q}(B_\varrho(\mathbf{x}))}^q + \|g\|_{L^{2q}(B_\varrho(\mathbf{x}))}^q) (\pi \varrho^2)^{\frac{1}{2}} \\ &\quad + 2\pi \alpha \varrho + C \alpha \varrho, \end{aligned}$$

hence we achieve the proof. \square

3. JOINING AND MATCHING BETWEEN LUNULAE

In this Section we prove some technical tools aimed to the proof of partial regularity at boundary points.

Lemma 3.1. (Joining between lunulae)

Assume (1.3), (1.4), (1.17), z, u in $GSBV^2(\tilde{\Omega})$, $\mathbf{x}_0 = (x_0, y_0) \in \partial\Omega$, $0 < d < \sigma < s < t < 1$ and $\sigma - d < t - s$ s.t. $\overline{B_t(\mathbf{x}_0)} \subset \tilde{\Omega}$, $\partial\Omega \cap B_t(\mathbf{x}_0) \in C^2$ and \mathbf{d} denotes the inner normal to $\partial\Omega$ at \mathbf{x}_0 . Set

$$(3.1) \quad B_t^d = B_t(\mathbf{x}_0) \cap \{(\mathbf{x} - \mathbf{x}_0) \cdot \mathbf{d} > d\}, \quad M_{t,s}^{d,\sigma} = \overline{B_t^d} \setminus B_s^\sigma.$$

Then for every $\theta \in (0, 1)$ there are $c = c(\theta) > 0$ and a cut-off function Ψ in $C^2(B_t(\mathbf{x}_0)) \cap C_0^2(B_t^d(\mathbf{x}_0))$ s.t. $\Psi \equiv 1$ in a neighborhood of $B_s^\sigma = B_s(\mathbf{x}_0) \cap \{(\mathbf{x} - \mathbf{x}_0) \cdot \mathbf{d} > \sigma\}$ and, by setting

$$U = \Psi u + (1 - \Psi)z$$

we have

$$\begin{aligned} \mathcal{F}(U, \overline{B_t^d}) &\leq (1 + \theta) \left(\mathcal{F}(u, \overline{B_t^d}) + \mathcal{F}(z, \overline{B_t^d} \setminus B_s^\sigma) \right) + \\ &+ \frac{c}{(\sigma - d)^2} \left(\int_{B_t^d \setminus B_s^\sigma} |\nabla(u - z)|^2 d\mathbf{x} + \frac{c}{\theta d^2 (\sigma - d)^2} \int_{B_t^d \setminus B_s^\sigma} |u - z|^2 d\mathbf{x} \right) \end{aligned}$$

and

$$\begin{aligned} \mathcal{E}(U, \overline{B_t^d}) &\leq (1 + \theta) \left(\mathcal{E}(u, \overline{B_t^d}) + \mathcal{E}(z, \overline{B_t^d} \setminus B_s^\sigma) \right) + \\ &+ \frac{c}{(\sigma - d)^2} \left(\int_{B_t^d \setminus B_s^\sigma} |\nabla(u - z)|^2 d\mathbf{x} + \frac{c}{\theta d^2 (\sigma - d)^2} \int_{B_t^d \setminus B_s^\sigma} |u - z|^2 d\mathbf{x} \right) \end{aligned}$$

Proof. Again it will be enough proving the estimates for the terms containing $|\nabla \cdot|^2$. Without loss of generality we assume $\mathbf{x}_0 = \mathbf{0}$ and $\mathbf{d} = \mathbf{e}_2$ so that

$$\{(\mathbf{x} - \mathbf{x}_0) \cdot \mathbf{d} > d\} = \{y > d\}$$

We fix $\theta \in (0, 1)$ and $N = N(\theta)$ where $N = 1 + [C/\theta]$ and C is a suitable constant. Let

$$s_j = s + j \frac{t - s}{N}, \quad j = 0, \dots, N,$$

$$d_0 = \sigma, \quad d_j = d_0 - j \frac{\sigma - d}{N}, \quad j = 0, \dots, N,$$

and ψ_j ($j = 0, \dots, N - 1$) a list of C^2 cut-off functions between B_{s_j} and $B_{s_{j+1}}$ (say $0 \leq \psi_j \leq 1$, $\psi_j \equiv 1$ in a neighborhood of B_{s_j} , ψ_j vanishes outside $\overline{B_{s_{j+1}}}$) with

$$|D\psi_j| \leq \frac{2N}{\sigma - d}, \quad |D^2\psi_j| \leq \frac{8N^2}{d(\sigma - d)^2}, \quad \text{in } C_j \stackrel{\text{def}}{=} B_{s_{j+1}} \setminus B_{s_j}$$

and a list of 1-dimensional cut-off functions η_j , $j = 1, \dots, N$, between $\{y > d_j\}$ and $\{y > d_{j+1}\}$ (say $0 \leq \eta_j \leq 1$, $\eta_j \equiv 1$ in a neighborhood of $\{y > d_j\}$, η_j vanishes outside $\{y > d_{j+1}\}$) with

$$|D\eta_j| \leq \frac{2N}{\sigma - d}, \quad |D^2\eta_j| \leq \frac{2N^2}{(\sigma - d)^2}, \quad \text{in } E_j \stackrel{\text{def}}{=} \{d_{j+1} < y < d_j\}$$

Then define

$$U_j = \eta_j \psi_j u + (1 - \eta_j \psi_j) z.$$

For any w ,

$$\nabla^2 (\eta_j \psi_j w) = \eta_j \nabla^2 (\psi_j w) + \begin{pmatrix} 0 & (\eta_j)_y (\psi_j w)_x \\ (\eta_j)_y (\psi_j w)_x & (\eta_j)_{yy} \psi_j w \end{pmatrix}$$

We introduce the handles $M_j \stackrel{\text{def}}{=} (C_j \cap \{y > d_{j+1}\}) \cup (E_j \cap \{|\mathbf{x}| < s_{j+1}\})$ for $j = 1, \dots, N-1$ and the lunula $M_0 = E_0 \cap \{|\mathbf{x}| < s\}$: we notice that the sets M_j , $j = 0, \dots, N-1$, are pair-wise disjoint ($j \neq k \Rightarrow M_j \cap M_k = \emptyset$) and their union covers the whole lunula $\{|\mathbf{x}| < t\} \cap \{y > d\}$ up to a set of measure 0.

Since ψ_j is a radial function we obtain, for every j ,

$$\begin{aligned} \int_{B_t^d} |\nabla^2 U_j|^2 d\mathbf{x} &\leq \int_{B_{s_j}^d} |\nabla^2 u|^2 d\mathbf{x} + \int_{B_t^d \setminus B_{s_j}^{d_{j+1}}} |\nabla^2 z|^2 d\mathbf{x} \\ &\quad + \int_{M_j} |\eta_j (\psi_j \nabla^2 u + (1 - \psi_j) \nabla^2 z + 2D\psi_j \odot \nabla(u - z) + D^2\psi_j (u - z)) \\ &\quad + (\mathbf{e}_1 \otimes \mathbf{e}_2 + \mathbf{e}_2 \otimes \mathbf{e}_1) (\eta_j)_y (\psi_j u)_x + \mathbf{e}_2 \otimes \mathbf{e}_2 (\eta_j)_{yy} \psi_j u \\ &\quad + (\mathbf{e}_1 \otimes \mathbf{e}_2 + \mathbf{e}_2 \otimes \mathbf{e}_1) (\eta_j)_y ((1 - \psi_j)z)_x + \mathbf{e}_2 \otimes \mathbf{e}_2 (\eta_j)_{yy} (1 - \psi_j)z|^2 d\mathbf{x} \\ &\leq \int_{B_t^d} |\nabla^2 u|^2 d\mathbf{x} + \int_{B_t^d \setminus B_s^\sigma} |\nabla^2 z|^2 d\mathbf{x} \\ &\quad + C \int_{M_j} \left(|\nabla^2 u|^2 + |\nabla^2 z|^2 + |D\psi_j|^2 |\nabla(u - z)|^2 + |D^2\psi_j|^2 |u - z|^2 \right. \\ &\quad \left. + |D\eta_j|^2 |D\psi_j|^2 |u - z|^2 + |D\eta_j|^2 |\nabla(u - z)|^2 + |D^2\eta_j|^2 |u - z|^2 \right) d\mathbf{x}. \end{aligned}$$

By taking into account that $\text{spt}(\eta_j \psi_j) \subset \bigcup_{k=0}^{j+1} \overline{M_k}$ we add the last inequalities with respect to j from 0 to $N-1$:

$$\begin{aligned} \min_j \int_{B_t^d} |\nabla^2 U_j|^2 d\mathbf{x} &\leq \int_{B_t^d} |\nabla^2 u|^2 d\mathbf{x} + \int_{B_t^d \setminus B_s^\sigma} |\nabla^2 z|^2 d\mathbf{x} \\ &\quad + \frac{C}{N} \int_{B_t^d \setminus B_s^\sigma} \left(|\nabla^2 u|^2 + |\nabla^2 z|^2 + \left(\frac{2N}{\sigma - d} \right)^2 |\nabla(u - z)|^2 + \left(\frac{12N^2}{d(\sigma - d)^2} \right)^2 |u - z|^2 \right) d\mathbf{x} \end{aligned}$$

We select the index j achieving such minimum and set $U = U_j$. Hence

$$\begin{aligned} \int_{B_t^d} |\nabla^2 U|^2 d\mathbf{x} &\leq \int_{B_t^d} |\nabla^2 u|^2 d\mathbf{x} + \int_{B_t^d \setminus B_s^\sigma} |\nabla^2 z|^2 d\mathbf{x} \\ &\quad + \theta \int_{B_t^d \setminus B_s^\sigma} \left(|\nabla^2 u|^2 + |\nabla^2 z|^2 + \left(\frac{2(C+1)}{\theta(\sigma - d)} \right)^2 |\nabla(u - z)|^2 + \left(\frac{12(C+1)^2}{\theta d(\sigma - d)^2} \right)^2 |u - z|^2 \right) d\mathbf{x} \end{aligned}$$

and the thesis follows by inequalities with $c = 12(C+1)^2/\theta$, $\Psi = \eta_j \psi_j$, since the terms not containing ∇^2 fulfill the inequality in the thesis with $\theta = 0$:

$$\begin{aligned} \mathcal{H}^1 \left(S_U \cap \overline{B_t^d} \right) &= \mathcal{H}^1 \left(S_u \cap \overline{B_{s_j}^{d_j}} \right) + \mathcal{H}^1 \left(S_z \cap (\overline{B_t^d} \setminus B_{s_{j+1}^{d_{j+1}}}) \right) \\ (3.2) \quad &\quad + \mathcal{H}^1 \left(S_u \cap (\overline{B_{s_{j+1}^{d_{j+1}}}} \setminus B_{s_j}^{d_j}) \right) + \mathcal{H}^1 \left(S_z \cap (\overline{B_{s_{j+1}^{d_{j+1}}}} \setminus B_{s_j}^{d_j}) \right), \end{aligned}$$

$$\begin{aligned}
& \mathcal{H}^1 \left((S_{\nabla U} \setminus S_U) \cap \overline{B_t^d} \right) = \\
(3.3) \quad & = \mathcal{H}^1 \left((S_{\nabla u} \setminus S_u) \cap \overline{B_{s_j}^{d_j}} \right) + \mathcal{H}^1 \left((S_{\nabla z} \setminus S_z) \cap (\overline{B_t^d} \setminus B_{s_{j+1}}^{d_{j+1}}) \right) \\
& + \mathcal{H}^1 \left((S_{\nabla u} \setminus S_v) \cap (\overline{B_{s_{j+1}}^{d_{j+1}}} \setminus B_{s_j}^{d_j}) \right) + \mathcal{H}^1 \left((S_{\nabla z} \setminus S_z) \cap (\overline{B_{s_{j+1}}^{d_{j+1}}} \setminus B_{s_j}^{d_j}) \right),
\end{aligned}$$

$$\begin{aligned}
(3.4) \quad \int_{B_t^d} |U - g|^q d\mathbf{x} & \leq \int_{B_{s_j}^{d_j}} |u - g|^q d\mathbf{x} + \int_{B_t^d \setminus B_{s_{j+1}}^{d_{j+1}}} |z - g|^q d\mathbf{x} \\
& + \int_{\overline{B_{s_{j+1}}^{d_{j+1}}} \setminus B_{s_j}^{d_j}} (\psi_j |u - g|^q + (1 - \psi_j) |z - g|^q) d\mathbf{x}.
\end{aligned}$$

□

Lemma 3.2. (Matching with lunulae) Let $\mathbf{x}_0 = (x_0, y_0)$, $z, v \in GSBV^2(\tilde{\Omega})$, $\overline{B_t}(\mathbf{x}_0) \subset \tilde{\Omega}$ and

$$\mathcal{H}^1(S_z \cap \partial B_t^d) = \mathcal{H}^1(S_{\nabla z} \cap \partial B_t^d) = \mathcal{H}^1(S_v \cap \partial B_t^d) = \mathcal{H}^1(S_{\nabla v} \cap \partial B_t^d) = 0$$

where $B_t^d = B_t(\mathbf{x}_0) \cap \{y > y_0 + d\}$. Then, by setting

$$u = \begin{cases} z & \text{in } B_t^d \\ v & \text{in } \tilde{\Omega} \setminus B_t^d \end{cases}$$

we have

$$\begin{aligned}
\mathcal{F}_g(u, \mu, \alpha, \beta, \tilde{\Omega}) & \leq \mathcal{F}_g(z, \mu, \alpha, \beta, B_t^d) + \mathcal{F}_g(v, \mu, \alpha, \beta, \tilde{\Omega} \setminus B_t^d) + \\
& + \alpha \mathcal{H}^1(\{\tilde{z} \neq \tilde{v}\} \cap \partial B_t^d) + \beta \mathcal{H}^1\left(\left(\{\tilde{\nabla} z \neq \tilde{\nabla} v\} \setminus \{\tilde{z} \neq \tilde{v}\}\right) \cap \partial B_t^d\right), \\
\mathcal{E}(u, \alpha, \beta, \tilde{\Omega}) & \leq \mathcal{E}(z, \alpha, \beta, B_t^d) + \mathcal{E}(v, \alpha, \beta, \tilde{\Omega} \setminus B_t^d) + \\
& + \alpha \mathcal{H}^1(\{\tilde{z} \neq \tilde{v}\} \cap \partial B_t^d) + \beta \mathcal{H}^1\left(\left(\{\tilde{\nabla} z \neq \tilde{\nabla} v\} \setminus \{\tilde{z} \neq \tilde{v}\}\right) \cap \partial B_t^d\right).
\end{aligned}$$

Proof. The thesis follows by the definitions. □

Lemma 3.3. Let $v \in GSBV^2(\Omega)$ s.t.

$$\mathcal{F}_g(v, \mu, \alpha, \beta, T) < +\infty \quad \forall \text{ compact set } T \subset \Omega.$$

Then

$$\lim_{\varrho \rightarrow 0} \varrho^{-1} \mathcal{F}_g(v, \mu, \alpha, \beta, B_\varrho(\mathbf{x})) = 0 \quad \text{for } \mathcal{H}^1 \text{ a.e. } \mathbf{x} \in \Omega \setminus (S_v \cup S_{\nabla v}).$$

Proof. Apply the same argument of Lemma 2.6 in [20]. □

4. TRUNCATION, POINCARÉ INEQUALITIES AND COMPACTNESS PROPERTIES IN $GSBV$ AND $GSBV^2$

We recall a Poincaré-Wirtinger type inequality in the class $GSBV$ which was proven in [9] allowing surgical truncations of non integrable functions of several variables and we refine its statement with the aim of taming blow-up at boundary points in case of functions vanishing in a full sector. We emphasize that $v \in GSBV^2(\Omega)$ does not even entail that either v or ∇v belongs to $L^1_{loc}(\Omega)$.

Let B be an open ball in \mathbb{R}^2 . For every measurable function $v : B \rightarrow \mathbb{R}$ we define the least median of v in B as

$$\text{med}(v, B) = \inf \{ t \in \mathbb{R}; |\{v < t\} \cap B| \geq \frac{1}{2}|B| \}.$$

We remark that $\text{med}(\cdot, B)$ is a non linear operator and in general it has no relationship with the averaged integral $\int_B \cdot dy / |B|$.

Obviously we have $\text{med}(v\chi_{B \setminus E} + \text{med}(v, B)\chi_E, B) = \text{med}(v, B)$ for every $E \subset B$. For every $v \in GSBV(B)$ and $a \in \mathbb{R}$ with $(2\gamma_2\mathcal{H}^1(S_v))^2 \leq a \leq \frac{1}{2}|B|$, we set

$$\tau'(v, a, B) = \inf \{ t \in \mathbb{R}; |\{v < t\}| \geq a \},$$

$$\tau''(v, a, B) = \inf \{ t \in \mathbb{R}; |\{v \geq t\}| \leq a \},$$

here γ_2 is the isoperimetric constant relative to the balls of \mathbb{R}^2 , i.e.

$$\min\{|E \cap B|^{\frac{1}{2}}, |B \setminus E|^{\frac{1}{2}}\} \leq \gamma_2 P(E, B) \quad \forall \text{ Borel set } E,$$

and $P(E, B)$ denotes the perimeter of E in $B : P(E, B) = \int_B |D\chi_E|$.

For $\eta \geq 0$ we define the truncation operator

$$(4.1) \quad T(v, a, \eta) = (\tau'(v, a, B) - \eta) \vee v \wedge (\tau''(v, a, B) + \eta).$$

We get easily $T(T(v, a, \eta), a, \eta) = T(v, a, \eta)$, $\text{med}(T(v, a, \eta), B) = \text{med}(v, B)$ and $T(\lambda v, a, \lambda\eta) = \lambda T(v, a, \eta)$ for every $\lambda > 0$. Moreover $|\nabla T(v, a, \eta)| \leq |\nabla v|$ a.e. on B and

$$(4.2) \quad |\{v \neq T(v, a, \eta)\}| \leq 2a.$$

In case v is vector-valued the operators med and T are defined componentwise.

For any given function in $GSBV$, we define an affine polynomial correction such that both median and gradient median vanish.

Let $B_r(\mathbf{x}) \subset \Omega$ and $v \in GSBV(B_r(\mathbf{x}))$; for every $\mathbf{y} \in \mathbb{R}^2$ we set

$$(4.3) \quad (M_{\mathbf{x}, r} v)(\mathbf{y}) = \text{med}(\nabla v, B_r(\mathbf{x})) \cdot (\mathbf{y} - \mathbf{x})$$

$$(4.4) \quad (\mathcal{P}_{\mathbf{x}, r} v)(\mathbf{y}) = (M_{\mathbf{x}, r} v)(\mathbf{y}) + \text{med}(v - M_{\mathbf{x}, r} v, B_r(\mathbf{x})).$$

Since $\text{med}(v - c, B_r(\mathbf{x})) = \text{med}(v, B_r(\mathbf{x})) - c$ for every $c \in \mathbb{R}$ and $\nabla(\mathcal{P}_{\mathbf{x}, r} v) = \nabla(M_{\mathbf{x}, r} v) = \text{med}(\nabla v, B_r(\mathbf{x}))$ then we have $\mathcal{P}_{\mathbf{x}, r}(v - \mathcal{P}_{\mathbf{x}, r} v) = 0$, say

$$\text{med}(v - \mathcal{P}_{\mathbf{x}, r} v, B_r(\mathbf{x})) = 0, \quad \text{med}(\nabla(v - \mathcal{P}_{\mathbf{x}, r} v), B_r(\mathbf{x})) = \mathbf{0}.$$

We notice that there are v such that $\text{med}(v, B_r(\mathbf{x})) \neq \text{med}(\mathcal{P}_{\mathbf{x}, r} v, B_r(\mathbf{x}))$, take e.g. $v(x, y) = (x^2 - x)H(-x) - \frac{x}{2}H(x)$, where H is the Heaviside function.

The following statement was proven by Theorem 4.1 in [9].

Theorem 4.1. (Poincaré-Wirtinger inequality for $GSBV$ functions in a ball)

Let $B \subset \mathbb{R}^2$ be an open ball, $v \in GSBV(B)$ and $a \in \mathbb{R}$ with

$$(4.5) \quad (2\gamma_2\mathcal{H}^1(S_v))^2 \leq a \leq \frac{1}{2}|B|,$$

let $\eta \geq 0$ and $T(v, a, \eta)$ as in (4.1). Then

$$(4.6) \quad \int_B |DT(v, a, \eta)| \leq 2|B|^{\frac{1}{2}} \left(\int_B |\nabla T(v, a, \eta)|^2 dy \right)^{\frac{1}{2}} + 2\eta\mathcal{H}^1(S_v).$$

We have also, for every $s \geq 2$,

$$(4.7) \quad \int_B |T(v, a, \eta) - \text{med}(v, B)|^s dy \leq 2^{s-1} (\gamma_2 s)^s \left(\int_B |\nabla T(v, a, 0)|^2 dy \right)^{\frac{s}{2}} |B| + (2\eta)^s a.$$

Theorem 4.2. (Classical Poincaré inequality in BV [22] Thm. 5.6.1(iii))
For any $\mathbf{x} \in \mathbb{R}^2$, $r > 0$, and $0 < \vartheta \leq 1$ there is K_ϑ such that

$$(4.8) \quad \|v\|_{L^2(B_r(\mathbf{x}))} \leq K_\vartheta \int_{B_r(\mathbf{x})} |Dv| \quad \forall v \in BV(B_r(\mathbf{x})) \text{ s.t.}$$

$$(4.9) \quad |\{\mathbf{y} \in B_r(\mathbf{x}) : v(\mathbf{y}) = 0\}| / |B_r(\mathbf{x})| \geq \vartheta.$$

Theorem 4.3. (Poincaré-Wirtinger inequality for GSBV functions vanishing in a sector)

Let $B \subset \mathbb{R}^2$ be an open ball, $v \in GSBV(B)$ s.t. (4.9) holds true and $a \in \mathbb{R}$ with

$$(4.10) \quad (2\gamma_2 \mathcal{H}^1(S_v))^2 \leq a \leq \frac{1}{2}|B|,$$

let $\eta \geq 0$ and $T(v, a, \eta)$ as in (4.1). Then

$$(4.11) \quad \int_B |DT(v, a, \eta)| \leq 2|B|^{\frac{1}{2}} \left(\int_B |\nabla T(v, a, \eta)|^2 dy \right)^{\frac{1}{2}} + 2\eta \mathcal{H}^1(S_v).$$

We have also, for every $s \geq 2$,

$$(4.12) \quad \int_B |T(v, a, \eta)|^s dy \leq 2^{s-1} (K_\vartheta s)^s \left(\int_B |\nabla T(v, a, 0)|^2 dy \right)^{\frac{s}{2}} |B| + (2\eta)^s a.$$

Proof. Identical to the proof of Theorem 4.1 in [9] except for the use of Theorem 4.2 instead of classical Poincaré-Wirtinger inequality ((4.12) in [9]), since we do not need to force vanishing of least median of v . \square

Theorem 4.4. (Compactness and lower semicontinuity for GSBV² functions vanishing in a set of full measure)

Assume $B_r(\mathbf{x}) \subset \mathbb{R}^2$, $u_h \in GSBV^2(B_r(\mathbf{x}))$, $0 < \vartheta \leq 1$

$$(4.13) \quad |\{\mathbf{y} \in B_r(\mathbf{x}) : u_h(\mathbf{y}) = 0\}| / |B_r(\mathbf{x})| \geq \vartheta,$$

$$(4.14) \quad \sup_h \int_{B_r(\mathbf{x})} |\nabla^2 u_h|^2 dy < +\infty$$

and

$$(4.15) \quad \lim_h L_h = 0, \quad \text{where } L_h = \mathcal{H}^1(S_{u_h} \cup S_{\nabla u_h}).$$

Then there are a positive constant c (dependent on the left-hand side of (4.14)), $u_\infty \in H^2(B_r(\mathbf{x}))$ and a sequence $z_h \in GSBV^2(B_r(\mathbf{x}))$ (whose explicit construction is given by (4.23)-(4.28)) s.t., up to a finite number of indices,

$$(4.16) \quad |\{z_h \neq u_h\}| \leq c L_h^2$$

$$(4.17) \quad P(\{z_h \neq u_h\}, B_r(\mathbf{x})) \leq c L_h$$

and there is a subsequence z_{h_k} such that

$$(4.18) \quad \lim_k z_{h_k} = u_\infty \quad \text{strongly in } L^p(B_r(\mathbf{x})), \quad \forall p \geq 1,$$

$$(4.19) \quad \lim_k \nabla z_{h_k} = Du_\infty \quad \text{strongly in } L^p(B_r(\mathbf{x})), \quad \forall p \geq 1,$$

$$(4.20) \quad \int_{B_r(\mathbf{x})} |D^2 u_\infty|^2 d\mathbf{y} \leq \liminf_k \int_{B_r(\mathbf{x})} |\nabla^2 z_{h_k}|^2 d\mathbf{y} \leq \liminf_k \int_{B_r(\mathbf{x})} |\nabla^2 u_{h_k}|^2 d\mathbf{y},$$

$$(4.21) \quad \lim_k u_{h_k} = u_\infty \quad \text{a.e. in } B_r(\mathbf{x}),$$

$$(4.22) \quad \lim_k \nabla u_{h_k} = Du_\infty \quad \text{a.e. in } B_r(\mathbf{x}).$$

Proof. Identical to the proof of Theorem 4.3 in [9], except for the fact that we can avoid forcing least median of u_h and ∇u_h to vanish since we can use Theorem 4.3 for functions vanishing in a sector instead of *GSBV* Poincaré-Wirtinger inequality given by Theorem 4.1 in [9].

For reader convenience we recall the explicit construction of the sequence z_h : by setting $a_h = 4\gamma_2^2 L_h^2$ we have $a_h \leq |B_r|/2$ for large h . Hence there is c dependent on the left-hand side of (4.14) and there are $\eta_h^k \in (0, 1)$, $h \in \mathbb{N}$, $k = 1, 2$, s.t.

$$(4.23) \quad |\{T(\nabla_k u_h, a_h, \eta_h^k) \neq \nabla_k u_h\}| \leq c L_h^2$$

$$(4.24) \quad P(\{T(\nabla_k u_h, a_h, \eta_h^k) \neq \nabla_k u_h\}, B_r) \leq c(L_h + \mathcal{H}^1(S_{\nabla_k u_h}))$$

Referring to definition (4.1) of truncating operator T , we set

$$(4.25) \quad E_h = \bigcup_{k=1,2} \{\mathbf{y} \in B_r : T(\nabla_k u_h, a_h, \eta_h^k) \neq \nabla_k u_h\}$$

$$(4.26) \quad \xi_h = u_h \chi_{B_r \setminus E_h}$$

$$(4.27) \quad b_h = 4K_\vartheta^2 (\mathcal{H}^1(S_{\xi_h} \cup S_{\nabla \xi_h}))^2 \leq \frac{1}{2} |B_r|$$

$$(4.28) \quad z_h = T(\xi_h, b_h, \eta_h)$$

□

5. HESSIAN DECAY FOR BI-HARMONIC FUNCTIONS IN HALF DISK

In this Section we prove that any function which is bi-harmonic in a half-disk and vanishes together with its normal derivative on the diameter has a suitable decay of hessian L^2 -norm.

Theorem 5.1. (*L^2 -hessian decay for bi-harmonic functions in half-disk which vanish together with normal derivative along diameter*)

Set $B_1^+ = B_1(\mathbf{0}) \cap \{(x, y) \in \mathbb{R}^2 : y > 0\} \subset \mathbb{R}^2$.

Assume $z \in H^2(B_1^+)$, $\Delta^2 z = 0$ on B_1^+ , $z = \partial z / \partial y = 0$ on $B_1(\mathbf{0}) \cap \{(x, y) \in \mathbb{R}^2 : y = 0\}$.

Then

$$(5.1) \quad \|D^2 z\|_{L^2(B_\vartheta^+)}^2 \leq \varrho^2 \|D^2 z\|_{L^2(B_1^+)}^2.$$

Moreover there exists a unique extension Z of z in whole B_1 such that $\Delta^2 Z \equiv 0$ and both z, Z have the following expansion in polar coordinates, which is strongly convergent in $L^2(B_1)$ and strongly convergent in $H^2(B_1^+)$:

$$(5.2) \quad Z(x, y) = \sum_{k=0}^{\infty} (a_k \cos(k\vartheta) + b_k \sin(k\vartheta) + (\alpha_k \cos(k\vartheta) + \beta_k \sin(k\vartheta)) r^2) r^k .$$

Proof. Since z belongs to $H^2(B_1^+)$, is bi-harmonic in B_1^+ and $z = \partial z / \partial y = 0$ on $B_1(\mathbf{0}) \cap \{y = 0\}$, then z solves a boundary value problem in B_1^+ for the bilaplacian operator with homogeneous Dirichlet boundary conditions on the diameter; hence regularity properties at a flat portion of the boundary (see [25], Chap.7) entail $z \in C^1(B_1^+(\mathbf{0}) \cup (B_1(\mathbf{0}) \cap \{y = 0\}))$. So the classical Duffin formula [21] holds true: z has a bi-harmonic extension Z in B_1 defined by $Z(x, y) = z(x, y)$ in B_1^+ and by

$$Z(x, -y) = -z(x, y) + 2yz_y(x, y) - y^2 \Delta z(x, y), \quad \forall (x, y) \in B_1^+ .$$

Function Z belongs to $L^2(B_1)$ by construction. Z is bi-harmonic in B_1 , hence by Almansi decomposition [1], there exist two harmonic functions ψ, φ in $L^2(B_1)$ such that $Z = \psi + (x^2 + y^2)\varphi$: in polar co-ordinates,

$$(5.3) \quad \varphi(r, \vartheta) = \frac{1}{4r} \int_0^r \Delta z(\varrho, \vartheta) d\varrho, \quad \psi = z - r^2 \varphi .$$

Hence Z can be represented, with suitable coefficients, by the expansion (5.2) which is strongly convergent in $L^2(B_1)$ and hence in $H^2(B_\varrho)$ for all $\varrho < 1$.

Notice that only suitable combinations of terms in expansion (5.2), say

$$(5.4) \quad \begin{cases} v_k = r^{k+1} \left(\sin((k-1)\vartheta) - \frac{k-1}{k+1} \sin((k+1)\vartheta) \right), & k = 2, 3, 4, \dots \\ \omega_k = r^{k+1} \left(\cos((k-1)\vartheta) - \cos((k+1)\vartheta) \right), & k = -1 \text{ and } 1, 2, 3, \dots \end{cases}$$

fulfill also conditions on diameter, nevertheless we disregard this complicate relationship on coefficients (though it is implicitly understood) which is useless in the following since system (5.4) is strongly entangled and far from providing orthogonal basis either in $H^2(B_r^+)$ or in $L^2(B_r^+)$.

By denoting f_k the k -th term of the expansion (5.2), we compute the second derivatives of f_0 and f_1 :

$$\begin{aligned} D_{xx}^2 f_0 &= 2\alpha_0, & D_{xy}^2 f_0 &= 0 & D_{yy}^2 f_0 &= 2\alpha_0, \\ D_{xx}^2 f_1 &= 2r(3\alpha_1 \cos(\vartheta) + \beta_1 \sin(\vartheta)), & D_{xy}^2 f_1 &= 2r(\beta_1 \cos(\vartheta) + \alpha_1 \sin(\vartheta)), \\ D_{yy}^2 f_1 &= 2r(\alpha_1 \cos(\vartheta) + 3\beta_1 \sin(\vartheta)), \end{aligned}$$

then we compute the second derivatives of f_k , with $k \geq 2$:

$$\begin{aligned} D_{xx}^2 f_k &= r^{k-2} (k(a_k(k-1) + \alpha_k(k+1)r^2) \cos((k-2)\vartheta) + 2\alpha_k(k+1)r^2 \cos(k\vartheta) \\ &\quad + k(b_k(k-1) + \beta_k(k+1)r^2) \sin((k-2)\vartheta) + 2\beta_k(k+1)r^2 \sin(k\vartheta)), \\ D_{xy}^2 f_k &= kr^{k-2} ((b_k(k-1) + \beta_k(k+1)r^2) \cos((k-2)\vartheta) \\ &\quad - (a_k(k-1) + \alpha_k(k+1)r^2) \sin((k-2)\vartheta)), \\ D_{yy}^2 f_k &= r^{k-2} (- (k(a_k(k-1) + \alpha_k(k+1)r^2) \cos((k-2)\vartheta) + 2\alpha_k(k+1)r^2 \cos(k\vartheta) \\ &\quad - k(b_k(k-1) + \beta_k(k+1)r^2) \sin((k-2)\vartheta) + 2\beta_k(k+1)r^2 \sin(k\vartheta)). \end{aligned}$$

Hence for suitable coefficients $c_k = c_k^{i,j}, d_k = d_k^{i,j}, \gamma_k = \gamma_k^{i,j}, \delta_k = \delta_k^{i,j}$, any second derivative of z has the following strongly $L^2(B_\varrho^+)$ convergent expansion, for every $\varrho < 1$ and $i, j = 1, 2$:

$$(5.5) \quad D_{ij}^2 z = \sum_{k=0}^{\infty} (c_k \cos(k\vartheta) + d_k \sin(k\vartheta) + (\gamma_k \cos(k\vartheta) + \delta_k \sin(k\vartheta)) r^2) r^k,$$

Due to strong convergence, we can select partial sums in (5.5) as follows, by splitting terms with different arguments in trigonometric functions,

$$(5.6) \quad \begin{aligned} D_{ij}^2 z &= c_0 + \gamma_0 r^2 \\ &+ (c_1 \cos(\vartheta) + d_1 \sin(\vartheta)) r + (\gamma_1 \cos(\vartheta) + \delta_1 \sin(\vartheta)) r^3 \\ &+ (c_2 \cos(2\vartheta) + d_2 \sin(2\vartheta)) r^2 + (\gamma_2 \cos(2\vartheta) + \delta_2 \sin(2\vartheta)) r^4 \\ &+ (c_3 \cos(3\vartheta) + d_3 \sin(3\vartheta)) r^3 + (\gamma_3 \cos(3\vartheta) + \delta_3 \sin(3\vartheta)) r^5 \\ &+ (c_4 \cos(4\vartheta) + d_4 \sin(4\vartheta)) r^4 + (\gamma_4 \cos(4\vartheta) + \delta_4 \sin(4\vartheta)) r^6 \\ &+ \dots \end{aligned}$$

Since the system $\{\cos(2k\vartheta), \sin(2k\vartheta)\}_{k \in \mathbb{N}}$ is an orthogonal complete system in $L^2(0, \pi)$ we have that odd lines in (5.6) are mutually orthogonal also in $L^2(B_r^+)$ and we can expand all the trigonometric functions with odd multiple of ϑ with respect to this system, in such a way that even lines will be absorbed by odd ones. This is carefully performed by suppressing even lines (the ones where $(2k+1)\vartheta$ appears) in (5.6) one at a time and taking into account of $L^2(B_r^+)$ orthogonal splitting $L^2(B_r^+) = V \oplus V^\perp$, where V is the space

$$V \stackrel{\text{def}}{=} \text{span} \{1, r, r^2, r^{2k+1}, k = 1, 2, \dots\}.$$

At first the $L^2(0, \pi)$ convergent expansions

$$\cos(\vartheta) = \sum_{n=1}^{\infty} \xi_n^1 \sin(2n\vartheta), \quad \sin(\vartheta) = \phi_0^1 + \sum_{n=1}^{\infty} \phi_n^1 \cos(2n\vartheta)$$

allow to cancel second line (related to ϑ) in (5.6) by allocating all terms with trigonometric functions evaluated at $2n\vartheta$ on $(2n+1)$ -th line and, taking into account r powers and convergence properties, writing

$$D_{ij}^2 z = S_1 + \Sigma_1, \quad \text{with } S_1 \in V, \quad \text{and } \Sigma_1 \text{ with empty first and second line of (5.6) :}$$

$$S_1 = c_0 + \gamma_0 r^2 + d_1 \phi_0^1 r + \delta_1 \phi_0^1 r^3.$$

Then the $L^2(0, \pi)$ convergent expansions

$$\cos(3\vartheta) = \sum_{n=1}^{\infty} \xi_n^3 \sin(2n\vartheta), \quad \sin(3\vartheta) = \phi_0^3 + \sum_{n=1}^{\infty} \phi_n^3 \cos(2n\vartheta)$$

allow to cancel fourth line (related to 3ϑ) in (5.6) by allocating all terms with trigonometric functions evaluated at $2n\vartheta$ on $(2n+1)$ -th line and, taking into account r powers and convergence properties, writing

$$D_{ij}^2 z = S_2 + \Sigma_2 \quad \text{with } S_2 \in V, \quad \text{and } \Sigma_2 \text{ with empty second and fourth lines of (5.6) :}$$

$$S_2 = S_1 + d_3 \phi_0^3 r^3 + \delta_3 \phi_0^3 r^5.$$

By iteration (after expanding $\sin((2k+1)\vartheta), \cos((2k+1)\vartheta), k = 0, \dots, n-1$, and getting $D_{ij}^2 z = S_{n-1} + \Sigma_{n-1}$), we expand $\sin((2n+1)\vartheta), \cos((2n+1)\vartheta)$, and get

$$D_{ij}^2 z = S_n + \Sigma_n \quad \text{with } S_n \in V, \quad \text{and } \Sigma_n \text{ with empty first } n \text{ odd lines,}$$

where S_n is a finite sum in the space V :

$$S_n = S_{n-1} + d_{2n-1} \phi_0^{2n-1} r^{2n-1} + \delta_{2n-1} \phi_0^{2n-1} r^{2n+1}.$$

Though Σ_n might not belong to V^\perp , by exploiting $L^2(B_r^+)$ convergence in (5.6) we denote by Ξ_n the modified odd lines from the third odd line (say the fifth one of (5.6)) to the n -th odd line, explicitly (referring to the lines position in (5.6)):

$$\begin{aligned} \Xi_1 &= \text{expansion of the second line,} \\ \Sigma_1 &= \Xi_1 + \text{all the lines after the second,} \\ \Xi_2 &= \Xi_1 + \text{third line} + \text{expansion of the fourth line,} \\ \Sigma_2 &= \Xi_2 + \text{all the lines after the fourth,} \\ \Xi_n &= \Xi_{n-1} + (2n-1)\text{-th line} + \text{expansion of the } (2n)\text{-th line,} \\ \Sigma_n &= \Xi_n + \text{all the lines after the } (2n)\text{-th line.} \end{aligned}$$

Then

$$(5.7) \quad D_{ij}^2 z = S_n + \Xi_n + \varepsilon_n \quad S_n \in V, \quad \Xi_n \in V^\perp, \quad \varepsilon_n \rightarrow 0 \text{ strongly in } L^2(B_r^+).$$

$$(5.8) \quad S_n \rightarrow S \text{ strongly in } L^2(B_r^+), \quad \Xi_n \rightarrow \Xi \text{ strongly in } L^2(B_r^+).$$

Hence

$$D_{ij}^2 z = \sum_{k=0}^{\infty} \left(\left(\sum_{h=0}^{\infty} (A_{h,k} + B_{h,k} r^2) r^h \right) \cos(2k\vartheta) + \left(\sum_{h=0}^{\infty} (C_{h,k} + D_{h,k} r^2) r^h \right) \sin(2k\vartheta) \right)$$

where the expansion is strongly $L^2(B_\varrho^+)$ convergent, $\forall \varrho \in (0, 1)$.

Since $\int_0^\varrho r dr = \varrho^2/2$ and

$$\|1 = \cos 0\|_{L^2(0,\pi)}^2 = \pi, \quad \|\cos(2n\vartheta)\|_{L^2(0,\pi)}^2 = \|\sin(2n\vartheta)\|_{L^2(0,\pi)}^2 = \pi/2 \quad n = 1, 2, \dots,$$

by setting $\lambda_2 = \lambda_2^{(i,j)} = \frac{\pi}{2} \left(\sum_{h=0}^{\infty} (A_{h,0}^{i,j})^2 + \sum_{h=0}^{\infty} (B_{h,0}^{i,j})^2 \right)$, $\Lambda_2 = \sum_{ij} \lambda_2^{(i,j)}$,

via Plancherel identity in $L^2(B_r^+)$, we get

$$\begin{aligned} \|D_{ij}^2 z\|_{L^2(B_\varrho^+)} &= \int_0^\varrho \left\{ \pi \left| \sum_{h=0}^{\infty} (A_{h,0} + B_{h,0} r^2) r^h \right|^2 + \right. \\ &+ \left. \frac{\pi}{2} \sum_{k=1}^{\infty} \left(\left| \sum_{h=0}^{\infty} (A_{h,k} + B_{h,k} r^2) r^h \right|^2 + \left| \sum_{h=0}^{\infty} (C_{h,k} + D_{h,k} r^2) r^h \right|^2 \right) \right\} r dr = \\ &= \lambda_2 \varrho^2 + \sum_{l=3}^{\infty} \lambda_l \varrho^l \end{aligned}$$

but this power sum with positive coefficients is convergent (so the inner sums do converge) and is estimated (uniformly in $\varrho < 1$) by $\|D_{ij}^2 z\|_{L^2(B_1^+)} < +\infty$; then it is (absolutely) convergent even for $\varrho = 1$, and the sum is estimated in the same way. Then $D^2 z \in L^2(B_1^+)$ and has the same expansion since coefficients $A_{h,k}, B_{h,k}, C_{h,k}, D_{h,k}$ are independent of $\varrho \in (0, 1]$. Moreover $(S_n + \Xi_n)$ converges strongly in $L^2(B_1^+)$ to $D_{ij}^2 z$, together with every reordering of its.

By summarizing the following expansion is strongly $L^2(B_\varrho^+)$ convergent, $\forall \varrho \in (0, 1]$:

$$D_{ij}^2 z = \sum_{k=0}^{\infty} \left(\left(\sum_{h=0}^{\infty} (A_{h,k} + B_{h,k} r^2) r^h \right) \cos(2k\vartheta) + \left(\sum_{h=0}^{\infty} (C_{h,k} + D_{h,k} r^2) r^h \right) \sin(2k\vartheta) \right).$$

So, if $\|D^2z\|_{L^2(B_1^+)} \neq 0$, then

$$\frac{\|D^2z\|_{L^2(B_\varrho^+)}}{\|D^2z\|_{L^2(B_1^+)}} = \left(\Lambda_2 \varrho^2 + \sum_{l=3}^{\infty} \Lambda_l \varrho^l \right) / \left(\Lambda_2 + \sum_{l=3}^{\infty} \Lambda_l \right) \leq \varrho^2.$$

□

Remark 5.2. Since $\lambda_2^{(1,2)} = 0$, by the proof of Theorem 5.1 we get a faster decay of mixed derivative:

$$\|D_{xy}^2z\|_{L^2(B_\varrho^+)}^2 \leq \varrho^3 \|D_{xy}^2z\|_{L^2(B_1^+)}^2.$$

Remark 5.3. No nontrivial harmonic function fulfil assumptions of Theorem 5.1. Precisely any $z \in H^2(B_1^+)$ s.t. $\Delta^2z = 0$ on B_1^+ and $z = \partial z / \partial y = 0$ on $B_1(\mathbf{0}) \cap \{y = 0\}$ satisfies also $\Delta z = 0$ in B_1^+ if and only if $z \equiv 0$.

Nevertheless there are (simple) examples with $\Delta z \neq 0 = \Delta^2z$ on B_1^+ , $z = \partial z / \partial y = 0$ on $B_1(\mathbf{0}) \cap \{y = 0\}$ with non trivial harmonic part in Almansi decomposition: e.g. $z(x, y) = y^2 = (y^2 - x^2)/2 + (x^2 + y^2)/2$.

Remark 5.4. Theorem 5.1 cannot be deduced by Schwarz reflection principle for harmonic functions vanishing on the diameter, since the Almansi decomposition on the half-disk B_1^+ ([16], [1]) may not respect the vanishing value on the diameter: e.g. $\varrho^3(\cos \vartheta - \cos(3\vartheta)) = \varrho^2\varphi + \psi$ where $\varphi = x$, $\psi = 3xy^2 - x^3$ are both harmonic but do not vanish on the diameter $\{y = 0\}$ (see [1] and Theorem 3.2 of [17]).

Remark 5.5. While Schwarz reflection for harmonic functions vanishing on the diameter is bounded by 1 as a linear operator from $H^1(B_1^+)$ to $H^1(B_1^-)$, Duffin extension map for bi-harmonic functions vanishing on the diameter together with normal derivative provides a poor control of $H^2(B_1^-)$ in term of $H^2(B_1^+)$ as shown by the following example: referring to (5.4), if we choose $z = \omega_2 - v_3 + \omega_4 - v_5$ then $\|D^2z\|_{L^2(B_1^-)} \approx 12.5761 \|D^2z\|_{L^2(B_1^+)}$.

This depends on the fact that bi-harmonic extension of z may be either even in y (e.g. $z = y^2$) or odd in y (e.g. $z = r^3(3 \sin \vartheta - \sin(3\vartheta)) = 4y^3$) or a mixing of the two (e.g. $z = \omega_2 - v_3$).

Remark 5.6. Bi-harmonic functions like $\varrho^3(\cos \vartheta - \cos(3\vartheta))$ and, quite surprisingly, also combinations of multi-valued functions like $\varrho^{3/2}(\cos(\vartheta/2) - \cos(3\vartheta/2))$ or like $\varrho^{5/3}(\cos(\vartheta/3) - \cos(5\vartheta/3))$ actually turn out to be ($H^2(B_1^+)$ strongly convergent) infinite sums of kind given by (5.2) above: hence they have single-valued analytic (and bi-harmonic) extension to the whole disk B_ϱ and fulfil decay property (5.1).

Remark 5.7. We remark that, by setting $\varphi_t(\vartheta) = (\sin(t\vartheta) - \sin((t-2)\vartheta))t/(t-2)$, $\psi_\tau(\vartheta) = (\cos(\tau\vartheta) - \cos((\tau-2)\vartheta))$, both $r^t\varphi_t(\vartheta)$, $r^\tau\psi_\tau(\vartheta)$, though built with polydromic functions, do have (unique) bi-harmonic extension to the whole disk $B_\varrho(\mathbf{0})$: in fact $\partial_\vartheta^h \varphi_t(\vartheta)|_{\vartheta=0} = \partial_\vartheta^h \varphi_t(\vartheta)|_{\vartheta=2\pi} \forall h$ (due to 2π periodicity of sin and cos), so that their gluing at 2π is not only continuous but also analytic. The same argument holds true for ψ_τ .

6. BLOW-UP AND DECAY AT BOUNDARY POINTS

In this section we analyze the boundary locally around any point belonging to $\partial\Omega \setminus (T_0 \cup T_1 \cup M)$. At first (Theorems 6.1, 6.2) we perform a blow-up of functionals

\mathcal{F} and \mathcal{E} around the origin under the additional assumption that $\mathbf{0}$ belongs to $\partial\Omega$. Then we exploit this results (by translating and scaling) to estimate the decay of these functionals when evaluated on local minimizers around boundary points (Theorems 6.3, 6.4).

Theorem 6.1. (Blow-up of functional \mathcal{F} at boundary points)

Assume (1.4), (1.5), (1.6), (1.7), (1.8) and:

$\mathbf{0} \in \partial\Omega \setminus (T_0 \cup T_1 \cup M)$, $B_r(\mathbf{0}) \subset \tilde{\Omega}$, $\psi_h \in C^2(-r, r)$ with $\psi_h \rightarrow 0$ in $W^{2,\infty}(-r, +r)$, $\omega_h \in C^2(B_r)$ with $\omega_h \rightarrow \omega_\infty \equiv 0$ in $W^{2,\infty}(B_r(\mathbf{0}))$

$$(6.1) \quad \begin{cases} \psi_h \in C^2(-r, r), & \psi_h(0) = 0, & \psi_h'(0) = 0 & \text{Lip}(\psi_h') \leq 1, \\ B^{\psi_h+} \stackrel{\text{def}}{=} B_r(\mathbf{0}) \cap \{y > \psi_h(x)\}, & B^{\psi_h-} \stackrel{\text{def}}{=} B_r(\mathbf{0}) \cap \{y < \psi_h(x)\}, \\ B_\varrho^\tau = \{\mathbf{x} = (x, y) : |\mathbf{x}| < \varrho, y > \tau\} & \text{for } 0 < \tau < \varrho < r. \end{cases}$$

$\gamma_h \in L^q(\tilde{\Omega}) \cap L_{loc}^{2q}(\tilde{\Omega})$, let α_h, β_h, μ_h , three sequences of positive numbers with $\beta_h \leq \alpha_h$, and let $v_\infty \in H^2(B_r(\mathbf{0}))$ s.t. $v_\infty \equiv 0$ in $B_r^-(\mathbf{0})$. Assume $v_h \in GSBV^2(\tilde{\Omega}) \cap L^q(\tilde{\Omega})$, $v_h = \omega_h$ a.e. in B^{ψ_h-} and

- (i) v_h are Ω local minimizers of $\mathcal{F}_{\gamma_h, \omega_h}(\cdot, \mu_h, \alpha_h, \beta_h, B_r(\mathbf{0}))$,
- (ii) $\lim_h \mathcal{H}^1((S_{v_h} \cup S_{\nabla v_h}) \cap B_r(\mathbf{0})) = 0$,
- (iii) $\exists \lim_h \mathcal{F}_{\gamma_h, \omega_h}(v_h, \mu_h, \alpha_h, \beta_h, \overline{B_\varrho^\tau}) \stackrel{\text{def}}{=} \delta(\varrho, \tau) \leq 1$
for a.e. $\varrho, \tau \in (0, r)$ with $\tau < \varrho$, and set $\delta(\varrho, \tau) = 0$ if $\varrho < \tau$.
- (iv) $\lim_h v_h = v_\infty$ a.e. in $B_r(\mathbf{0})$,
- (v) $\lim_h \mu_h = 0$, $\lim_h \mu_h \|\gamma_h\|_{L^q(B_r(\mathbf{0}))}^q = 0$.

Then, for every $\varrho \in (0, r)$, $\tau \in (0, \varrho)$, v_∞ minimizes the functional

$$(6.2) \quad \int_{B_\varrho^\tau(\mathbf{0})} |D^2 v|^2 \, d\mathbf{x}$$

over $\{v \in H^2(B_r(\mathbf{0})) : v = v_\infty \text{ in } B_r(\mathbf{0}) \setminus B_\varrho^\tau; \text{ in particular } v = 0 \text{ in } \overline{B_r^-(\mathbf{0})}\}$.

Moreover

$$(6.3) \quad \delta(\varrho, \tau) = \int_{B_\varrho^\tau(\mathbf{0})} |D^2 v_\infty|^2 \, d\mathbf{x} \quad \text{for almost all } \varrho, \tau : 0 < \tau < \varrho < r.$$

In particular $\Delta^2 v_\infty = 0$ in $B_r^+(\mathbf{0})$, $v_\infty = 0 = \partial v_\infty / \partial y$ in $B_r(\mathbf{0}) \cap \{y = 0\}$, and $v_\infty \in C^1(B_r(\mathbf{0}))$.

Proof. By convergence assumptions on ψ_h , for any $\kappa \in (0, r/2)$ we can assume $|\psi_h| < \kappa < r/2$ for large h , that is hypo-graph of ψ_h contains a fixed sector (of the disk B_r) where $v_h = \omega_h$, hence $u_h = v_h - \omega_h$ fulfil assumption (4.13) uniformly in h with $\vartheta \geq 1/3$, while Dirichlet datum ω_h is not imposed on the portion of the disk where $y > \kappa$.

By (iv) (v) we get, up to subsequences,

$$\lim_h \mu_h^{1/q} |v_h - \gamma_h| = 0 \quad \text{a.e in } B_r.$$

By (ii) (iii) sequence $u_h = v_h - \omega_h$ fulfils all the assumptions of Theorem 4.4:

$$(6.4) \quad \sup_{\varrho, \tau} \sup_h \int_{B_\varrho^\tau} |\nabla^2 v_h|^2 \, d\mathbf{x} \leq 1.$$

Then we can build a sequence z_h as in (4.23)-(4.28), choose subsequences (without relabeling) $z_h, u_h, v_h = u_h + \omega_h$ and $u_\infty \in H^2$ s.t. (4.16)-(4.20) hold true. Since $\omega_h \rightarrow 0$ and $v_h \rightarrow v_\infty$ a.e., we get $u_h \rightarrow u_\infty = v_\infty$, a.e. By (4.20), $\omega_h \rightarrow 0$ in $W^{2,\infty}$ and by (iii) we obtain, for a.e. $\varrho, \tau, 0 < \tau < \varrho < r$,

$$\begin{aligned} \int_{B_\varrho^\tau} |D^2 v_\infty|^2 d\mathbf{y} &\leq \liminf_h \int_{B_\varrho^\tau} \left(|\nabla^2 u_h|^2 + \mu_h |u_h - \gamma_h|^q \right) d\mathbf{y} \leq \\ &\leq \liminf_h \int_{B_\varrho^\tau} \left(|\nabla^2 v_h|^2 + \mu_h |v_h - \gamma_h|^q \right) d\mathbf{y} \leq \lim_h \mathcal{F}_{\gamma_h \omega_h}(v_h, \overline{B_\varrho^\tau}) = \delta(\varrho, \tau). \end{aligned}$$

To achieve the proof we have to show that for a.e. $\varrho, \tau, 0 < \tau < \varrho < r$, for every $\kappa \in (0, r/2)$, and every $u \in H^2(B_r)$ with $u = v_\infty$ in $B_r^+(\mathbf{0}) \setminus B_\varrho^{2\kappa}$ (hence $u = 0$ in $B_r^-(\mathbf{0})$):

$$(6.5) \quad \int_{B_\varrho^\tau} |D^2 u|^2 d\mathbf{y} \geq \delta(\varrho, \tau).$$

In fact (6.5) implies $\Delta v_\infty = 0, B_\varrho^+$ and $v_\infty = \partial v_\infty / \partial y = 0$ on $B_1 \cap \{y = 0\}$; hence $v_\infty \in C^1(B_r^+ \cup (B_r \cap \{y = 0\}))$ ([25]). We prove the inequality (6.5) for fixed $\kappa \in (0, r/2)$: the convergence property of ψ_h allows to repeat the proof for any such κ by selecting large enough h .

Map δ is monotone non decreasing in ϱ and monotone non increasing in τ , hence:

for any frozen τ , map δ is continuous up to a countable set of values for ϱ ,

for any frozen ϱ , map δ is continuous up to a countable set of values for τ .

For any selection of ϱ, τ s.t. δ is *separately continuous* at ϱ, τ , we get by monotonicity that actually δ is a (two-variables) continuous map at ϱ, τ . This continuity property holds true for a.e. $\varrho, \tau \in (0, r)$.

Assume by contradiction there exist $u \in H^2(B_r), \varepsilon > 0, s, \sigma$, s.t. $2\kappa < \sigma < s < r$, δ is continuous at $\varrho = s, \tau = \sigma$, $u = v_\infty$ in $B_r^+ \setminus B_s^{2\kappa}$ (hence $u = 0$ in B_r^-) and

$$(6.6) \quad \int_{B_s^\sigma} |D^2 u|^2 d\mathbf{y} \leq \delta(s, \sigma) - \varepsilon.$$

From now on we fix η, κ s.t.

$$(6.7) \quad s < \eta < r, \quad 0 < \kappa < \frac{1}{2} \sigma < \frac{1}{2} \sqrt{r^2 - \eta^2}$$

Referring to (4.25), (4.27), we set

$$(6.8) \quad L_h = \mathcal{H}^1((S_{v_h} \cup S_{\nabla v_h}) \cap B_r(\mathbf{0}))$$

$$(6.9) \quad \Xi_h = \{ \mathbf{y} \in B_r : z_h \neq \xi_h \},$$

$$(6.10) \quad A_h = \{ z_h \neq u_h \}.$$

In particular $A_h = E_h \cup \Xi_h$.

In order to get a contradiction we will paste together u_h and z_h along the boundary of a suitably chosen lunula (the energy addition will tend to 0 as $h \rightarrow \infty$ due to (iii), (4.17) and Matching Lemma 3.2), then we will join such new function in a smaller lunula with u (which has less squared hessian energy). Before some preliminary estimates are needed.

We emphasize that

$$(6.11) \quad \mathcal{H}^1(S_{v_h} \cap \partial B_\varrho) = \mathcal{H}^1(S_{\nabla v_h} \cap \partial B_\varrho) = 0 \quad \text{for a.e } \varrho \in (0, r)$$

and by (4.17)

$$(6.12) \quad P(A_h, B_r) \leq c L_h.$$

By integrating first in polar coordinates, then in cartesian coordinates and taking into account isoperimetric inequality we get

$$(6.13) \quad \alpha_h \int_0^r \mathcal{H}^1(A_h \cap \partial B_\varrho) d\varrho = \alpha_h |A_h \cap B_r| \leq \alpha_h (\gamma_2 P(A_h, B_r))^2 \leq c^2 \gamma_2^2 \alpha_h L_h^2$$

$$(6.14) \quad \begin{aligned} & \alpha_h \int_0^{\sqrt{r^2-t^2}} \mathcal{H}^1(A_h \cap \{(x, y) : |x| \leq t\}) dy = \\ & \alpha_h \left| A_h \cap \{|x| \leq t, 0 \leq y \leq \sqrt{r^2-t^2}\} \right| \leq \\ & \leq \alpha_h |A_h \cap B_r| \leq \alpha_h (\gamma_2 P(A_h, B_r))^2 \leq c^2 \gamma_2^2 \alpha_h L_h^2 \end{aligned}$$

since the sequence $\alpha_h L_h$ is bounded by (iii), then (ii) entails

$$\lim_h \alpha_h L_h^2 = 0.$$

Hence, by (6.13), we have, up to subsequence and without relabeling,

$$(6.15) \quad \exists \lim_h \alpha_h \mathcal{H}^1(A_h \cap \partial B_t^+) = 0 \text{ for a.e. } t \in (0, r).$$

By assumption (6.7), the interval $(2\kappa, \sqrt{r^2 - \eta^2})$ is not empty and contains σ . For any choice of $t \in (s, r)$ as above (fulfilling (6.15)) and for a.e. $d \in (2\kappa, \sqrt{r^2 - \eta^2})$ (thanks to (6.14)) we have

$$(6.16) \quad \lim_h \alpha_h \mathcal{H}^1(A_h \cap \{|x| \leq t, y = d\}) = 0;$$

by summarizing, for any t fulfilling (6.15) for a.e. $d \in (2\kappa, \sqrt{r^2 - t^2})$ both (6.15), (6.16) hold true, so that, up to subsequence and without relabeling,

$$(6.17) \quad \exists \lim_h \alpha_h \mathcal{H}^1(A_h \cap \partial B_t^d) = 0 \text{ for a.e. } t, d,$$

and by (6.11), (6.17), (ii) and $\beta_h \leq \alpha_h$ we get,

$$(6.18) \quad \begin{aligned} & \lim_h (\alpha_h \mathcal{H}^1(S_{z_h} \cap \partial B_t^d) + \beta_h \mathcal{H}^1((S_{\nabla z_h} \setminus S_{z_h}) \cap \partial B_t^d)) = 0 \\ & \text{for a.e } t \in (s, \eta) \text{ and a.e } d \in (2\kappa, \sqrt{r^2 - t^2}), \end{aligned}$$

notice that the interval $(2\kappa, \sqrt{r^2 - t^2})$ is not empty since it contains σ due to (6.7). By continuity of δ at $\varrho = s, \tau = \sigma$ and by (6.17), (6.18) and (iii) we can choose $t \in (s, \eta)$ close to s as needed, $d \in (2\kappa, \sigma)$ close to σ as needed (and let them fixed in the following) and $\tilde{h} \in \mathbb{N}$ s.t. $\sigma - d < t - s$ and, setting $M_{t,s}^{d,\sigma} = \overline{B_t^d} \setminus B_s^\sigma$, the following list of inequalities hold true:

$$(6.19) \quad \delta(t, d) - \delta(s, \sigma) < \varepsilon/6,$$

$$(6.20) \quad \alpha_h \mathcal{H}^1(A_h \cap \partial M_{t,s}^{d,\sigma}) \leq \varepsilon/6 \quad h > \tilde{h},$$

$$(6.21) \quad \alpha_h \mathcal{H}^1(S_{z_h} \cap \partial M_{t,s}^{d,\sigma}) + \beta_h \mathcal{H}^1((S_{\nabla z_h} \setminus S_{z_h}) \cap \partial M_{t,s}^{d,\sigma}) \leq \varepsilon/6 \quad h > \tilde{h},$$

$$(6.22) \quad \int_{B_t^d \setminus B_s^\sigma} |D^2 u|^2 d\mathbf{x} < \varepsilon/6,$$

$$(6.23) \quad \mathcal{F}\left(v_h, \overline{B_t^d} \setminus B_s^\sigma\right) \leq 2\varepsilon/6 \quad h > \tilde{h};$$

In fact (6.19) express the continuity of δ at (s, σ) ; feasibility of choices (6.20), (6.21) follows by (6.17), (6.18); inequality (6.22) follows by the absolute continuity of $\int_A |D^2 u|^2 d\mathbf{x}$ with respect to the Lebesgue measure of A , eventually (6.23) follows by

$$\lim_h \mathcal{F}\left(v_h, \overline{B_t^d} \setminus B_s^\sigma\right) = \delta(t, d) - \delta(s, \sigma) + \lim_h \mathcal{F}(v_h, \partial B_s^\sigma)$$

which is estimated by $2\varepsilon/6$ thanks to (6.19), (6.21).

We fix the matching:

$$(6.24) \quad \zeta_h = u_h \chi_{B_r \setminus \overline{B_t^d}} + z_h \chi_{\overline{B_t^d}},$$

hence (6.17) and Lemma 3.2 entail, for a.e. $\varrho \in (0, r)$, $\tau \in (d, \sigma)$,

$$(6.25) \quad \lim_h \mathcal{F}_{\gamma_h \omega_h}(\zeta_h, \mu_h, \alpha_h, \beta_h, B_\varrho^\tau) = \lim_h \mathcal{F}_{\gamma_h \omega_h}(v_h, \mu_h, \alpha_h, \beta_h, B_\varrho^\tau) = \delta(\varrho, \tau) \leq 1.$$

Eventually we perform the joining of $u + \omega_h$ and $\zeta_h + \omega_h$ between lunulae B_t^d and B_s^σ : by referring to Lemma 3.1, we choose $\Psi \equiv 1$ in a neighborhood of B_s^σ and set

$$(6.26) \quad \tau_h = \Psi(u + \omega_h) + (1 - \Psi)(\zeta_h + \omega_h),$$

so that

$$(6.27) \quad \tau_h = u_h + \omega_h = v_h \quad \text{in } B_r \setminus \overline{B_t^d}, \quad \tau_h = u + \omega_h \quad \text{in } \overline{B_s^\sigma},$$

hence

$$(6.28) \quad \mathcal{F}(\tau_h, B_r \setminus \overline{B_t^d}) = \mathcal{F}(v_h, B_r \setminus \overline{B_t^d}).$$

Then by Lemma 3.1 we obtain, for any $\theta > 0$,

$$(6.29) \quad \begin{aligned} \mathcal{F}(\tau_h, \overline{B_t^d}) &\leq (1 + \theta) \left(\mathcal{F}(u + \omega_h, \overline{B_t^d}) + \mathcal{F}(\zeta_h + \omega_h, \overline{B_t^d} \setminus B_s^\sigma) \right) + \\ &+ \frac{c}{(\sigma - d)^2} \left(\int_{B_t^d \setminus B_s^\sigma} |\nabla(u - \zeta_h)|^2 d\mathbf{x} + \frac{c}{\theta d^2 (\sigma - d)^2} \int_{B_t^d \setminus B_s^\sigma} |u - \zeta_h|^2 d\mathbf{x} \right) \end{aligned}$$

By compactness Theorem 4.4, with our choice for d, σ fulfilling $2\kappa < d < \sigma$:

$$(6.30) \quad \lim_h \int_{B_t^d \setminus B_s^\sigma} |\nabla(v_\infty - \zeta_h)|^2 d\mathbf{x} = \lim_h \int_{B_t^d \setminus B_s^\sigma} |v_\infty - \zeta_h|^2 d\mathbf{x} = 0,$$

hence, thanks to $u = v_\infty$ in $B_r^+ \setminus B_s^{2\kappa}$, possibly by extracting subsequences without relabeling and letting $h \rightarrow +\infty$ in (6.29) we obtain

$$(6.31) \quad \lim_h \mathcal{F}(\tau_h, \overline{B_t^d}) \leq (1 + \theta) \left(\lim_h \mathcal{F}(u + \omega_h, \overline{B_t^d}) + \lim_h \mathcal{F}(\zeta_h + \omega_h, \overline{B_t^d} \setminus B_s^\sigma) \right).$$

By convergence $\omega_h \rightarrow 0$ in $W^{2, \infty}(B_r)$ there is $h_0 \geq \tilde{h}$ s.t.

$$(6.32) \quad \left| \int_{B_t^d \setminus B_s^\sigma} |D^2(u + \omega_h)|^2 d\mathbf{x} - \int_{B_t^d \setminus B_s^\sigma} |D^2 u|^2 d\mathbf{x} \right| < \frac{\varepsilon}{6} \quad \text{for } h > h_0.$$




FIGURE 1. Joining along handle $M_{t,s}^{d,\sigma} = \overline{B_t^d} \setminus B_s^\sigma$

By (4.26), $\omega_h \rightarrow 0$ in $W^{2,\infty}$, (6.23), (6.27) we get

$$(6.33) \quad \begin{aligned} \lim_h \mathcal{F} \left(\zeta_h + \omega_h, \overline{B_t^d} \setminus B_s^\sigma \right) &= \\ &= \lim_h \mathcal{F} \left(z_h + \omega_h, \overline{B_t^d} \setminus B_s^\sigma \right) \leq \lim_h \mathcal{F} \left(v_h, \overline{B_t^d} \setminus B_s^\sigma \right) \leq 2 \frac{\varepsilon}{6} \end{aligned}$$

By letting $\vartheta \rightarrow 0$ in (6.31), taking into account (6.20)-(6.24), (6.27), (6.32) and (6.33) we get

$$(6.34) \quad \mathcal{F} \left(\tau_h, \overline{B_t^d} \setminus B_s^\sigma \right) \leq 4\varepsilon/6 \quad h > h_0;$$

By exploiting (6.6), (6.20), (6.21), (6.23), (6.34), $\tau_h = u + \omega_h$ in B_s^σ and eventually δ monotonicity with respect to inclusion of sets, we get the contradiction:

$$(6.35) \quad \begin{aligned} \delta(t,d) &= \lim_h \mathcal{F} \left(v_h, \overline{B_t^d} \right) \stackrel{\text{minimality of } v_h}{\leq} \lim_h \mathcal{F} \left(\tau_h, \overline{B_t^d} \right) = \\ &= \lim_h \mathcal{F} \left(\tau_h, \overline{B_s^\sigma} \right) + \lim_h \mathcal{F} \left(\tau_h, \overline{B_t^d} \setminus B_s^\sigma \right) - \lim_h \mathcal{F} \left(\tau_h, \partial B_s^\sigma \right) \stackrel{(6.34)}{\leq} \\ &\leq \lim_h \mathcal{F} \left(\tau_h, \overline{B_s^\sigma} \right) + 4 \frac{\varepsilon}{6} = \\ &= \lim_h \mathcal{F} \left(u + \omega_h, \overline{B_s^\sigma} \right) + 4 \frac{\varepsilon}{6} = \lim_h \int_{B_s^\sigma} |D^2(u + \omega_h)|^2 dx + 4 \frac{\varepsilon}{6} \stackrel{(6.32)}{\leq} \\ &\leq \int_{B_s^\sigma} |D^2 u|^2 dx + 5 \frac{\varepsilon}{6} \leq \delta(s,\sigma) - \varepsilon + 5 \frac{\varepsilon}{6} \leq \delta(t,d) - \frac{\varepsilon}{6}. \end{aligned}$$

□

Theorem 6.2. (Blow-up of functional \mathcal{E} at boundary points)

Assume (1.4), (1.5), (1.6), (1.7), (1.8), (1.9), (1.10), (1.11), $\mathbf{0} \in \partial\Omega \setminus (T_0 \cup T_1 \cup M)$ and $B_r(\mathbf{0}) \subset \tilde{\Omega}$, let α_h, β_h , two sequences of positive numbers with $\beta_h \leq \alpha_h$,

$\psi_h \in C^2(-r, r)$ with $\psi_h \rightarrow 0$ in $W^{2,\infty}(-r, +r)$, $\omega_h \in C^2(B_r(\mathbf{0}))$ with $\omega_h \rightarrow \omega_\infty \equiv 0$ in $W^{2,\infty}$ and let $v_\infty \in H^2(B_r(\mathbf{0}))$ s.t. $v_\infty \equiv 0$ in $B_r^-(\mathbf{0})$. Assume (6.1), $v_h \in GSBV^2(\tilde{\Omega})$, $v_h = \omega_h$ a.e. in $B^{\psi_h^-}$ and

- (i) v_h are Ω local minimizers of $\mathcal{E}_{\omega_h}(\cdot, \alpha_h, \beta_h, B_r(\mathbf{0}))$,
- (ii) $\lim_h \mathcal{H}^1((S_{v_h} \cup S_{\nabla v_h}) \cap B_r(\mathbf{0})) = 0$,
- (iii) $\exists \lim_h \mathcal{E}_{\omega_h}(v_h, \alpha_h, \beta_h, \overline{B_\varrho}(\mathbf{0})) \stackrel{\text{def}}{=} \delta(\varrho, \tau) \leq 1$
for a.e. $\varrho, \tau \in (0, r)$ with $\tau < \varrho$, and set $\delta(\varrho, \tau) = 0$ if $\varrho < \tau$.
- (iv) $\lim_h v_h = v_\infty$ a.e. in $B_r(\mathbf{0})$.

Then, for every $\varrho, \tau : 0 < \tau < \varrho < r$, v_∞ minimizes the functional

$$(6.36) \quad \int_{B_r^+(\mathbf{0})} |D^2 u|^2 dx$$

over $\{u \in H^2(B_\varrho(\mathbf{0})) : u = 0 \text{ in } B_r^-(\mathbf{0}); u = v_\infty \text{ in } B_r(\mathbf{0}) \setminus B_\varrho^+(\mathbf{0})\}$.

Moreover

$$(6.37) \quad \delta(\varrho, \tau) = \int_{B_\varrho^+(\mathbf{0})} |D^2 v_\infty|^2 dx \quad \text{for almost all } \tau, \varrho : 0 < \tau < \varrho < r.$$

In particular $\Delta^2 v_\infty = 0$ in $B_r^+(\mathbf{0})$, $v_\infty = 0 = \partial v_\infty / \partial y$ in $B_r(\mathbf{0}) \cap \{y = 0\}$.

Proof. Repeat the proof of the previous Theorem with $\mu_h = 0$. \square

Due to (1.5), (1.6) for any sequence of points $\mathbf{x}_h \in \partial\Omega \setminus (T_0 \cup T_1 \cup M)$ possibly after suitable rotations of coordinates around each $\mathbf{x}_h = (x_h, y_h)$, we can find ϱ_h and φ_h s.t., by setting

$$(6.38) \quad \Omega_{\varphi_h^+} = \Omega \cap B_{\varrho_h}(\mathbf{x}_h), \quad \Omega_{\varphi_h^-} = B_{\varrho_h}(\mathbf{x}_h) \setminus \overline{\Omega},$$

we have

$$(6.39) \quad \begin{cases} w \in C^2(B_{\varrho_h}(\mathbf{x}_h)), & \Omega_{\varphi_h^+} = B_{\varrho_h}(\mathbf{x}_h) \cap \{y > \varphi_h(x)\}, \\ \varphi_h \in C^2(x_h - \varrho_h, x_h + \varrho_h), & \varphi_h(x_h) = y_h, \varphi_h'(x_h) = 0, \text{Lip}(\varphi_h') \leq C. \end{cases}$$

Referring to (6.38), (6.39) we re-scale and translate at the origin sets Ω_{φ_h} and choose the graphs ψ_h to be used in the application of blow-up Theorem (with $r = 1$) as follows:

$$(6.40) \quad \psi_h(x) = \varrho_h^{-1}(\varphi_h(x_h + \varrho_h x) - y_h)$$

$$(6.41) \quad B^{\psi_h^+} = B_1(\mathbf{0}) \cap \{(x, y) : y_h + \varrho_h y > \varphi_h(x_h + \varrho_h x)\} = B_1(\mathbf{0}) \cap \{y > \psi_h(x)\}$$

$$(6.42) \quad B^{\psi_h^-} = B_1(\mathbf{0}) \cap \{(x, y) : y_h + \varrho_h y < \varphi_h(x_h + \varrho_h x)\} = B_1(\mathbf{0}) \cap \{y < \psi_h(x)\};$$

we get

$$(6.43) \quad \left\{ \begin{array}{l} B^{\psi_h^\pm} = (\Omega_{\varphi_h^\pm} - \mathbf{x}_h) / \varrho_h \\ \psi_h(0) = 0, \psi_h'(0) = 0 \\ \psi_h'(x) = \varphi_h'(x_h + \varrho_h x) = \varrho_h x \varphi_h''(x_h) + o(\varrho_h) \\ \psi_h''(x) = \varrho_h \varphi_h''(x_h + \varrho_h x) \\ \text{Lip}(\psi_h) = \text{Lip}(\varphi_h), \quad \text{Lip}(\psi_h') = \varrho_h \text{Lip}(\varphi_h'). \end{array} \right.$$

Theorem 6.3. (Decay of functional \mathcal{F} at boundary points) Assume (1.3), (1.4), (1.5), (1.6), (1.7), (1.8), (1.9), (1.10), (1.17). Then, by referring to (2.13) and to (2.14) about the meaning of $\bar{\varrho}$ and c_0 ,

$$(6.44) \quad \forall k > 2, \forall \eta, \sigma \in (0, 1), \quad \exists \varepsilon_0 > 0, \exists \vartheta_0 > 0 \quad \text{such that}$$

for all $\varepsilon \in (0, \varepsilon_0]$, for any $\mathbf{x} \in \partial\Omega \setminus (T_0 \cup T_1 \cup M)$, for any u which is an $\overline{\Omega_{\varphi+}}$ local minimizer of $\mathcal{F}_{g w}(\cdot, \mu, \alpha, \beta, \overline{\Omega_{\varphi+}})$, for any ϱ s.t. $B_{\varrho}(\mathbf{x}) \subset \tilde{\Omega}$ (we can assume without restriction (6.39)), $0 < \varrho \leq (\varepsilon^k \wedge \bar{\varrho} \wedge (c_0 \vee 1)^{-1})$, $\int_{B_{\varrho}(\mathbf{x})} |g|^{2q} \leq \varepsilon^k$ and

$$(6.45) \quad \alpha \mathcal{H}^1(S_u \cap \overline{\Omega_{\varphi+}}) + \beta \mathcal{H}^1((S_{\nabla u} \setminus S_u) \cap \overline{\Omega_{\varphi+}}) < \varepsilon \varrho,$$

we have

$$(6.46) \quad \mathcal{F}_{g w}(u, B_{\eta \varrho}(\mathbf{x})) \leq \eta^{2-\sigma} \max \left\{ \mathcal{F}_{g w}(u, B_{\varrho}(\mathbf{x})), \varrho^2 \vartheta_0 \left((\text{Lip}(\varphi'))^2 + (\text{Lip}(Dw))^2 \right) \right\}.$$

Proof. Assume the Theorem is false. Then there are $k > 2$, $\eta, \sigma \in (0, 1)$; three sequences $\varrho_h, \varepsilon_h, \vartheta_h$ s.t. $0 < \varrho_h \leq (\bar{\varrho} \wedge (c_0 \vee 1)^{-1})$, $\varepsilon_h > 0$, $\vartheta_h > 0$, $\varepsilon_1 = 1$, $\varepsilon_h \downarrow 0$, $\lim_h \vartheta_h = +\infty$; a sequence $\mathbf{x}_h \in \partial\Omega \setminus (T_0 \cup T_1 \cup M)$; a sequence $w_h \in C^2(B_{\varrho_h}(\mathbf{x}_h))$ s.t. $|w_h| \leq C$, $\text{Lip}(Dw_h) \leq C$; a sequence $\varphi_h \in C^2((x_h - \varrho_h, x_h + \varrho_h))$ with $\varphi_h(x_h) = y_h$, $\varphi_h'(x_h) = 0$, $\text{Lip}(\varphi_h') \leq C$ and

$$\Omega_{\varphi_h+} = \Omega \cap B_{\varrho_h}(\mathbf{x}_h) \cap \{y > \varphi_h(x)\};$$

a sequence $u_h \in X(\tilde{\Omega})$ of Ω local minimizers of $\mathcal{F}_{g w_h}(\cdot, \mu, \alpha, \beta, B_{\varrho_h}(\mathbf{x}_h))$ among v s.t. $v = w_h$ on Ω_{φ_h-} ;

$$(6.47) \quad \varrho_h \leq \varepsilon_h^k, \quad \int_{B_{\varrho_h}(\mathbf{x}_h)} |g|^{2q} \leq \varepsilon_h^k;$$

$$(6.48) \quad \alpha \mathcal{H}^1(S_{u_h} \cap \overline{\Omega_{\varphi_h+}}) + \beta \mathcal{H}^1((S_{\nabla u_h} \setminus S_{u_h}) \cap \overline{\Omega_{\varphi_h+}}) < \varepsilon_h \varrho_h$$

and

$$(6.49) \quad \mathcal{F}_{g w_h}(u_h, \mu, \alpha, \beta, B_{\eta \varrho_h}(\mathbf{x}_h)) > \eta^{2-\sigma} \max \left\{ \mathcal{F}_{g w_h}(u_h, \mu, \alpha, \beta, B_{\varrho_h}(\mathbf{x}_h)), \varrho_h^2 \vartheta_h \left((\text{Lip}(\varphi_h'))^2 + (\text{Lip}(Dw_h))^2 \right) \right\}.$$

By translating \mathbf{x}_h to $\mathbf{0}$, re-scaling and applying a common affine linear correction to data and local minimizers, we set, for $\mathbf{y} \in B_1(\mathbf{0})$:

$$(6.50) \quad \omega_h(\mathbf{y}) = (\lambda_h \varrho_h^3)^{-1/2} \left(w_h(\mathbf{x}_h + \varrho_h \mathbf{y}) - \varrho_h Dw_h(\mathbf{x}_h) \cdot \mathbf{y} - w_h(\mathbf{x}_h) \right)$$

$$(6.51) \quad \gamma_h(\mathbf{y}) = (\lambda_h \varrho_h^3)^{-1/2} \left(g(\mathbf{x}_h + \varrho_h \mathbf{y}) - \varrho_h Dw_h(\mathbf{x}_h) \cdot \mathbf{y} - w_h(\mathbf{x}_h) \right)$$

$$(6.52) \quad v_h(\mathbf{y}) = (\lambda_h \varrho_h^3)^{-1/2} \left(u_h(\mathbf{x}_h + \varrho_h \mathbf{y}) - \varrho_h Dw_h(\mathbf{x}_h) \cdot \mathbf{y} - w_h(\mathbf{x}_h) \right)$$

where

$$(6.53) \quad \lambda_h = (\varrho_h^{-1} \mathcal{F}_{g w_h}(u_h, \mu, \alpha, \beta, B_{\varrho_h}(\mathbf{x}_h))) \vee \varepsilon_h.$$

Notice that, due to density upper bound in Theorem 2.10 and $\varrho_h \leq \bar{\varrho} \wedge (c_0 \vee 1)^{-1}$,

$$\lambda_h \leq c_0 \vee 1 < +\infty \quad \text{and} \quad \lambda_h \varrho_h \leq 1 \quad \forall h$$

and functions v_h and ω_h coincide on $B^{\psi_h^-} = B_1(\mathbf{0}) \cap \{y < \psi_h(x)\}$. Moreover, by uniform C^2 property of w_h , and applying Lagrange Theorem to each component of $D\omega_h$

$\forall \mathbf{y} \in B_1$, $i, j = 1, 2 \exists \tilde{t}(i) \in (0, 1)$ s.t. by setting $\tilde{\mathbf{y}}(i) = \tilde{t}(i)\mathbf{y}$, we get

$$D_j \omega_h(\mathbf{y}) = D_j \omega_h(\mathbf{0}) + \sum_i D_{ij} \omega_h(\tilde{\mathbf{y}}(i)) \mathbf{y}_i$$

say, by denoting $\widetilde{D^2 \omega_h}(\tilde{\mathbf{y}})$ the hessian of ω_h with i -th row evaluated at $\tilde{\mathbf{y}}(i)$,

$$D\omega_h(\mathbf{y}) = D\omega_h(\mathbf{0}) + \widetilde{D^2 \omega_h}(\tilde{\mathbf{y}}) \cdot \mathbf{y}$$

then

$$(6.54) \quad \left\{ \begin{array}{l} \omega_h(\mathbf{0}) = 0, \quad D\omega_h(\mathbf{0}) = \mathbf{0}, \\ D\omega_h(\mathbf{y}) = \widetilde{D^2 \omega_h}(\tilde{\mathbf{y}}) \cdot \mathbf{y} \\ D\omega_h(\mathbf{y}) = (\lambda_h \varrho_h)^{-1/2} \left(Dw_h(\mathbf{x}_h + \varrho_h \mathbf{y}) - Dw_h(\mathbf{x}_h) \right) \\ D^2 \omega_h(\mathbf{y}) = (\varrho_h / \lambda_h)^{1/2} D^2 w_h(\mathbf{x}_h + \varrho_h \mathbf{y}) \\ |D^2 \omega_h(\mathbf{y})| \leq (\varrho_h / \lambda_h)^{1/2} \text{Lip}(Dw_h) \\ |D\omega_h(\mathbf{y})| \leq |\widetilde{D^2 \omega_h}(\tilde{\mathbf{y}})| |\mathbf{y}| \leq (\varrho_h / \lambda_h)^{1/2} \text{Lip}(Dw_h) \\ \text{Lip}(D\omega_h) = (\varrho_h / \lambda_h)^{1/2} \text{Lip}(Dw_h) \end{array} \right.$$

hence

$$(6.55) \quad |D\omega_h(\mathbf{y})| \leq C \varepsilon_h^{(k-1)/2}, \quad |D^2 \omega_h(\mathbf{y})| \leq C \varepsilon_h^{(k-1)/2}.$$

Due to (1.6) φ_h are uniformly C^2 , hence (6.43) entails $\psi_h \rightarrow 0$ in $W^{2,\infty}(-1, 1)$.

Estimates (6.55) entail strong $W^{2,\infty}(B_1)$ convergence of ω_h to $\omega_\infty \equiv 0$.

Due to Remark 2.9 functions v_h are Ω local minimizers of $\mathcal{F}_{\gamma_h, \omega_h}(\cdot, \mu_h, \alpha_h, \beta_h, B_1(\mathbf{0}))$ among v with $v = \omega_h$ in $B^{\psi_h^-}$ where

$$(6.56) \quad \alpha_h = \frac{\alpha}{\lambda_h}, \quad \beta_h = \frac{\beta}{\lambda_h}, \quad \mu_h = \mu \lambda_h^{\frac{q}{2}-1} \varrho_h^{1+\frac{3}{2}q}.$$

By scaling Lemma 2.5, (6.49), last identity in (6.43), (6.54), and $\lambda_h \varrho_h \leq 1$ we have

$$(6.57) \quad \begin{cases} \mathcal{F}_{g w_h}(u_h, \mu, \alpha, \beta, B_{\varrho_h}(\mathbf{x}_h)) = \lambda_h \varrho_h \mathcal{F}_{\gamma_h \omega_h}(v_h, \mu_h, \alpha_h, \beta_h, B_1(\mathbf{0})) \\ \mathcal{F}_{g w_h}(u_h, \mu, \alpha, \beta, B_{\eta \varrho_h}(\mathbf{x}_h)) = \lambda_h \varrho_h \mathcal{F}_{\gamma_h \omega_h}(v_h, \mu_h, \alpha_h, \beta_h, B_\eta(\mathbf{0})) \end{cases}$$

and

$$(6.58) \quad \begin{aligned} \mathcal{F}_{\gamma_h \omega_h}(v_h, \mu_h, \alpha_h, \beta_h, B_\eta(\mathbf{0})) &= \mathcal{F}_{g w_h}(u_h, \mu, \alpha, \beta, B_{\eta \varrho_h}(\mathbf{x}_h)) / (\lambda_h \varrho_h) \\ &> \eta^{2-\sigma} \frac{\varrho_h}{\lambda_h} \vartheta_h \left(\left(\frac{\text{Lip}(\psi_h')}{\varrho_h} \right)^2 + \left(\frac{\text{Lip}(D\omega_h)}{\sqrt{\varrho_h / \lambda_h}} \right)^2 \right) \\ &= \eta^{2-\sigma} \frac{\vartheta_h}{\lambda_h} \left(\frac{(\text{Lip}(\psi_h'))^2}{\varrho_h} + \lambda_h (\text{Lip}(D\omega_h))^2 \right) \\ &\geq \eta^{2-\sigma} \vartheta_h \left((\text{Lip}(\psi_h'))^2 + (\text{Lip}(D\omega_h))^2 \right) \end{aligned}$$

so that by (6.53), (6.57)

$$(6.59) \quad \mathcal{F}_{\gamma_h \omega_h}(v_h, \mu_h, \alpha_h, \beta_h, B_1(\mathbf{0})) \leq 1,$$

$$(6.60) \quad \alpha_h \mathcal{H}^1(S_{v_h} \cap \overline{B^{\psi_h+}}) + \beta_h \mathcal{H}^1((S_{\nabla v_h} \setminus S_{v_h}) \cap \overline{B^{\psi_h+}}) < \varepsilon_h$$

and (6.49), (6.57), (6.58) entail

$$(6.61) \quad \mathcal{F}_{\gamma_h \omega_h}(v_h, \mu_h, \alpha_h, \beta_h, B_\eta(\mathbf{0})) > \eta^{2-\sigma} \mathcal{F}_{\gamma_h \omega_h}(v_h, \mu_h, \alpha_h, \beta_h, B_1(\mathbf{0})).$$

By (6.59) and Theorem 4.4, up to subsequences and without relabeling,

$$(6.62) \quad \exists v_\infty \in H^2(B_1) : \quad \lim_h v_h = v_\infty \quad \text{a.e. in } B_1$$

say hypothesis (iv) of Theorem 6.1, which we want to apply to handle v_h and v_∞ .

We proceed by checking the other assumptions of the Theorem 6.1.

Since u_h is an Ω local minimizer of $\mathcal{F}_{g \omega_h}$ then v_h is a $\varrho_h^{-1}(\Omega - \mathbf{x}_h)$ local minimizer of $\mathcal{F}_{\gamma_h \omega_h}$, that is (i) holds true; (6.60) entails (ii); we must verify (iii), (v) and the structural assumptions.

Now we prove (iii): choose a dense (in (0,1)) sequence of radii $\varrho_j = \tau_j$. Thanks to (6.59), for any pair $j, l \in \mathbb{N}$ such that $0 < \tau_l < \varrho_j < 1$ we can extract a subsequence of v_h and then diagonalize (without relabeling) in such a way that

$$\exists \text{ finite } \delta(\varrho_j, \tau_l) \stackrel{\text{def}}{=} \lim_h \mathcal{F}_{\gamma_h \omega_h}(v_h, \mu_h, \alpha_h, \beta_h, B_{\varrho_j}^{\tau_l}) \quad \forall j, l; \delta(\varrho_j, \tau_l) \in (0, 1).$$

Since $\varrho_j, \tau_l \rightarrow \delta(\varrho_j, \tau_l)$ is monotone non decreasing with respect to inclusion of lunulae, there is a (unique) monotone non decreasing with respect to inclusion and one-side continuous with respect to exterior approximation extension defined for all lunulae in the two parameters family, defined as follows:

$$\delta(\varrho, \tau) = \inf_{j, l} \{ \delta(\varrho_j, \tau_l) : \varrho_j > \varrho, \tau_l < \tau \}.$$

obviously:

$\forall \tau, \varrho \rightarrow \delta(\varrho, \tau)$ is right-continuous everywhere and continuous up to a countable set,

$\forall \varrho, \tau \rightarrow \delta(\varrho, \tau)$ is left-continuous everywhere and continuous up to a countable set.

Since separate continuity together with monotonicity entail continuity in 2 variables, we get

$\forall \tau, \varrho \rightarrow \delta(\varrho, \tau)$ is continuous for a.e. ϱ , $\forall \varrho, \tau \rightarrow \delta(\varrho, \tau)$ is continuous for a.e. τ

say, δ is continuous with respect to τ, ϱ almost everywhere in $0 < \tau < \varrho < r$.

Hence by monotonicity of $\mathcal{F}_{\gamma_h \omega_h}(v_h, \mu_h, \alpha_h, \beta_h, \cdot)$ with respect to inclusion of sets and the same monotonicity property of δ , together with the coincidence in a dense set of $\delta(\varrho, \tau)$ with the limit of $\mathcal{F}_{\gamma_h \omega_h}(v_h, \mu_h, \alpha_h, \beta_h, B_\varrho^\tau)$ we get the existence of such limit almost everywhere and its coincidence with δ almost everywhere.

Estimate (6.59) entails $\mathcal{F}_{\gamma_h \omega_h}(v_h, \mu_h, \alpha_h, \beta_h, B_\varrho^\tau) \leq 1$, hence $\delta(\varrho, \tau) \leq 1$, for all $\varrho, \tau, 0 \leq \tau \leq \varrho \leq r$. Hence also the estimate in (iii) of Theorem 6.1 holds true.

Now we prove (v): by (6.53) $\lambda_h \geq \varepsilon_h$, then by density upper bound in Theorem 2.10, and (6.47):

$$(6.63) \quad 0 < \mu_h \leq \frac{\varrho_h}{\lambda_h} \leq \frac{\varrho_h}{\varepsilon_h} \leq \varepsilon_h^{k-1} \quad \text{for large } h,$$

hence $\lim_h \mu_h = 0$; moreover, by (6.50), (6.51) and (6.56), changing variables, using Hölder inequality and (6.47) we find for large h :

$$\begin{aligned}
(6.64) \quad & \mu_h \int_{B_1} |\gamma_h - \omega_h|^q d\mathbf{x} = \\
& = \frac{\mu_h}{\varrho_h^2} \frac{1}{(\lambda_h \varrho_h^3)^{q/2}} \int_{B_{\varrho_h}(\mathbf{x}_h)} |g_h(\mathbf{x}_h) - w_h(\mathbf{x}_h) \pm (\varrho_h Dw_h(\mathbf{x}_h) \cdot \mathbf{y} - w_h(\mathbf{x}_h))|^q d\mathbf{y} \leq \\
& = \frac{\mu_h}{\varrho_h^2} \frac{1}{(\lambda_h \varrho_h^3)^{q/2}} \int_{B_{\varrho_h}(\mathbf{x}_h)} |g_h(\mathbf{x}_h) - w_h(\mathbf{x}_h)|^q d\mathbf{y} \leq \\
& \leq 2^{q-1} \mu \varrho_h^{-1} \lambda_h^{-1} \left(\int_{B_{\varrho_h}(\mathbf{x}_h)} |g_h|^q d\mathbf{y} + \int_{B_{\varrho_h}(\mathbf{x}_h)} |w_h|^q d\mathbf{y} \right) \leq \\
& \leq 2^{q-1} \mu \varrho_h^{-1} \varepsilon_h^{-1} \left(\int_{B_{\varrho_h}(\mathbf{x}_h)} |g_h|^{2q} d\mathbf{y} \right)^{1/2} \sqrt{\pi} \varrho_h + 2^{q-1} \mu \varrho_h^{-1} \varepsilon_h^{-1} C^q \pi \varrho_h^2 \leq \\
& \leq 2^{q-1} \mu \left(\sqrt{\pi} \varepsilon_h^{k/2-1} + \pi C^q \varepsilon_h^{k-1} \right) \leq 2^q \mu \sqrt{\pi} \varepsilon_h^{k/2-1}
\end{aligned}$$

We know $\mu_h \int_{B_1} |\omega_h|^q \rightarrow 0$ as $h \rightarrow \infty$ by the first statements in (6.54), (6.55). Hence (6.64) entails $\lim_h \mu_h \int_{B_1} |\gamma_h|^q d\mathbf{x} = 0$.

By Theorem 6.1, v_∞ is bi-harmonic in $B_1^+(\mathbf{0})$, $v_\infty = 0$ in $B_1^-(\mathbf{0})$ by (6.62) and

$$(6.65) \quad \int_{B_\varrho(\mathbf{0})} |D^2 v_\infty|^2 d\mathbf{x} = \int_{B_\varrho^+(\mathbf{0})} |D^2 v_\infty|^2 d\mathbf{x} \quad \varrho \in (0, 1),$$

hence, since $v_\infty \in H^2(B_1)$, (6.65) holds true also for $\varrho = 1$.

Since $v_\infty = 0$ in $B_1^-(\mathbf{0})$, traces continuity in H^2 entails $v_\infty = \partial v_\infty / \partial y = 0$ in $B_1(\mathbf{0}) \cap \{y = 0\}$. By (5.1) of Theorem 5.1 and (6.65) we get

$$\begin{aligned}
(6.66) \quad & \int_{B_\eta(\mathbf{0})} |D^2 v_\infty|^2 d\mathbf{x} = \int_{B_\eta^+(\mathbf{0})} |D^2 v_\infty|^2 d\mathbf{x} \leq \\
& \leq \eta^2 \int_{B_1^+(\mathbf{0})} |D^2 v_\infty|^2 d\mathbf{x} = \eta^2 \int_{B_1(\mathbf{0})} |D^2 v_\infty|^2 d\mathbf{x}.
\end{aligned}$$

Therefore, by exploiting (iii), (6.3) of Blow-up Theorem 6.1 and (6.66)

$$(6.67) \quad \limsup_h \mathcal{F}_{\gamma_h, \omega_h}(v_h, \mu_h, \alpha_h, \beta_h, B_\eta(\mathbf{0})) \leq \eta^2 \int_{B_1(\mathbf{0})} |D^2 v_\infty|^2 d\mathbf{x}$$

whereas, by (6.61),

$$\begin{aligned}
(6.68) \quad & \lim_h \mathcal{F}_{\gamma_h, \omega_h}(v_h, \mu_h, \alpha_h, \beta_h, B_\eta(\mathbf{0})) \geq \\
& \geq \eta^{2-\sigma} \lim_h \mathcal{F}_{\gamma_h, \omega_h}(v_h, \mu_h, \alpha_h, \beta_h, B_1(\mathbf{0})) = \eta^{2-\sigma} \int_{B_1(\mathbf{0})} |D^2 v_\infty|^2 d\mathbf{x}
\end{aligned}$$

contradicting the assumption on η and σ . \square

Theorem 6.4. (Decay of functional \mathcal{E} at boundary points) Assume (1.3), (1.4), (1.5), (1.6), (1.7), (1.8), (1.9), (1.10) and (1.11). Then,

$$(6.69) \quad \forall k > 2, \forall \eta, \sigma \in (0, 1), \quad \exists \tilde{\varepsilon} > 0, \exists \tilde{\vartheta} > 0 \quad \text{such that}$$

for any $\varepsilon \in (0, \tilde{\varepsilon}]$, any $\mathbf{x} \in \partial\Omega \setminus (T_0 \cup T_1 \cup M)$, any u an $\overline{\Omega_{\varphi+}}$ local minimizer of $\mathcal{E}(\cdot, \alpha, \beta, \overline{\Omega_{\varphi+}})$, any ϱ s.t. $B_\varrho(\mathbf{x}) \subset \tilde{\Omega}$ and (6.39), $0 < \varrho \leq (\varepsilon^k \wedge \bar{\varrho} \wedge (c_0 \vee 1)^{-1})$ and

$$(6.70) \quad \alpha \mathcal{H}^1(S_u \cap \overline{\Omega_{\varphi+}}) + \beta \mathcal{H}^1((S_{\nabla u} \setminus S_u) \cap \overline{\Omega_{\varphi+}}) < \varepsilon \varrho,$$

we have

$$(6.71) \quad \mathcal{E}(u, B_{\eta\varrho}(\mathbf{x})) \leq \eta^{2-\sigma} \max \left\{ \mathcal{E}(u, B_\varrho(\mathbf{x})), \varrho^2 \tilde{\vartheta} \left((\text{Lip}(\varphi'))^2 + (\text{Lip}(Dw))^2 \right) \right\}.$$

Proof. Straightforward consequence of Theorem 6.3 □

7. PROOF OF MAIN RESULTS

In this section we prove main results: say Theorems 1.1, 1.2, 1.3,

Proof of Theorem 1.2 - Assume $v \in GSBV^2(\tilde{\Omega}) \cap L^q(\tilde{\Omega})$ minimizes \mathcal{F} among $v \in GSBV^2(\tilde{\Omega}) \cap L^q(\tilde{\Omega})$ s.t. $v = w$ a.e. in $\tilde{\Omega} \setminus \Omega$. The existence of such v is proven by Theorem 2.1 and Remark 2.2.

First of all we notice that if $B \subset \tilde{\Omega}$ is an open ball and $\mathcal{H}^1(B \cap (S_v \cup S_{\nabla v})) = 0$ then v is smooth in B since $v \in H^2(B)$ so that by standard regularity theory we get $\tilde{v} \in C^2(B \cap \tilde{\Omega})$ (see [24]).

So we deduce $\tilde{v} \in C^2(\tilde{\Omega} \setminus \overline{(S_v \cup S_{\nabla v})})$.

Now we evaluate $\mathcal{H}^1(\tilde{\Omega} \cap (\overline{(S_v \cup S_{\nabla v})} \setminus (S_v \cup S_{\nabla v})))$. Set

$$(7.1) \quad \Omega_0 = \left\{ \mathbf{x} \in \tilde{\Omega} : \lim_{\varrho \rightarrow 0} \varrho^{-1} \mathcal{F}(v, B_\varrho(\mathbf{x})) = 0 \right\}$$

We are going to prove that Ω_0 is open.

Notice that $\Omega_0 \cap (\tilde{\Omega} \setminus \tilde{\Omega})$ is trivially open by assumptions (1.8), (1.9), (1.11) so that we have only to analyze points \mathbf{x} in $\tilde{\Omega} \cap \Omega_0$, and show that they are all in the interior part of Ω_0 .

The interior points $\mathbf{x} \in \Omega$ can be handled by applying Theorems 5.1, 5.4 in [9], to get

$$(7.2) \quad \mathcal{H}^1(\Omega \cap (\overline{(S_v \cup S_{\nabla v})} \setminus (S_v \cup S_{\nabla v}))) = 0,$$

so we have obtained the information that $\Omega_0 \cap \Omega$ is open, and thanks to (1.6) (1.7) we are left to show that all points $\mathbf{x} \in \Omega_0 \cap (\partial\Omega \setminus (T_0 \cup T_1 \cup M))$ are interior points of Ω_0 .

From now on we fix

$$(7.3) \quad \mathbf{x} \in \Omega_0 \cap (\partial\Omega \setminus (T_0 \cup T_1 \cup M)).$$

Let c_0 be the constant in the density upper bound Theorem 2.10. In order to apply Decay property of local minimizers (Theorem 6.3) fix $k > 2$, $\eta \in (0, 1)$, $\sigma \in (0, 1)$, and related constants $\varepsilon_0 = \varepsilon_0(\eta, \sigma, \alpha, \beta, \dots)$, $\vartheta_0 = \vartheta_0(\eta, \sigma, \alpha, \beta, \dots)$ whose existence

is warranted by Theorem 6.3 in the present paper and denote by $\tilde{\varepsilon}$ the constant ε_0 whose existence is warranted by Theorem 5.4 of [9], then set

$$(7.4) \quad L = (\text{Lip}(\varphi'))^2 + (\text{Lip}(Dw))^2 .$$

Choose $\eta' \in (0, \eta)$ s.t.

$$(7.5) \quad (\eta')^{1-\sigma} c_0 < \varepsilon_0 \wedge \tilde{\varepsilon} .$$

and denote by ε', ϑ' the related constants $\varepsilon' = \varepsilon'(\eta', \sigma, \alpha, \beta, \dots)$, $\vartheta' = \vartheta'(\eta', \sigma, \alpha, \beta, \dots)$ whose existence is warranted by Theorem 6.3.

Set $\varepsilon = \varepsilon_0 \wedge \tilde{\varepsilon} \wedge \varepsilon'$. Choose r s.t.

$$(7.6) \quad 0 < r^2 \vartheta_0 < \vartheta_0, \quad 0 < r < (\varepsilon^k \wedge \bar{\varrho} \wedge (c_0 \vee 1)^{-1}), \quad \int_{B_r(\mathbf{x})} |g|^{2q} d\mathbf{y} \leq \varepsilon^k,$$

$$(7.7) \quad r^2(\vartheta_0 \vee \vartheta') L < (c_0 \wedge \varepsilon) r, \quad B_r(\mathbf{x}) \cap (T_0 \cup T_1 \cup M) = \emptyset,$$

$$(7.8) \quad \mathcal{F}(v, B_r(\mathbf{x})) \leq \varepsilon \eta' r .$$

We claim that $B_{(1-\eta)r}(\mathbf{x}) \subset \Omega_0$. In fact, if $\mathbf{y} \in B_{(1-\eta)r}(\mathbf{x})$ there are 3 cases: if $\mathbf{y} \in \tilde{\Omega} \setminus \bar{\Omega}$ then v coincides with w which is $(C^2 \cap W^{2,\infty})(B_{\eta r}(\mathbf{y}))$ hence the functional has a nice decay, if $\mathbf{y} \in \Omega$ then we can repeat the argument in Section 6 of [9], if

$$(7.9) \quad \mathbf{y} \in \partial\Omega \cap B_{(1-\eta)r}(\mathbf{x})$$

additional analysis is required as follows.

In case (7.9) we have

$$(7.10) \quad \mathcal{F}(v, B_{\eta' r}(\mathbf{y})) \leq \mathcal{F}(v, B_r(\mathbf{x})) \leq \varepsilon \eta' r .$$

By (7.5)-(7.8), Theorems 6.3 with the choice $\varrho = \eta' r < r$, and density upper bound estimate (Theorem 2.10) we deduce

$$(7.11) \quad \begin{aligned} \mathcal{F}(v, B_{\eta' \varrho}(\mathbf{y})) &\leq (\eta')^{2-\sigma} (\mathcal{F}(v, B_{\varrho}(\mathbf{y})) \vee (\varrho^2 \vartheta' L)) \leq \\ &\leq (\eta')^{2-\sigma} ((c_0 \varrho) \vee ((c_0 \wedge \varepsilon) \varrho)) \leq \varepsilon_0 \eta' \varrho, \end{aligned}$$

$$(7.12) \quad \alpha \mathcal{H}^1(S_v \cap B_{\eta' \varrho}(\mathbf{y})) + \beta \mathcal{H}^1((S_{\nabla v} \setminus S_v) \cap B_{\eta' \varrho}(\mathbf{y})) < \varepsilon_0 \eta' \varrho$$

so that, by setting $\varrho' = \eta' \varrho$, Theorem 6.3 with the choice η entails

$$(7.13) \quad \begin{aligned} \mathcal{F}(v, B_{\eta \varrho'}(\mathbf{y})) &\leq \eta^{2-\sigma} (\mathcal{F}(v, B_{\varrho'}(\mathbf{y})) \vee ((\varrho')^2 \vartheta_0 L)) \\ &\leq \eta^{2-\sigma} (\mathcal{F}(v, B_{\varrho'}(\mathbf{y})) \vee (\varepsilon_0 \varrho')) . \end{aligned}$$

Inequalities (7.11),(7.13) together with $\varrho' = \eta' \varrho$ entail

$$(7.14) \quad \mathcal{F}(v, B_{\eta \varrho'}(\mathbf{y})) \leq \eta^{2-\sigma} \varepsilon_0 \varrho' .$$

In the same way we get: for any $h \in \mathbb{N}$

$$(7.15) \quad \mathcal{F}(v, B_{\eta^h \varrho'}(\mathbf{y})) \leq \eta^{h(2-\sigma)} \varepsilon_0 \varrho' .$$

entails

$$(7.16) \quad \begin{cases} \mathcal{F}(v, B_{\eta^{h+1} \varrho'}(\mathbf{y})) \leq \eta^{(h+1)(2-\sigma)} \varepsilon_0 \varrho' . \\ \alpha \mathcal{H}^1(S_v \cap B_{\eta^{h+1} \varrho'}(\mathbf{y})) + \beta \mathcal{H}^1((S_{\nabla v} \setminus S_v) \cap B_{\eta^{h+1} \varrho'}(\mathbf{y})) < \varepsilon_0 \eta^{h+1} \varrho' . \end{cases}$$

Since (7.15) holds true for $h = 1$ due to (7.14), by induction we know that (7.16) holds true for any $h \in \mathbb{N}$. Then $\forall h = 1, 2, \dots$

$$\begin{aligned}
(7.17) \quad & \mathcal{F}(v, B_{\eta^{h+1}\varrho'}(\mathbf{y})) \leq \eta^{(h+1)(2-\sigma)} \varepsilon_0 \varrho' = \\
& = \eta^{h(2-\sigma)} \eta^{2-\sigma} \varepsilon_0 \varrho' \leq \\
& \leq \eta^{h(2-\sigma)} \varepsilon_0 \eta \varrho' = \\
& = \eta^{h(1-\sigma)} \varepsilon_0 (\eta^{h+1} \varrho')
\end{aligned}$$

For every t s.t. $0 < t < \eta^2 \varrho'$ there is $j \geq 3$ s.t. $\eta^j \varrho' \leq t \leq \eta^{j-1} \varrho'$, so that, by (7.17),

$$\begin{aligned}
(7.18) \quad & t^{-1} \mathcal{F}(v, B_t(\mathbf{y})) \leq t^{-1} \mathcal{F}(v, B_{\eta^{j-1}\varrho'}(\mathbf{y})) \leq \\
& \leq t^{-1} \eta^{(j-2)(1-\sigma)} \varepsilon_0 \eta^{j-1} \varrho' = \\
& = (t^{-1} \eta^j \varrho') \eta^{(j-2)(1-\sigma)-1} \varepsilon_0 \leq \\
& \leq \eta^{(j-2)(1-\sigma)-1} \varepsilon_0
\end{aligned}$$

and passing to the limit as $t \rightarrow 0_+$ (say $j \rightarrow +\infty$) we get $\mathbf{y} \in \Omega_0$.

By summarizing we have shown that Ω_0 is an open set.

Since $S_v \cup S_{\nabla v}$ is countably $(\mathcal{H}^1, 1)$ rectifiable, by Theorem 3.2.19 in [23] we get $\mathcal{H}^1((S_v \cup S_{\nabla v}) \cap \Omega_0) = 0$, $\nabla v = Dv$ in Ω_0 , $\nabla^2 v = D^2 v$ in Ω_0 , so that $\tilde{v} \in C^2(\Omega_0)$ and $(S_v \cup S_{\nabla v}) \cap \Omega_0 = \emptyset$. Since Ω_0 is open then $\Omega_0 \cap \overline{(S_v \cup S_{\nabla v})} = \emptyset$. By Lemma 3.3 we have

$$(7.19) \quad \mathcal{H}^1\left(\tilde{\Omega} \cap \left(\overline{(S_v \cup S_{\nabla v})} \setminus (S_v \cup S_{\nabla v})\right)\right) = 0.$$

By setting

$$(7.20) \quad K_0 = \overline{S_v} \setminus (S_{\nabla v} \setminus S_v), \quad K_1 = \overline{S_{\nabla v}} \setminus S_v,$$

thanks to (1.6), (1.7), (1.8), (1.11), (7.19), (7.20) we obtain

$$(7.21) \quad K_0 \cup K_1 \text{ is closed, } \mathcal{H}^1(K_0 \cap \tilde{\Omega}) = \mathcal{H}^1(\overline{S_v}), \quad \mathcal{H}^1(K_1 \cap \tilde{\Omega}) = \mathcal{H}^1(\overline{S_{\nabla v}} \setminus S_v),$$

hence $K_0 \cap \tilde{\Omega}$ and $K_1 \cap \tilde{\Omega}$ are $(\mathcal{H}^1, 1)$ rectifiable, moreover

$$(7.22) \quad F(K_0, K_1, \tilde{v}) = \mathcal{F}(v) = \min\{\mathcal{F}(z) : z \in GSBV^2(\tilde{\Omega}) \cap L^q(\tilde{\Omega}), z = w \tilde{\Omega} \setminus \Omega\}.$$

Then (by Lemma 3.2 and Remark 3.3 of [9]) we conclude that (K_0, K_1, \tilde{v}) is a minimizing triplet for F in the class of admissible triplets.

By (7.20), (7.21), (7.22) and trace properties of $GSBV^2$ functions, we can say that properties (1.12), (1.13), (1.14), (1.15) hold true for the minimizing triplet (K_0, K_1, \tilde{v}) obtained by partial regularity of a weak minimizer for \mathcal{F} .

Eventually, by Lemma 3.2 of [9], we get for any other minimizing triplet $(\mathfrak{K}_0, \mathfrak{K}_1, u)$ of F

$$(7.23) \quad S_u \subset \mathfrak{K}_0, \quad (S_{\nabla u} \setminus S_u) \subset (\mathfrak{K}_1 \setminus \mathfrak{K}_0)$$

$$(7.24) \quad \mathcal{F}(u) \leq F(\mathfrak{K}_0, \mathfrak{K}_1, u) = F(K_0, K_1, v);$$

assume by contradiction that inequality in (7.24) is strict, then (by Theorem 2.1) there is $z \in GSBV^2(\tilde{\Omega}) \cap L^q(\tilde{\Omega})$ s.t.

$$(7.25) \quad \mathcal{F}(z) = \min \mathcal{F} \leq \mathcal{F}(u)$$

and, by repeating on z the regularization procedure previously performed on v , we find a minimizing triplet $(\mathfrak{Z}_0, \mathfrak{Z}_1, \tilde{z})$ for F fulfilling

$$(7.26) \quad F(K_0, K_1, \tilde{v}) = F(\mathfrak{Z}_0, \mathfrak{Z}_1, \tilde{z}) = \mathcal{F}(z).$$

Relationships (7.25) and (7.26) together contradict (7.24) with strict inequality; so in (7.24) we must have equality, hence properties (1.12), (1.13), (1.14), (1.15) hold true also for $(\mathfrak{K}_0, \mathfrak{K}_1, u)$. \square

Proof of Theorem 1.1 - The thesis follows immediately by Theorem 1.2 by dropping the term $\mu \int_{\tilde{\Omega}} |v - g|^q dx$ and exploiting Theorem 6.4 instead of Theorem 6.3. \square

Proof of Theorem 1.3 - The thesis follows immediately by Theorem 1.2 with the choice $K = \overline{S_v \cup S_{\nabla v}}$ where v minimizes \mathcal{F} and taking into account the assumption $\alpha = \beta$. \square

REFERENCES

- [1] E.Almansi, *Sull'integrazione dell'equazione differenziale $\Delta^{2n} = 0$* , Ann. Mat. Pura Appl., **III**, (1899), 1-51
- [2] L. Ambrosio, N.Fusco, D.Pallara, *Functions of Bounded Variation and Free Discontinuity Problems*, Oxford Math. Mon., Oxford Univ.Press, Oxford, 2000.
- [3] L.Ambrosio, L.Faina & R.March, *Variational approximation of a second order free discontinuity problem in computer vision*, SIAM J. Math. Anal., **32**, (2001), 1171-1197.
- [4] T.Boccellari & F.Tomarelli, *Generic uniqueness of minimizer for 1D Blake & Zisserman functional*, to appear.
- [5] A.Blake & A.Zisserman, *Visual Reconstruction*,The MIT Press, Cambridge, 1987.
- [6] M.Carriero, A.Leaci, *Existence theorem for a Dirichlet problem with free discontinuity set*, Nonlinear Analysis Th. Meth Appl., **15**, n.7, (1990), 661-677.
- [7] M.Carriero, A.Leaci & F.Tomarelli: *Free gradient discontinuities*, in "Calculus of Variations, Homogeneization and Continuum Mechanics", (Marseille 1993), 131-147, Ser.Adv.Math Appl.Sci., **18**, World Sci. Publishing, River Edge, NJ, 1994.
- [8] M.Carriero, A.Leaci & F.Tomarelli, *A second order model in image segmentation: Blake & Zisserman functional*, in "Variational Methods for Discontinuous Structures" (Como, 1994), Progr. Nonlinear Differential Equations Appl. **25**, Birkhäuser, Basel, (1996) 57-72.
- [9] M.Carriero, A.Leaci & F.Tomarelli, *Strong minimizers of Blake & Zisserman functional*, Ann. Scuola Norm. Sup. Pisa Cl.Sci. (4), **25**, (1997), n.1-2, 257-285.
- [10] M.Carriero, A.Leaci & F.Tomarelli, *Density estimates and further properties of Blake & Zisserman functional*, in "From Convexity to Nonconvexity", R.Gilbert & Pardalos Eds., Nonconvex Optim. Appl., **55**, Kluwer Acad. Publ., Dordrecht (2001), 381-392.
- [11] M.Carriero, A.Leaci & F.Tomarelli, *Second order functionals for image segmentation*, in: Advanced Mathematical Methods in Measurement and Instrumentation (Como 1998), Esculapio, (2000), 169-179.
- [12] M.Carriero, A.Leaci & F.Tomarelli: *Necessary conditions for extremals of Blake & Zisserman functional*, C. R. Math. Acad. Sci. Paris, **334**, n.4,(2002) 343-348.
- [13] M.Carriero, A.Leaci & F.Tomarelli: *Local minimizers for a free gradient discontinuity problem in image segmentation*, in "Variational Methods for Discontinuous Structures", Progr. Nonlinear Differential Equations Appl., **51**, Birkhäuser, Basel, (2002), 67-80.
- [14] M.Carriero, A.Leaci & F.Tomarelli, *Calculus of Variations and image segmentation*, J. of Physiology, Paris, **97**, n.2-3, (2003), 343-353.

- [15] M.Carriero, A.Leaci & F.Tomarelli, *Second Order Variational Problems with Free Discontinuity and Free Gradient Discontinuity*, in: “Calculus of Variations: Topics from the Mathematical Heritage of E. De Giorgi”, Quad. Mat., **14**, Dept. Math., Seconda Univ. Napoli, Caserta, (2004), 135–186.
- [16] M.Carriero, A.Leaci & F.Tomarelli, *Euler equations for Blake & Zisserman functional*, Calc. Var. Partial Diff.Eq., **32**, 1 (2008) 81-110.
- [17] M.Carriero, A.Leaci & F.Tomarelli, *Candidate local minimizer of Blake & Zisserman functional*, submitted to Arch. Rat.Mech.Anal.
- [18] M.Carriero, A.Leaci & F.Tomarelli, *Uniform density estimates for Blake & Zisserman functional*, to appear.
- [19] E. De Giorgi, Free discontinuity problems in calculus of variations, in “Frontiers in Pure & Appl.Math.”, R.Dautray Ed., North-Holland, Amsterdam, (1991), 55–61.
- [20] E. De Giorgi, M.Carriero & A.Leaci, *Existence theorem for a minimum problem with free discontinuity set*, Arch. Rational Mech. Anal., **108**, (1989), 195–218.
- [21] R.J.Duffin, *Continuation of biharmonic functions by reflection*, Duke Math. J., **22**, (1955), 313-324.
- [22] L.C.Evans & R.F.Gariepy, Measure Theory and Fine Properties of Functions, Studies in Advanced Math., CRC Press, Boca Raton, 1992.
- [23] H.Federer, Geometric Measure Theory, Springer, Berlin, 1969.
- [24] M.Giaquinta, *Multiple Integrals in the Calculus of Variations and Nonlinear Elliptic Systems*, Ann. Math. Stud., Princeton U. P., 1983.
- [25] P.Grisvard, *Elliptic Problems in Nonsmooth Domains*, Pitman, London, 1985.
- [26] F.A.Lops, F.Maddalena & S.Solimini, *Hölder continuity conditions for the solvability of Dirichlet problems involving functionals with free discontinuities*, Ann. Inst. H. Poincaré Anal. Non Linéaire, **18**, (2001), n.6, 639-673.
- [27] P.A. Markovich, *Applied Partial Differential Equations: a Visual Approach* Springer, 2007.
- [28] J.M.Morel & S.Solimini, *Variational Models in Image Segmentation*, Progr. Nonlinear Differential Equations Appl., **14**, Birkhäuser, Basel, 1995.

Acknowledgements -

*Research partially supported by PRIN 2006 M.U.R
(Progetto “Problemi variazionali con scale multiple”)*

POLITECNICO DI MILANO, DIPARTIMENTO DI MATEMATICA "FRANCESCO BRIOSCHI", ITALY,
DIPARTIMENTO DI MATEMATICA "ENNIO DE GIORGI" UNIVERSITÀ DEL SALENTO, ITALY

E-mail address: `michele.carriero@unisalento.it`

E-mail address: `antonio.leaci@unisalento.it`

E-mail address: `franco.tomarelli@polimi.it`